

Great Lakes Coastal Wetland Monitoring Program

Semiannual Progress Report October 1, 2024 – March 31, 2025

Prepared for:

U.S. EPA GLNPO (G-17J) 77 W. Jackson Blvd. Chicago, IL 60604-3590
Contract/WA/Grant No./Project Identifier: GL-00E00612-0



Prepared by:

Dr. Donald G. Uzarski, Principal Investigator
CMU Institute for Great Lakes Research
Central Michigan University
Mount Pleasant, MI 48859

Dr. Valerie J. Brady, QA Manager
Natural Resources Research Institute
University of Minnesota Duluth
Duluth, MN 55811-1442

Dr. Matthew J. Cooper, QA Manager
Department of Biology
Grand Valley State University
Allendale, MI 49401

Table of Contents

INTRODUCTION	4
SUMMARY OF SAMPLING SCHEDULE	4
PROGRAM ORGANIZATION.....	5
PROGRAM TIMELINE.....	5
SITE SELECTION	7
ORIGINAL DATA ON GREAT LAKES COASTAL WETLAND LOCATIONS	7
SITE MANAGEMENT SYSTEM	7
2025 SITE SELECTION	12
<i>Site Management System Improvements</i>	<i>14</i>
TRAINING.....	15
CERTIFICATION.....	16
DOCUMENTATION AND RECORD.....	17
WEB-BASED DATA ENTRY SYSTEM	17
RESULTS-TO-DATE (2011-2024, WITH EXCEPTIONS NOTED)	20
BIOTIC COMMUNITIES AND CONDITIONS.....	26
WETLAND CONDITION.....	38
PUBLIC ACCESS WEBSITE.....	48
COASTAL WETLAND MAPPING TOOL.....	50
OUTREACH TO MANAGERS	53
TEAM REPORTS.....	55
WESTERN BASIN BIRD/ANURAN TEAM AT THE NATURAL RESOURCES RESEARCH INSTITUTE, UNIVERSITY OF MINNESOTA DULUTH	55
WESTERN BASIN FISH, INVERTEBRATE AND WATER QUALITY TEAM AT THE NATURAL RESOURCES RESEARCH INSTITUTE, UNIVERSITY OF MINNESOTA DULUTH.....	60
TEAM NAME: WESTERN BASIN VEGETATION TEAM AT UNIVERSITY OF WISCONSIN-SUPERIOR	68
US CENTRAL BASIN BIRD & ANURAN TEAM AT THE COFRIN CENTER FOR BIODIVERSITY, UNIVERSITY OF WISCONSIN-GREEN BAY.....	70
US CENTRAL BASIN, CENTRAL MICHIGAN UNIVERSITY (CMU), BIRD/ANURAN TEAM.....	75
US CENTRAL BASIN FISH, INVERTEBRATE AND WATER QUALITY TEAM	82
US CENTRAL BASIN VEGETATION TEAM.....	95
CANADIAN CENTRAL/EASTERN BASIN BIRD/ANURAN TEAM AT BIRDS CANADA, PORT ROWAN/LONG POINT, ONTARIO.....	105
CANADIAN CENTRAL BASIN FISH, INVERTEBRATE AND WATER QUALITY TEAM AT THE UNIVERSITY OF WINDSOR AND UNIVERSITY OF WISCONSIN RIVER FALLS	111
CANADIAN CENTRAL BASIN VEGETATION TEAM AT THE UNIVERSITY OF WINDSOR AND UNIVERSITY OF WISCONSIN RIVER FALLS	115
CANADIAN EASTERN BASIN FISH, INVERTEBRATE AND WATER QUALITY TEAM AT CANADIAN WILDLIFE SERVICE	119
CANADIAN EASTERN BASIN VEGETATION TEAM AT CANADIAN WILDLIFE SERVICE	123
US EASTERN BASIN BIRD AND ANURAN TEAM AT SUNY BROCKPORT	127
US EASTERN BASIN FISH, INVERTEBRATE, AND WATER QUALITY TEAM AT SUNY BROCKPORT.....	130

US EASTERN BASIN VEGETATION TEAM AT SUNY BROCKPORT	134
ASSESSMENT AND OVERSIGHT	138
DATA VERIFICATION	140
EXAMPLE WATER QUALITY QC INFORMATION	143
COMMUNICATION AMONG PERSONNEL	146
OVERALL	147
LEVERAGED BENEFITS OF PROJECT (2010 – 2023)	147
SPIN-OFF PROJECTS (CUMULATIVE SINCE 2010).....	148
SUPPORT FOR UN-AFFILIATED PROJECTS (2011 – 2023)	159
REQUESTS FOR ASSISTANCE COLLECTING MONITORING DATA (2011 – 2023)	160
STUDENT RESEARCH SUPPORT (2011 – 2023).....	162
<i>Graduate Research with Leveraged Funding:</i>	162
<i>Undergraduate Research with Leveraged Funding:</i>	164
<i>Graduate Research without Leveraged Funding:</i>	165
<i>Undergraduate Research without Leveraged Funding:</i>	167
JOBS CREATED/RETAINED (2020)	168
JOBS CREATED/RETAINED (CUMULATIVE SINCE 2011, LAST UPDATED 2020).....	169
PRESENTATIONS ABOUT THE COASTAL WETLAND MONITORING PROJECT (INCEPTION THROUGH 2023).....	170
PUBLICATIONS/MANUSCRIPTS (INCEPTION THROUGH 2023)	193
REFERENCES.....	199
APPENDIX	202

INTRODUCTION

Monitoring the biota of Great Lakes coastal wetlands began as a project funded under the Great Lakes Restoration Initiative on 10 September 2010. The project had the primary objective of implementing a standardized basin-wide coastal wetland monitoring program. Our first five years of sampling (2011-2015) set the baseline for future sampling years and showed the power of the datasets that can be used to inform decision-makers on coastal wetland conservation and restoration priorities throughout the Great Lakes basin. During round one, we 1) developed a database management system; 2) developed a standardized sample design with rotating panels of wetland sites to be sampled across years, accompanied by sampling protocols, QAPPs, and other methods documents; and 3) developed background documents on the indicators.

We have completed three five-year rounds of monitoring and this summer will be year 4 of the third five-year sampling round (2021-2025). This is our first full 5-year sampling round as a sampling program rather than a project. During the second round (2016-2020) we combated high water levels that made wetland sampling challenging and drowned out some wetlands. Fortunately, Great Lakes water levels have moderated for round 3. In addition, we continue to support wetland restoration projects by providing data, information, and context.


SUMMARY OF SAMPLING SCHEDULE

Our yearly sampling schedule proceeds in this manner: During the winter, PIs and crew chiefs meet to discuss issues, update each other on progress, and ensure that everyone is staying on track for QA/QC. Sites are selected by March using the on-line site selection database system, and field crew training takes place from March – June, depending on sampling type. Anuran sampling typically begins in late March/early April with bird sampling beginning in April or May, and finally vegetation, fish, macroinvertebrate, and water quality sampling begins in June. Sampling start dates are weather and temperature dependent. Phenology is followed across the basin so that the most southerly sites are sampled earlier than more northerly sites. In the fall and early winter, data are entered into the database, unknown fish and plants are identified, and macroinvertebrates are identified. The goal is to have all data entered and QC'd by March. Metrics and IBIs are calculated in late March in preparation for the spring report to US EPA GLNPO.

Full summaries of the first two 5-year rounds of sampling have been submitted to US EPA and are available at <http://www.greatlakeswetlands.org/Reports-Publications.vbhtml>.

PROGRAM ORGANIZATION

Figure 1 shows our current organization. Our project management team has not changed.



PROGRAM TIMELINE

The program timeline remains unchanged and we are on schedule (Table 1). During the next project period we will sample the sites selected for this summer. In addition, we have shifted to the new Site Management System, hosted on the CMU servers, which currently host the Data Management System. This will be our first field season using the new Site Management System.

Table 1. Timeline of tasks and deliverables for the Great Lakes Coastal Wetland Monitoring Program.

Tasks	2021				2022				2023				2024				2025				2026			
	W	Sp	Su	F	W	Sp	Su	F	W	Sp	Su	F	W	Sp	Su	F	W	Sp	Su	F	W	Sp	Su	F
Funding received			X																					
PI meeting	X				X				X				X				X				X			
Site selection system updated	X				X				X				X				X							
Site selection for summer		X			X				X				X				X							
Sampling permits acquired		X				X				X				X				X						
Field crew training		X	X			X	X			X	X			X	X			X	X					
Wetland sampling		X	X			X	X			X	X			X	X			X	X					
Mid-season QA/QC evaluations			X				X				X				X				X					
Sample processing & QC				X	X			X	X			X	X				X	X			X	X		
Data QC & upload to GLNPO					X	X			X	X			X	X			X	X			X	X	X	
Report to GLNPO		X		X		X		X		X		X		X		X		X		X		X		X
Re-code Site Management System								X	X															

Table 2. GLRI Action Plan II of Measure of Progress. Wetlands are sampled during the summer.

GLRI Action Plan II of Measure of Progress		Reporting Period (Oct 1, 2024 – March 31, 2025)		Project Status* (February 2021 – January 2026)	
		Number	Percent	Number	Percent
4.1.3	Number of Great Lakes coastal wetlands assessed for biotic condition	0	0%	718	80%

* (Not Started; Started; Paused; 25% Completed; 50% Completed; 75% Completed; 95% Completed; and 100% Completed)

SITE SELECTION

Year fifteen site selection was completed in March 2025. We have completed our 5-year sampling scheme twice (round 1: 2011-2015; round 2: 2016-2020) and completed the fourth year of round 3 sampling (2021-2025) through our list of Great Lakes coastal wetlands.

Differences in the site list between successive sampling rounds are most often associated with special benchmark sites or changes due to lake levels and our ability to access sites safely and with permission. Benchmark sites (sites of special interest for restoration or protection) can be sampled more than once in the five-year sampling rotation, may need to be sampled in a different year to accommodate restoration work and may be sites that were not on the original sampling list. The dramatic change in Great Lakes water levels has also affected what wetlands we are able to sample for which biota. The list of wetlands to be sampled this year (2025) was previously sampled in 2015 and 2020, with some differences due to benchmarks, safe access, water levels, and the fact that 2020 was the first year of Covid and teams could not sample many sites due to travel restrictions.

ORIGINAL DATA ON GREAT LAKES COASTAL WETLAND LOCATIONS

The GIS coverage used was a product of the Great Lakes Coastal Wetlands Consortium (GLCWC) and was downloaded from the Great Lakes Commission website on December 6, 2010. See <http://www.glc.org/wetlands/inventory.html> for details.

SITE MANAGEMENT SYSTEM (formerly called the Site Selection System), original system completed in 2011, now the new system is in full use for 2025.

Background

In 2011, a web-based database application was developed to facilitate site identification, stratified random site selection, and field crew coordination. This database is housed at NRRI and backed up routinely. It is also password-protected. Using this database, potential wetland polygons from the GLCWC GIS coverage were reviewed by PIs and those that were greater than four hectares, had herbaceous vegetation, had (or appeared to have) a lake connection navigable by fish, and were influenced by lake water levels were placed into the site selection random sampling rotation (Table 3). That is, these 1014 wetlands became our wetland sampling universe, with minor modifications and additions for benchmark sites, as previously described, and some sites being dropped due to lack of any crew ever being able to access them. See the

QAPP for a thorough description of site selection criteria. Note that the actual number of sampleable wetlands fluctuates year-to-year with lake level, continued human activity and safe access for crews. Based on the number of wetlands that proved to be sampleable thus far, we expect that the total number of sampleable wetlands will be around 900 in any given year; we sample roughly 180 of these (one fifth) per year.

Table 3. Counts, areas, and proportions of the 1014 Great Lakes coastal wetlands deemed sampleable in 2011 following Great Lakes Coastal Wetland Consortium protocols based on review of aerial photography. Area in hectares.

Country	Site count	Site percent	Site area	Area percent
Canada	386	38%	35,126	25%
US	628	62%	105,250	75%
Totals	1014		140,376	

This wetland coverage shows more wetlands in the US than in Canada, with an even greater percent of wetland area in the US (Table 3). We speculate that this is partly due to poor representation of Georgian Bay (Lake Huron) wetlands in the sampleable wetland database. This area is also losing wetlands rapidly due to a combination of glacial rebound and topography that limits the potential for coastal wetlands to migrate downslope during periods of low lake levels and to recover with rising water levels. Another component of this US/CA discrepancy is the lack of coastal wetlands along the Canadian shoreline of Lake Superior due to the rugged topography and geology. A final possibility is unequal loss of wetlands between the two countries, but this has not been investigated.


Strata

Geomorphic classes

Geomorphic classes (riverine, barrier-protected, and lacustrine) were determined for each site in the original coastal wetland GIS coverage. Many wetlands inevitably combine aspects of multiple classes, with an exposed coastal region transitioning into protected backwaters bisected by riverine elements. Wetlands were classified according to their predominant geomorphology. Note that we typically do not revisit or change the class originally assigned to a wetland during our 2011 initial site review process.

Regions

Existing ecoregions (Omernik 1987, Bailey and Cushwa 1981, CEC 1997) were examined for stratification of sites. None were found which stratified the Great Lakes' shoreline in a manner that captured a useful cross section of the physiographic gradients in the basin. To achieve the intended stratification of physiographic conditions, a simple regionalization was adopted that divided each lake into northern and southern components, with Lake Huron being split into three parts and Lake Superior being treated as a single region (Figure 2). The north-south splitting of Lake Michigan is common to all major ecoregion systems (Omernik / Bailey / CEC).



Panelization

Randomization

To create our stratified random wetland site sampling design, the first step was the assignment of selected sites from each of the project's 30 strata (10 regions x 3 geomorphic wetland types) to a random year or panel in the five-year rotating panel. Because the number of sites in some strata was quite low (in a few cases less than 5, more in the 5-20 range), simple random assignment would not produce the desired even distribution of sites within each strata over time.

Instead it was necessary to assign the first fifth of the sites within a stratum, defined by their pre-defined random ordering, to one year, and the next fifth to another year, etc. All sites were assigned to panels in 2011, prior to the first round of sampling.

In 2012, sites previously assigned to panels for sampling were assigned to sub-panels for re-sampling. The project's sampling design requires that 10% of sites are re-sampled the year after they were sampled based on their main panel designation to help determine interannual variability and the effects of changing water levels. This design requires five primary panels, A-E, one for each year of a five-year rotation, and ten sub-panels, α -j, for the 10% resample sites. If 10% of each panel's sites were simply randomly assigned to sub-panels in order α -j, sub-panel j would have a low count relative to other sub-panels. To avoid this, the order of sub-panels

was randomized for each panel during site-to-sub-panel assignment, as can be seen in the random distribution of the '20' and '21' values in Table 4.

For the first five-year cycle, sub-panel *a* was re-sampled in each following year, so the 20 sites in sub-panel *a* of panel *A* were candidates for re-sampling in 2012. The 20 sites in sub-panel *a* of panel *B* were candidates for re-sampling in 2013, and so on. In 2016, panel *A* was sampled for the second time, so the 21 sites in sub-panel *a* of panel *E* became the re-sample sites. This past summer (2024), panel *D* was sampled for the third time and the sites in sub-panel *c* of panel *C* comprised the re-sample sites. The total panel and sub-panel rotation covers 50 years.

Table 4. Sub-panel re-sampling, showing year of re-sampling for sub-panels *a-c*.

Main Panel	Subpanel										
	a	b	c	d	e	f	g	h	i	j	TOTAL
A: 2011 2016 2021	20/2012	21/2017	21/2022	20	21	20	21	21	21	21	207
B: 2012 2017 2022	20/2013	20/2018	20/2023	21	20	21	21	20	21	21	205
C: 2013 2018 2023	21/2014	21/2019	21/2024	21	21	20	21	21	21	21	209
D: 2014 2019 2024	22/2015	21/2020	21/2025	21	21	21	21	21	21	21	211
E: 2015 2020 2025	21/2016	20/2021	21/2026	21	21	21	20	21	21	21	208

Workflow states

Each site is assigned a particular 'workflow' status. During the field season, sites selected for sampling in the current year move through a series of sampling states in a logical order, as shown in Table 5. The *data_level* field is used for checking that all data have been received and their QC status. Users set the workflow state for sites in the web tool, although some states can also be updated by querying the various data entry databases. In 2020 we ran into the problem of being unable to sample sites because of the global pandemic, Covid-19. The site status code “could not sample” was added as a workflow state in the site selection list for crews to have more options to indicate problems sampling sites. “Could not access” is used to indicate when a crew cannot safely get to a site for some reason, while “could not sample” is used to indicate the inability to sample a site even though they can get to it (e.g., water is too deep for their sampling gear; for Covid, this would be things like no access onto tribal lands, etc.).

Team assignment

With sites assigned to years and randomly ordered within years, specific sites were then assigned to specific teams. Sites were assigned to teams initially based on expected zones of

logistic practicality, and the interface described in the ‘Site Status’ section is used to exchange sites between teams for efficiency and to better assure that distribution of effort matches each team’s sampling capacity.

Field maps


Multi-page PDF maps are generated for each site for field crews each year. The first page depicts the site using aerial imagery and a road overlay with the wetland site polygon boundary. The image also shows the location of the waypoint provided for navigation to the site via GPS. The second page indicates the site location on a road map at local and regional scales. The remaining pages list information from the database for the site, including site informational tags, team assignments, and the history of comments made on the site, including information from previous field crew visits intended to help future crews find boat launches and learn about any hazards a site poses.

Table 5. Workflow states for sites listed in the Site Status table within the web-based site selection system housed at NRRI. This system tracks site status for all taxonomic groups and teams for all sites to be sampled in any given year. Values have the following meanings: -1: site will not generate data, 0: site may or may not generate data, 1: site should generate data, 2: data received, 3: data QC’d.

Name	Description	Data_level
too many	Too far down randomly-ordered list, beyond sampling capacity for crews.	-1
Not sampling BM	Benchmark site that will not be sampled by a particular crew.	-1
listed	Place holder status; indicates status update needed.	0
web reject	Rejected based on regional knowledge or aerial imagery in web tool.	-1
will visit	Indicates site assignment to a team with intent to sample.	0
could not access site	Site proved impossible to access safely.	-1
could not sample	Added in 2020; indicates inability of crew to sample for some reason other than safety or lack of an appropriate wetland.	-1
visit reject	Visited in field, and rejected (no lake influence, no wetland present, etc.).	-1
data quarantined	Data quarantined due to non-compliance of sampling with QAPP/SOP	-1
will sample	Interim status indicating field visit confirmed sampleability, but sampling has not yet occurred.	1
sampled	Sampled, field work done.	1
entered	Data entered into database system.	2
checked	Data in database system QC-checked.	3


2025 SITE SELECTION

For 2025, 201 sites have been selected for sampling (Figure 3). Of these, 10 are benchmark sites. Another 18 sites are re-sample sites and 18 are pre-sample sites, which will be re-sample sites next year (2026). Benchmark, re-sample, and pre-sample sites are sorted to the top of the sampling list because they are the highest priority sites to be sampled. By sorting next year's resample sites to the top of the list, this helps ensure that most crews sample them, allowing more complete comparison of year-to-year variation when the sites are sampled again the next year. Because this is our third sampling round, crews are familiar with most of the sites on the 2025 site list.



Benchmark sites (Figure 4) are sites that are not on the site list, are special interest sites that were too far down the site list and risked not being sampled by all crews, or are sites that are considered a reference of some type and are being sampled more frequently. Sites that were

not on the site list typically are too small, disconnected from lake influence, or are not a wetland at this time, and thus do not fit the protocol. These sites are added back to the sampling list by request of researchers, agencies, or others who have specific interest in the sites. Many of these sites are scheduled for restoration, and the groups who will be restoring them need baseline data against which to determine restoration success. Each year, Coastal Wetland Monitoring (CWM) researchers get a number of requests to provide baseline data for restoration work.



We now have approximately 100 sites for which at least a portion of sampling is designated as “benchmark.” Of these sites, about 40 are to evaluate restoration efforts and about a dozen serve as reference sites for their area or for nearby restoration sites. The rest are more intensive monitoring sites at which the extra data will help provide long-term context and better ecological understanding of coastal wetlands.

Wetlands have a “clustered” distribution around the Great Lakes due to geological and topographic differences along the Great Lakes coastline. As has happened each sampling

season so far, several teams ended up with fewer sites than they had the capacity to sample, while other teams' assigned sites exceeded their sampling capacity. Within reason, teams with excess sampling capacity expanded their sampling boundaries to assist neighboring over-capacity teams to maximize the number of wetlands sampled. The Site Management System facilitates these exchanges.

Site Management System Improvements

The original Site Selection System had been in use for almost 15 years and had recently experienced multiple failures, with each fix becoming more tenuous due to old software and incompatibility issues with newer servers, image sources, and browser software. For the future integrity of the monitoring program, we completely re-constructed the system to become a Site Management System and move it to the servers that host the main CWMP website and Data Management System at Central Michigan University. The Site System problems and associated down time emphasized the critical importance of this system to the running of our program because it allows us to allocate sites correctly and efficiently across teams and the basin each sampling year in a manner that upholds the statistical design of our sampling program. It also allows us to track and note conditions and safety issues at each site as well as maintain notes on why sites are benchmarks and what we know about their benchmark and restoration status and progress.

The new Site Management System was thoroughly tested over the winter against the old Site Selection System and produced yearly site sampling lists that exactly matched the old system. We brought the new system online in February 2025 and used it to generate the 2025 site sampling list, establish benchmark sites, resample and presample sites, and allocate sites to teams across the basin. It did all of this correctly, with enhanced functionality and, most importantly, stability compared to the old system. The Site Management System is fully integrated into the main CWMP web application and database that support the Data Management System. The integrated system is currently housed on a dedicated CMU server. Going forward, the new, integrated Site Management System will provide opportunities for more effective and efficient data verification workflows because it is now possible to conduct real-time checks of the Data Management System data against the information in the Site Management System.

TRAINING

All personnel responsible for sampling invertebrates, fish, macrophytes, birds, anurans, and water quality received training and were certified prior to this sampling program beginning in 2011. During that first year, teams of experienced trainers held training workshops at several locations across the Great Lakes basin to ensure that all PIs and crews were trained in Coastal Wetland Monitoring methods. Now that PIs and crew leaders are experienced, field crew training is being handled by each PI at each regional location, with more experienced trainers providing assistance, including in-person training by the management team, as necessary when major personnel changes take place (e.g., new field crew leader, new PI). As is true every field season, all crew members still have to pass all training tests. Mid-season QC will also be conducted. As has become standard protocol, the trainers are always available via phone and email to answer any questions that arise during training sessions or during the field season.

The following is a synopsis of the training conducted by PIs each spring. See the individual team reports for information on how each team conducted crew training. Some crews are trained by the crew leader; some crews use primarily experienced personnel who have worked for the project for years and needed minimal retraining. In general, each PI or field crew leader trains all field personnel on meeting the data quality objectives for each element of the project; this includes reviewing the most current version of the QAPP, covering site verification procedures, providing hands-on training for each sampling protocol, and reviewing record-keeping and archiving requirements, data auditing procedures, and certification exams for each sampling protocol. All field crew members have to pass all training certifications before they are allowed to work unsupervised. Those who do not pass all training aspects are only allowed to work under the supervision of a crew leader who has passed all training certifications.

Training for bird and anuran field crews includes tests on anuran calls, bird vocalizations, and bird visual identification. These tests are based on an online system established at the University of Wisconsin, Green Bay – see <http://www.birdercertification.org/GreatLakesCoastal>. In addition, individuals are tested for proficiency in completing field sheets, and audio testing is done to ensure their hearing is within the normal ranges. Field training is also completed to ensure guidelines in the QAPP are followed: rules for site verification, safety issues including caution regarding insects (e.g., Lyme's disease), GPS and compass use, and record keeping.

Fish, macroinvertebrate, and water quality crews are trained on field and laboratory protocols. Field training included selecting appropriate sampling points within each site, setting fyke nets, identifying fish, sampling and sorting macroinvertebrates, and collecting water quality and

habitat covariate data. Laboratory training includes preparing water samples, titrating for alkalinity, and filtering for chlorophyll. Other training includes GPS use, safety and boating issues, field sheet completion, and GPS and records uploading. All crew members are required to be certified in each respective protocol prior to working independently.

Training for fish and invertebrate crews now includes specific instructions for sampling in deep water. These techniques were trialed in 2019 and found to work to allow sampling in at least somewhat deeper water than we have been sampling. Specifically, to sample macro-invertebrates in depths greater than 1 m, D-frame dip net handles can be extended and sampling can be done from the boat by moving around the boat and by allowing the boat to swing around one of its anchors. To set fyke nets in deeper water, the boat can be used to set the cod end of the net in deep water and the frame can be set underwater, using rock bag anchors to weight the cod end. These deep-set fyke net data are still considered experimental at this point and data are coded accordingly.

Vegetation crew training also includes both field and laboratory components. Crews are trained in field sheet completion, transect and point location and sampling, GPS use, and plant curation. Plant identification is tested following phenology through the first part of the field season. All crew members are certified in all required aspects of sampling before starting in the field unless supervised.

Training on data entry and data QC was provided by Valerie Brady and Terry Brown through a series of conference calls/webinars during the late summer, fall, and winter of 2011. All co-PIs and crew leaders responsible for data entry participated in these training sessions and each regional laboratory has been successfully inputting data for many years. Additional training on data entry, data uploading, and data QC was provided in 2016 with the implementation of the updated version of the data entry/data archiving system by Todd Redder at LimnoTech.

Training on data entry and QC continues via webinar as needed for new program staff and was done in both 2017 and 2018 as new staff joined the program. Additional training on data entry is now provided as needed.

CERTIFICATION

To be certified in a given protocol, individuals must pass a practical exam. Certification exams are conducted in the field in most cases, either during training workshops or during site visits early in the season. When necessary, field exams are supplemented with photographs (for fish and vegetation) or audio recordings (for bird and anuran calls). Passing a given exam certifies the individual to perform the respective sampling protocol(s). Since not every individual is

responsible for conducting every sampling protocol, crew members are only tested on the protocols for which they are responsible. Personnel who are not certified (e.g., part-time technicians, new students, volunteers) are not allowed to work independently nor to do any taxonomic identification except under the direct supervision of certified staff members. Certification criteria are listed in the project QAPP. For some criteria, demonstrated proficiency during field training workshops or during site visits is considered adequate for certification. Training and certification records for all participants are collected by regional team leaders and copied to Drs. Brady and Cooper (QC managers) and Uzarski (lead PI). Note that the training and certification procedures explained here are separate from the QA/QC evaluations explained in the following section. However, failure to meet project QA/QC standards requires participants to be re-trained and re-certified.

DOCUMENTATION AND RECORD

All site selection and sampling decisions and comments are archived in the site selection system (see “site selection”). These include comments and revisions made during the QC oversight process. These records are preserved in the new Site Management System. Regional team leaders archive copies of the testing and certification records of all field crew members. Summaries of these records are also archived with the QC managers (Brady and Cooper).

WEB-BASED DATA ENTRY SYSTEM

The CWMP uses a web-based data management system (DMS) that was originally developed by NRRRI in 2011 to collect field and laboratory data and then redeveloped by LimnoTech during 2015-16. The current web-based system uses Microsoft’s Active Server Pages .NET (ASP.NET) web application framework running on a Windows Server 2019 Datacenter and hosted on a virtual machine at Central Michigan University (CMU). The open source PostgreSQL Relational Database Management System (RDMS) with PostGIS spatial extensions is used to provide storage for all CWMP data, including both the DMS and the Site Management System, on the same Windows 2019 server that hosts the web application.

The CWMP database includes collections of related tables for each major taxonomic group, including vegetation, fish and macroinvertebrates, anurans, and birds. Separate data entry/editing forms are created for data entry based on database table schema information that is stored in a separate PostgreSQL schema. Data entry/editing forms are password-

protected and can only be accessed by users that have “Project Researcher” or “Admin” credentials associated with their CWMP user account and permissions for specific taxa group(s).

Specific features of note for the CWMP data management system include:

- Automated processes for individual users to request and confirm accounts;
- An account management page where a limited group of users with administrative privileges can approve and delete user accounts and change account settings as needed;
- Numerous validation rules employed to prevent incorrect or duplicate data entry on the various data entry/editing forms;
- Custom form elements to mirror field sheets (e.g. the vegetation transects data grid), which makes data entry more efficient and minimizes data entry errors;
- Domain-specific “helper” utilities, such as generation of fish length records based on fish count records;
- Dual-entry inconsistency highlighting for anuran and bird groups who use dual-entry for quality assurance;
- Tools for adding new taxa records or editing existing taxa records for the various taxonomic groups; and
- GPS waypoint file (*.gpx) uploading utilities and waypoint processing to support matching of geographic (latitude/longitude) coordinates to sampling points.

The CWMP data management system also provides separate webpages that allow researchers to download “raw” data for the various taxonomic groups as well as execute and download custom queries that are useful for supporting dataset review and QA/QC evaluations as data entry proceeds during and following each field season. Users from state management agencies are able to access the separate download pages for raw data and custom queries. Such organizations include GLNPO and its subcontractors and Michigan EGLE. Index of Biological Integrity (IBI) metrics are currently included as a download option based on static scores that reflect data collection through the 2024 field season. Over the past few years, a standalone .NET-based program has been developed and fully tested to automate the calculation of IBI metric scores for vegetation, invertebrates and fish on an annual (spring) schedule after data have been entered and gone through QA/QC.

Raw data downloads are available in both Microsoft (MS) Excel spreadsheet and MS Access database formats, while custom query results are available in spreadsheet format only. All available data/query export and download options are automatically regenerated every night, and users have the option of either downloading the last automated export or generating a new export that provides a snapshot of the database at the time the request is made (the former option is much faster). Currently, datasets for the major taxonomic groups must be downloaded individually; however, a comprehensive export of all pertinent data tables is generated in a single MS Access database file and provided to GLNPO on a bi-annual schedule in fall and spring of each program year.

In addition to providing CWMP researchers with data entry and download access, the CWMP data management team is providing ongoing technical support and guidance to GLNPO to support its internal management and application of the QA/QC'ed monitoring datasets. GLNPO, with support from subcontractors, maintains a separate, offline version of the CWMP monitoring database within the Microsoft Access relational database framework. In addition to serving as an offline version of the database, this version provides additional querying and reporting options to support GLNPO's specific objectives and needs under GLRI. CWMP data management support staff generate and provide to GLNPO and its contractors a "snapshot" of the master CWMP PostgreSQL database as a Microsoft Access database twice per year, corresponding to a spring and fall release schedule. This database release is then used by GLNPO and its contractors to update the master version of the Microsoft Access database used to support custom querying and reporting of the monitoring datasets.

A full backup of the CWMP PostgreSQL database is created each night at 3:00 AM Eastern time using a scheduled backup with the PostgreSQL Backup software application. Nightly database backups are automatically uploaded to a dedicated folder on LimnoTech's Sharefile system where they are maintained on a 30-day rolling basis. In the event of significant database corruption or other failure, a backup version can be restored within an hour with minimal data loss. The server that houses the DMS has also been configured to use CMU's Veeam Backup Solution. This backup solution provides end-to-end encryption including data at rest. Incremental backups are performed nightly and stored at secure locations (on premise and offsite). Nightly backup email reports are generated and sent to appropriate CMU IT staff for monitoring purposes. Incremental backups are kept indefinitely and restores can be performed for whole systems, volumes, folders and individual files upon request.

RESULTS-TO-DATE (2011-2024, WITH EXCEPTIONS NOTED)

A total of 176 wetlands were sampled in 2011, with 206 sampled in 2012, 201 in 2013, 216 in 2014, and 211 in 2015 our 5th and final summer of sampling for the first project round. Overall, 1010 Great Lakes coastal wetland sampling events were conducted in the first round of sampling (2011-2015; Tables 6 and 7), and we have completed sampling these wetlands a second time for the second complete round of coastal wetland assessment, 2016-2020. Note that this total number is not the same as the number of unique wetlands sampled because of temporal re-sampling events and benchmark sites that are sampled in more than one year per 5-year sampling round. For the second round of sampling, we sampled 192 wetlands in 2016, 209 wetlands in 2017, 192 wetlands in 2018, 211 wetlands in 2019, and 174 wetlands in 2020 (fewer wetlands sampled due to the global pandemic).

Round 3 (2021-2025) began summer 2021 with teams sampling 175 wetlands (again, fewer than in Round 2 due to the pandemic; Tables 6 and 7). In 2022 teams sampled 188 wetlands. In 2023, teams sampled 174 wetlands, and in 2024 teams sampled 180 wetlands (Tables 6 and 7, Figures 5 and 6).

In all years, more wetlands are sampled on the US side due to the uneven distribution of wetlands between the two countries. The wetlands on the US side also tend to be larger (see area percentages, Tables 6 and 7). When compared to the total number of wetlands targeted to be sampled by this project (Table 3), we are achieving our goals of sampling 20% of US wetlands per year, both by count and by area. However, each year 60-65% of total sites sampled are US coastal wetlands, with 75-80% of the wetland area sampled on the US side. Overall, we have sampled most of the large, surface-connected Great Lakes coastal emergent wetlands by count and by area. A few wetlands cannot currently be sampled due to a lack of safe access or a lack of permission to cross private lands.

Table 6. Counts, areas, and proportions of US Great Lakes coastal wetlands sampled in Round 1 (2011 – 2015), Round 2 (2016 – 2020) and Round 3 (2021 – 2025) sampling by the Coastal Wetland Monitoring Program. Percentages are of overall total sampled each year. Area in hectares.

US	Site count	Site %	Site area	Area %
Round 1 (2011 – 2015)				
2011	126	72%	22,008	87%
2012	124	60%	21,845	73%
2013	130	65%	18,939	73%
2014	144	67%	26,836	80%
2015	134	64%	26,681	73%
US total Round 1	658	65%	116,309	77%
Round 2: 2016 – 2020				
2016	129	67%	24,446	85%
2017	139	67%	30,703	80%
2018	125	65%	17,715	82%
2019	135	64%	30,281	80%
2020	119	69%	29,325	77%
US total Round 2	647	66%	132,470	82%
Round 3: 2021 – 2025				
2021	122	70%	24,734	85%
2022	128	68%	29,625	82%
2023	112	64%	18,648	82%
2024	117	63%	24,695	75%
US total Round 3	479	67%	97,702	81%

Table 7. Counts, areas, and proportions of Canadian Great Lakes coastal wetlands sampled in Round 1 (2011 – 2015), Round 2 (2016 – 2020) and Round 3 (2021 – 2025) sampling by the Coastal Wetland Monitoring Program. Percentages are of overall total sampled each year. Area in hectares.


Canada	Site count	Site %	Site area	Area %
Round 1: 2011 - 2015				
2011	50	28%	3,303	13%
2012	82	40%	7,917	27%
2013	71	35%	7,125	27%
2014	72	33%	6,781	20%
2015	77	36%	10,011	27%
CA total Round 1	352	35%	35,137	23%
Round 2: 2016 - 2020				
2016	63	33%	4,336	15%
2017	70	33%	7,801	20%
2018	67	35%	3,356	18%
2019	76	36%	7,746	20%
2020	55	32%	8,603	23%
CA total Round 2	331	34%	31,843	18%
Round 3: 2021 - 2025				
2021	53	30%	4,264	15%
2022	59	32%	6,637	18%
2023	62	36%	4,097	18%
2024	63	35%	8,137	25%
CA total Round 3	237	33%	23,135	19%
Overall Totals Round 1	1010		151,446	
Overall Totals Round 2	978		164,312	
Overall Totals Round 3	536		120,837	

Ability to sample sites depends not only on access but also on water levels. Teams were able to sample more sites in 2014 due to higher lake levels on Lakes Michigan and Huron, which allowed crews to access sites and areas that have been dry or inaccessible in previous years. By 2015 water depths in some coastal wetlands had become so deep that crews had difficulty


finding areas shallow enough to set fish nets in zones typically sampled for fish (cattail, bulrush, SAV, floating leaf, etc.). In 2017 Lake Ontario levels reached highs not seen in many decades. Water levels were again near historic highs in 2019 and 2020 and crews continued to report sampling challenges due to the high water, with coastal wetlands flooded out and only beginning to migrate upslope into areas that remain covered by terrestrial vegetation (shrubs, trees, etc.) or being blocked in this upslope migration by human land use or shoreline hardening. This highlights the difficulty of precisely determining the number of sampleable Great Lakes coastal wetlands in any given year, and the challenges crews face with rising and falling water levels.

In 2021, water levels had moderated slightly and crews reported fewer difficulties in sampling. This trend continued in 2022 and 2023, with some crews finding water levels low enough in some wetlands to impact sampling due to low water. The sites sampled in 2023 are shown in Figures 5 and 6 and are color coded by which taxonomic groups were sampled at the sites and by wetland types, respectively. Many sites were sampled for all taxonomic groups. Sites not sampled for birds and anurans typically were sites that were impossible to access safely, often related to private property access issues, or, during the pandemic, due to border closures. Most bird and anuran crews do not operate from boats since they need to arrive at sites in the dark or stay until well after dark. There are also a number of sites sampled only by bird and anuran crews because these crews can complete their site sampling more quickly and thus have the capacity to sample more sites than do the fish, macroinvertebrate, and vegetation crews. In both 2022 and 2023, bird and anuran crews faced a very cold, late spring across much of the region, compressing fieldwork into a shorter timeframe. Spring of 2024 was also slow to warm up, and in some areas of the Great Lakes was followed by an unseasonably cool and wet early summer.

Wetland types are not distributed evenly across the Great Lakes due to fetch, topography, and geology (Figure 6). Lacustrine wetlands occur in more sheltered areas of the Great Lakes within large bays or adjacent to islands. Barrier-protected wetlands occur along harsher stretches of coastline, particularly in sandy areas, although this is not always the case. Riverine wetlands are somewhat more evenly distributed around the Great Lakes. Low water levels in 2011-2013 and much higher water levels from 2014 – 2020 require that indicators be relatively robust to Great Lakes water level variations, or that data users are very cognizant of water level effects on indicators.



Benchmark sites are sites that are not on the site list, are special interest sites that were too far down the site list and risked not being sampled by all crews, or are sites that are considered a reference of some type and are being sampled more frequently. Sites that were not on the site list typically are too small, disconnected from lake influence, or are not a wetland at this time, and thus do not fit the protocol. These sites are added back to the sampling list by request of researchers, agencies, or others who have specific interest in the sites. Many of these sites are scheduled for restoration, and the groups who will be restoring them need baseline data against which to determine restoration success. Each year, Coastal Wetland Monitoring (CWM) researchers get a number of requests to provide baseline data for restoration work.



We now have about 100 sites that are or have been sampled as a “benchmark.” Of these, about 40 are to evaluate restoration efforts and about a dozen serve as reference sites for their area or for nearby restoration sites. The rest are more intensive monitoring sites at which the extra data will help provide long-term context, help us adjust indicators to be robust against water level fluctuations, and gain better ecological understanding of coastal wetlands. Almost all benchmark sites are in the US, with a few Canadian benchmark sites recently added.

Determining whether some of these benchmark sites would have been sampled at some point as part of the random site selection process is difficult because several of the exclusion conditions are not easy to assess without site visits. Our best estimate is that approximately 60% of the 17 benchmark sites from 2011 would have been sampled at some point, but they were marked “benchmark” to either sample them sooner (to get ahead of restoration work for baseline sampling) or so that they could be sampled more frequently. Thus, about 40% of 2011 benchmark sites were either added new because they were not (yet) wetlands, are small, or were missed in the wetland coverage, or would have been excluded for lack of connectivity.

This percentage decreased in 2012, with only 20% of benchmark sites being sites that were not already in the list of wetlands scheduled to be sampled. In 2013, 30% of benchmark sites were not on the list of random sites to be sampled by CWMP researchers in any year, and most were not on the list for the year 2013. For 2014, 26% of benchmark sites were not on the list of sampleable sites, and only 20% of these benchmark sites would have been sampled in 2014. These tend to be sites that are degraded former wetlands that no longer appear on any wetland coverage but for which restoration is a goal or, in a few cases, wetlands that are diked and the dike is being breached for restoration. There are a number of benchmark sites that are being sampled every year or every other year to collect extra data on these locations. At this point we are adding relatively few new sites as benchmarks each year (for 2023, only 2 new benchmarks were added; these are sites [7078, 7079] with major restorations planned for them). In 2024 we added a single new benchmark site (7080) in order to sample important wetlands on the upstream edge of the St. Louis River estuary that were missed in original site selection.

BIOTIC COMMUNITIES AND CONDITIONS (based on 2011-2024 data)

We can now compile good statistics on Great Lakes coastal wetland biota because we have sampled nearly 100% of the medium and large herbaceous coastal wetlands that have a surface water connection to the Great Lakes, are hydrologically influenced by lake levels, and can be safely accessed by crews in small boats. The following indicators and information are from data collected through 2024.

Wetlands average about 23-26 bird species; richness at high quality sites was as great as 54 bird species (Table 8). There are many fewer calling amphibian species (anurans) in the Great Lakes (8 total), and coastal wetlands averaged about 4 species per wetland, with some benchmark wetlands containing no anurans (Table 8). However, there were wetlands where 8 anuran species were heard over the three sampling dates.

Table 8. Bird and anuran species in wetlands; summary statistics by country. Data from 2011 through 2024, using only the latest year sampled for each wetland.

Country	Site count	Mean	Max	Min	St. Dev.
<i>Birds</i>					
Can.	254	26.9	55	9	10.2
U.S.	463	23.2	54	5	8.9
<i>Anurans</i>					
Can.	234	4.4	8	0	1.6
U.S.	407	4.1	8	0	1.4

Bird and anuran data in Great Lakes coastal wetlands by lake (Table 9) shows that wetlands on most lakes had an average number of bird species in the mid-twenties. The greatest number of bird species at a wetland occurred on lakes Erie, Huron and Ontario. These data include some benchmark sites, many of which are in need of or are undergoing restoration, so the minimum number of species found at a site can be quite low.

Calling anuran species counts show less variability among lakes simply because fewer of these species occur in Great Lakes coastal wetlands. Wetlands averaged about four calling anuran species regardless of lake (Table 9). Similarly, there was little variability by lake in maximum or minimum numbers of species. At some benchmark sites, and occasionally during unusually cold spring weather and/or at benchmark sites, no calling anurans were heard.

Table 9. Bird and anuran species found in Great Lakes coastal wetlands by lake. Mean, maximum, and minimum number of species per wetland for wetlands sampled from 2011 through 2024, using only data from the latest year sampled for each wetland.


Lake	Sites	Birds			Sites	Anurans		
		Mean	Max	Min		Mean	Max	Min
Erie	89	25.0	54	5	81	4.1	7	1
Huron	219	24.4	54	8	192	4.2	8	0
Michigan	130	23.9	50	5	116	4.0	7	0
Ontario	194	25.7	55	6	186	4.5	8	1
Superior	85	22.7	41	5	66	3.8	7	1

An average of 9 to 13 fish species were collected in Canadian and US Great Lakes coastal wetlands, respectively (Table 10). Again, these data include sites in need of restoration, and some had very few species. On the other hand, the wetlands with the highest richness had as many as 20 (CA) or 28 (US) fish species. The average number of non-native fish species per wetland was approximately one, though some wetlands had as many as 5. An encouraging sign is that there are wetlands in which no non-native fish species were caught in fyke nets, although some non-native fish are adept at net avoidance (e.g., common carp).

Table 10. Total fish species in wetlands, and non-native species; summary statistics by country for sites sampled from 2011 through 2024, using only data from the latest year sampled for each wetland.

Country	Sites	Mean	Max	Min	St. Dev.
<i>Overall</i>					
Can.	143	8.8	20	1	3.6
U.S.	253	12.8	28	0	4.7
<i>Non-natives</i>					
Can.	143	0.8	3	0	0.8
U.S.	253	1.0	5	0	1.1

From 2016-2020, we collected no non-native fish in 44% of Great Lakes coastal wetlands sampled, and we caught only one non-native fish species in 40% of Great Lakes coastal wetlands (Figure 7). We caught more than one non-native fish species in far fewer wetlands. It is important to note that the sampling effort at sites was limited to one night using passive capture nets, so these numbers are likely quite conservative, and wetlands where we did not catch non-native fish may actually harbor them.



Total fish species did not differ greatly by lake, averaging 10-13 species per wetland (Table 11). Lakes Erie and Huron had the most species of fish in a wetland, 28 species; the other lakes had a maximum of 22-26 species. Because sites in need of restoration are included, some of these sites had very few fish species or none at all. Wetlands averaged 1 non-native fish species captured. Having very few or no non-native fish is a positive and all lakes had some wetlands in which we caught no non-native fish. This result does not necessarily mean that these wetlands are free of non-natives. Our single-night net sets do not catch all fish species in wetlands, and some species are quite adept at avoiding passive capture gear. There are well-documented biases associated with each type of fish sampling gear. For example, active sampling gears (e.g., electrofishing) are better at capturing large active fish, but perform poorly at capturing smaller fish, forage fish, and young fish that are sampled well by our passive gear.

Table 11. Fish total species and non-native species found in Great Lakes coastal wetlands by lake. Mean, maximum, and minimum number of species per wetland. Data from 2011 through 2024, using only data from the most recent year sampled for each wetland.

Lake	Sites	Fish (Total)			Non-native		
		Mean	Max	Min	Mean	Max	Min
Erie	48	11.4	28	4	1.5	5	0
Huron	143	11.3	28	1	0.7	4	0
Michigan	56	12.2	26	0	1.1	4	0
Ontario	95	10.2	25	3	0.9	3	0
Superior	54	12.7	22	3	1.1	4	0

The average number of macroinvertebrate taxa (taxa richness) per site was about 36 (Table 12), but some wetlands had more than twice this number. Sites scheduled for restoration and other taxonomically poor wetlands had fewer taxa. On a more positive note, the average number of non-native invertebrate taxa found in coastal wetlands was less than 1, with a maximum of no more than 5 taxa (Table 12). Note that our one-time sampling may not be capturing all of the non-native taxa at wetland sites. In addition, some non-native macroinvertebrates are quite cryptic, resembling native taxa, and may not yet be recognized as invading the Great Lakes.

Table 12. Total macroinvertebrate taxa in Great Lakes coastal wetlands, and non-native species; summary statistics by country. Data from 2011 through 2023, using only data from the most recent year sampled for each wetland.

Country	Sites	Mean	Max	Min	St. Dev.
<i>Overall</i>					
Can.	181	36.7	71	18	10.1
U.S.	310	36.7	68	9	12.0
<i>Non-natives</i>					
Can.	181	0.7	4	0	0.9
U.S.	310	0.8	5	0	1.1


There is little variability among lakes in the mean number of macroinvertebrate taxa per wetland, with averages ranging from 31-42 taxa with Lake Erie having a lower average than the other lakes (Table 13). The maximum number of invertebrate taxa was lowest in Lake Erie wetlands (54) with the most invertebrate-rich wetlands in the other lakes having a maximum of 65-71 taxa. Wetlands with the fewest taxa are sites in need of restoration. Patterns are likely being driven by differences in habitat complexity, which may in part be due to the loss of wetland habitats. This has been documented in numerous peer-reviewed publications.

Table 13. Macroinvertebrate total taxa and non-native species found in Great Lakes coastal wetlands by lake. Mean, maximum, and minimum number of taxa per wetland. Data from 2011 through 2024, using only data from the most recent year sampled for each wetland.


Lake	Sites	Macroinvertebrates (Total)			Non-native		
		Mean	Max	Min	Mean	Max	Min
Erie	63	33.1	54	12	1.1	5	0
Huron	169	40.0	68	13	0.6	4	0
Michigan	77	35.2	66	9	1.1	4	0
Ontario	114	31.8	71	15	0.7	3	0
Superior	68	41.9	68	19	0.5	4	0

There is little variability among lakes in non-native taxa occurrence (Table 13). In each lake there were some wetlands in which we found no non-native macroinvertebrates. As noted above, however, this does not necessarily mean that these sites do not contain non-native macroinvertebrates.

We found zero non-native aquatic macroinvertebrates in 55% of Great Lakes coastal wetlands sampled from 2016-2020 (Figure 8), but in a handful of wetlands we found as many as 5 non-native invertebrate taxa.



In 2014 we realized that we are finding some non-native, invasive species in significantly more locations around the Great Lakes than are being reported on nonindigenous species tracking websites such as the USGS's Nonindigenous Aquatic Species (NAS) website (<http://nas.er.usgs.gov/>). Locations of aquatic macroinvertebrates are particularly under-reported. The best example of the difference is shown in Figures 9 and 10 for the faucet snail, *Bithynia tentaculata*. Figure 9 shows the range portrayed on the USGS website for this snail before we reported our findings. Figure 10 shows the locations where our crew found this snail. Finally, Figure 11 shows the USGS website map after it was updated with our crews' reported findings.



The faucet snail is of particular interest to USFWS and others because it carries parasites that can cause disease and die-offs of waterfowl. Because of this, we produced numerous press releases reporting our findings (collaborating universities produced their own press releases). The Associated Press ran the story and about 40 articles were generated in the news that we are aware of. See Appendix for a mock-up of our press release and a list of articles that ran based on this press release.

One reason that we were able to increase the geographic range and total number of known locations occupied by faucet snails is the limited number of ecological surveys occurring in the Great Lakes coastal zone. Furthermore, those surveys that do exist tend to be at a much smaller scale than ours and sample wetlands using methods that do not detect invasive species with the precision of our program.

In collaboration with the Great Lakes Environmental Indicators project and researchers at the USEPA Mid-Continent Ecology Division in Duluth and at the University of Wisconsin Superior, a note was published in the Journal of Great Lakes Research about the spread of *Bithynia* in Lake Superior (Trebitz *et al.* 2015).





Figure 10. Locations of *Bithynia tentaculata* found by CWM crews, 2011 - 2013.

We also routinely provide data on other non-native macroinvertebrates, fish, and aquatic vegetation to Great Lakes databases and websites that track this information.

On average, there were 42-43 macrophyte species per wetland (Table 14) with a maximum number of nearly 100 species at exceptionally diverse sites. Some sites were quite depauperate in plant taxa (some having none), particularly in highly impacted areas that were no longer wetlands but were sampled because they are designated for restoration and because of high water levels along higher energy coastlines.



Non-native vegetation is commonly found in Great Lakes coastal wetlands. We have updated our plant taxa lists to ensure that we are correctly coding all non-native macrophyte taxa, even those that are not currently considered invasive. This update changed the numbers of non-native species for many wetlands because in the past we had focused more on the non-natives that are invasive and are problematic in wetlands.

Coastal wetlands averaged 4-5 non-native species (Table 14). Some wetlands contained as many as 17 non-native macrophyte species, but there were wetlands in which no non-native plant species were found. It is unlikely that our sampling strategy would miss significant non-native plants invading a wetland. However, small patches of cryptic or small-stature non-natives could be missed. Invasive species are a particularly important issue for restoration work. Restoration groups often struggle to keep restored wetland sites from becoming dominated by invasive plant species.

Table 14. Total macrophyte species and non-native macrophytes in Great Lakes coastal wetlands; summary statistics by country. Data from 2011 through 2024, using only data from the most recent year sampled for each wetland.

Country	Site count	Mean	Max	Min	St. Dev.
<i>Overall</i>					
Can.	187	42.9	88	5	18.8
U.S.	354	43.2	95	0	18.7
<i>Non-native</i>					
Can.	187	5.5	17	0	3.5
U.S.	354	4.3	17	0	3.5

Lake Erie wetlands had the lowest mean number of macrophyte species (31, Table 15), with the other lakes' wetlands having higher mean numbers of species (34-48, Table 15). Average numbers of non-native species were highest in Lake Ontario (8 species) and lowest in Lake Superior wetlands (1 species; Table 15). Lake Superior had the lowest maximum number of non-native macrophytes in a wetland (8) and Lake Ontario had the highest maximum number with 17. There are wetlands on all lakes in which we did not detect invasive plants.

Table 15. Macrophyte total species and non-native species found in Great Lakes coastal wetlands by lake. Mean, maximum, and minimum number of species per wetland. Data from 2011 through 2024, using only data from the most recent year sampled for each wetland.

Lake	Sites	Macrophytes (Total)			Non-native		
		Mean	Max	Min	Mean	Max	Min
Erie	60	30.7	70	4	6.0	15	0
Huron	196	47.1	95	3	3.3	13	0
Michigan	89	42.3	82	4	4.5	11	0
Ontario	126	48.0	85	12	8.4	17	0
Superior	70	34.5	63	0	1.4	8	0

Our macrophyte data have reinforced our understanding of the numbers of coastal wetlands that contain non-native plant species (Figure 12, based on 2016-2020 data). Only 7% of 556 sampled wetlands lacked non-native species, leaving 93% with at least one. Sites were most commonly invaded by up to 7 non-native plant species and 13% of sites contained 8 or more non-native species. Detection of non-native species is more likely for plants than for organisms that are difficult to collect such as fish and other mobile fauna, but we may still be missing small patches of non-natives in some wetlands.




Figure 12. Number of Great Lakes coastal wetlands containing invasive plant species based on 2016 through 2020 data.

As an example for the state of Michigan, we also looked at wetlands with both invasive plants and plant species considered “at risk” (Figure 13). We found that there were a few wetlands at all levels of invasion that also had at-risk plant populations. This information will be useful to groups working to protect at-risk populations by identifying wetlands where invasive species threaten sensitive native species.





Figure 13. Number of state of Michigan Great Lakes coastal wetlands containing both invasive plant species and “at risk” plant species, based on 2011 through 2014 data.

We created a map of invasion status of Great Lakes coastal wetlands using all invasive species data we collected through 2014 for all taxonomic groups combined (Figure 14). Unfortunately, this shows that most sites have some level of invasion, even on Isle Royale. However, the more remote areas clearly have fewer invasives than the more populated areas and areas with relatively intense human use.



WETLAND CONDITION (based on 2011 – 2024 data unless otherwise noted)

In the fall of 2012 we began calculating metrics and IBIs for various taxa. We are evaluating coastal wetland condition using a variety of biota (wetland vegetation, aquatic macroinvertebrates, fish, birds, and anurans [calling amphibians]).

Macrophytic vegetation has been used for many years as an indicator of wetland condition (only large plants; algal species were not included). One very common and well-recognized indicator is the Floristic Quality Index (FQI); this evaluates the quality of a plant community using all of the plants at a site. Each species is given a Coefficient of Conservatism (C) score based on the level of disturbance that characterizes each plant species' habitat. A species found in only undisturbed, high quality sites will have a high C score (maximum 10), while a weedy species will have a low C score (minimum 0). We also give invasive and non-native species a rank of 0. These C scores have been determined for various areas of the country by plant

experts; we used the published C values for the midwest. The FQI is an average of all of the C scores of the species growing at a site, divided by the square root of the number of species. The CWM wetland vegetation index uses C scores for wetland species, among other metrics.

This IBI has been updated and adjusted multiple times since the start of the project, accounting for the shift in condition scores for some sites. The first adjustment was necessary to reflect changes in the taxonomic treatment of many marsh plants in the 2012 Michigan Flora and Flora of North America. In spring 2020, Dr. Dennis Albert, with assistance from Allison Kneisel, reviewed the data input file for the plants, looking at each individual species (taxa) on the list and observing how many records of each taxon were in the database. First, redundant entries were removed; some taxa had several synonyms in the database. The next step was to remove species that had no occurrences over 9 years of data collection; this eliminated 2082 species or 49.6% of the original species from the data input file.

A final step was to review the database for upland species or species that were outside of their accepted range. Some of these were clearly errors that resulted from the dropdown menu. For example, *Carex oligosperma*, a common northern wetland sedge, was recorded along several transects over several years in a Lake Superior wetland, but then *Carex oligocarpa*, an upland sedge immediately next to *C. oligosperma* on the dropdown list, was recorded at several points along a single transect. This was clearly a data recording error. Similar errors were identified for a handful of species. Another type of error that was identified and corrected in the database occurred when a species was noted that had a range north or south of the Great Lakes but appears very similar to a Great Lakes species so was identified in error. Similarly, cases were found in which an upland species was selected instead of the correct wetland species with very similar characteristics; this was also a rare situation involving less than 10 species.

Collectively, these revisions reduced the plant data input list from 4192 species to 1724 species, a reduction of 59%, which should both speed up and reduce errors in data input.

Allison Kneisel reviewed and modified the existing non-native species list. This process resulted in the addition of 9 species to the non-native species list. For computation of the IBI scores, many of the best-studied non-native species are used in computation of specific IBI metrics. For many of the species that were added to the non-native species list, there are few studies documenting what individual species are responding to, whether the response is to wetland dry down, increased nutrient loading, turbidity tolerance, or other factors.

In 2023 we are debuted a draft vegetation-based IBI; this IBI was originally developed by Dr. Dennis Albert during the early stage of Great Lakes-wide biotic sampling for the USEPA (Albert 2008) and is now updated (see Dybiec *et al.* 2020). The structure and many of the metrics of the

new IBI are shared with the original, but the new IBI has increased the number of metrics used and refined the metrics for the submergent zone. The original submergent zone metrics were difficult to compute.

Both the old and new IBIs were calculated by vegetation zone, making it possible to identify the source of degradation in a wetland. In many cases the impact of land or water use can result in the level of degradation in one zone being very different than that in other zones, and identifying the degraded zones can facilitate more effective restoration efforts. The advantage of the Dybiec *et al.* (2020) version is that the zonal scores are more easily accessible than in the original IBI, and the submergent zone metrics are much more dependable and easier to compute. The zonal scores in both IBIs are combined to create a site-wide score, and these site-wide scores are what are used in individual lake (Erie, Huron, Michigan, Ontario, and Superior) comparisons and long-term tracking of wetland quality change for the individual lakes and the entire Great Lakes.

The scores of the old and new IBIs are strongly correlated for the site-wide scores, with $R^2 = 0.65$ for the entire plant database between 2011-2022 (Figure 15), with a similar $R^2 = 0.63$ for the high-water years of 2021-2022 (Figure 16). It appears that the IBI scores of some of the most open lacustrine sites that had the highest IBI scores (5) with the original IBI, scored much lower with the new IBI, especially during high-water years of 2021 and 2022. Our interpretation is that the new IBI is providing a more effective evaluation of the submergent zone, a weakness in the original IBI.





Figure 15. Comparison of original vs. revised vegetation IBI (2011-2022).


Using the new IBI, the site-wide scores appear to be slightly lower for the most degraded sites (old IBI scores <2) and slightly higher for the less degraded sites (old IBI scores >2). This is likely the result of adding metrics based on specific taxa - *Carex* spp. for the Wet Meadow and Cyperaceae cover for the emergent zone - both taxonomic groups well represented in less degraded wetlands and often groups missing from highly degraded wetlands.



Lake-wide comparison of the old and new IBIs produce similar results. The order of lake-wide quality remains the same, with Lake Superior having the highest IBI scores, followed in order by Lake Huron, Lake Michigan, Lake Ontario, and Lake Erie.

The map (Figure 17) shows the distribution of Great Lakes coastal wetland vegetation index scores across the basin. Note that there are long stretches of Great Lakes coastline that do not have coastal wetlands due to topography and geology. Sites with low IBI scores are concentrated in the southern Great Lakes, where there are large amounts of both agriculture and urban development, and where water levels may be more tightly regulated (e.g., Lake Ontario), while sites with high IBI scores are concentrated in the northern Great Lakes. Even in


the north, an urban area like Duluth, MN may have high quality wetlands in protected sites and lower quality degraded wetlands in the lower reaches of estuaries (drowned river mouths) where there are legacy effects from the pre-Clean Water Act era, along with nutrient enrichment or heavy siltation from industrial development and/or sewage effluent. Benchmark sites in need of restoration will also have lower condition scores.




Another of the IBIs that was developed by the Great Lakes Coastal Wetlands Consortium uses the aquatic macroinvertebrates found in several of the most common vegetation types in Great Lakes coastal wetlands: sparse bulrush (*Schoenoplectus*), dense bulrush (*Schoenoplectus*), and wet meadow (multi-species) zones (Figure 18). In 2019 we had a major shift in the taxonomy of some invertebrates (primarily snails and mollusks) used in the calculation of some indicator metrics due to taxonomic updates and revisions. Thus, the invertebrate IBI map (Figure 18) in

this report should not be compared to the maps shown in previous reports. However, this IBI has been calculated for all sites with appropriate zones and invertebrate data for all years.

The lack of sites on lakes Erie and Ontario and southern Lake Michigan is due to either a lack of wetlands (southern Lake Michigan) or because these areas do not contain any of the three specific vegetation zones that GLCWC used to develop and test the invertebrate IBI. Many areas contain dense cattail stands (e.g., southern Green Bay, much of Lake Ontario), for which we do not yet have a published macroinvertebrate IBI. We are developing IBIs for additional vegetation zones to cover these sites, but these IBIs have not yet been validated so they are not included here.



Our fish IBI scores for wetland sites now contain bulrush, cattail, lily, or SAV zones (Figure 19). Because of the prevalence of these vegetation types in wetlands throughout the Great Lakes basin, this indicator provides more site scores than the macroinvertebrate indicator. Because these are updated and adjusted indicators, the map image in this report should not be compared to fish IBI map images in previous reports. However, all sites reporting fish data from zones applicable to the new fish IBIs are shown here, regardless of the year they were sampled.



To develop the most recent fish IBI, fish community metrics were evaluated against numerous indices of anthropogenic disturbance derived from measurements of water quality and surrounding land cover. Disturbance indices included individual land cover and water quality variables, principal components combining land cover and water quality variables, a previously published landscape-based index (SumRel; Danz *et al.* 2005), and a rank-based index combining land cover and water quality variables (SumRank; Uzarski *et al.* 2005). Multiple disturbance

indices were used to ensure that IBI metrics captured various dimensions of human disturbances.


We divided fish, water quality, and land cover data (2011-2015 data) into separate “development” and “testing” sets for metric identification/calibration and final IBI testing, respectively. Metric identification and IBI development generally followed previously established methods (e.g., Karr *et al.* 1981, USEPA 2002, Lyons 2012) in which 1) a large set of candidate metrics was calculated; 2) metrics were tested for response to anthropogenic disturbance or habitat quality; 3) metrics were screened for responses to anomalous catches of certain taxa, for adequate range of responses, and for highly redundant metrics; 4) scoring schemes were devised for each of the final metrics; 5) the final set of metrics was optimized to improve the fit of the IBI to anthropogenic disturbance gradients; and 6) the final IBI was validated against an independent data set.

Final IBIs were composed of 10-11 fish assemblage metrics for each of four vegetation types (bulrush [*Schoenoplectus* spp.], cattail [*Typha* spp.], water lily [*Brassenia*, *Nuphar*, *Nymphaea* spp.], and submersed aquatic vegetation [SAV, primarily *Myriophyllum* or *Ceratophyllum* spp.]). Scores of all IBIs correlated well with values of anthropogenic disturbance indices using the development and testing data sets. Correlations of IBIs to disturbance scores were also consistent among each of the five years. A manuscript describing development and testing of this IBI has been published (Cooper *et al.* 2018).

In 2024 we began using a new method for calculating the condition of Great Lakes coastal wetlands based on birds and anurans. The new method, called the Index of Biotic Condition (Howe *et al.* 2023), is qualitatively like our previous metric (Index of Ecological Condition) but is much simpler to calculate and therefore invites broader applications by state and local conservation agencies. We have back-calculated all point indices (IBC values), so our trend estimates are truly “apples-to-apples” comparisons. The IBC and IEC are highly correlated, and both are scaled to a range of 0 (poorest possible condition) to 10 (ideal condition). The Index of Biotic Condition (IBC), however, is more stable when few species are present and is more highly correlated with species richness. The IBC reaches a maximum value only when a full complement of indicator species is present at a site, generally leading to lower absolute values. In other words, using this method, biotic condition at Great Lakes wetlands based on birds (Figure 20) looks quite different than did these condition maps in previous reports.

Unlike the IEC method, the highest IBC value is achieved by an “ideal” species assemblage, which might not occur in the sampled data set (i.e., in any Great Lakes coastal wetland). The IBC and IEC use the same maximum likelihood method to quantify the sensitivity (biotic

response) of species to an explicit reference gradient defined by wetland size and the “human footprint” in the surrounding landscape and watershed. Unlike the IEC, the IBC assigns “weights” to different species based on parameters of the biotic response functions. These weights are applied to the simple arithmetic formula reflecting the number and environmental sensitivity (“quality”) of species present.



Coastal Wetland Monitoring field teams have recorded 13 species of anurans (2 toads and 11 frogs) since 2011, but 4 of these (northern [Blanchard’s] cricket frog, *Acris crepitans*; Fowler’s toad, *Anaxyrus fowleri*; mink frog, *Lithobates septentrionalis*; and pickerel frog, *Lithobates palustris*) are seldom observed. Cope’s gray treefrog (*Dryophytes chrysoscelis*) and eastern gray treefrog (*Dryophytes versicolor*) are sibling species that are difficult to differentiate in the field, so we combined records into a single taxon. We also did not separate geographically distinct species of chorus frogs, *Pseudacris*. IEC calculations for anurans therefore were based on 8 taxa

(American toad or Fowler's Toad, *Anaxyrus spp.*; gray treefrogs, *Dryophytes spp.*; bullfrog, *Lithobates catesbeianus*; northern leopard frog, *Lithobates pipiens*; green frog, *Lithobates clamitans*; wood frog, *Lithobates sylvaticus*; chorus frogs, *Pseudacris spp.*, and spring peeper, *Pseudacris crucifer*). A ninth category combines other less-common species such as pickerel frog and mink frog (*Lithobates spp.*). Wetland condition based on anuran communities as calculated by the new IBC method is shown in Figure 21.





Figure 31. Condition of coastal wetland calling anuran communities based on the IBC method and using data from the most recent visit to each wetland.

Finally, we have developed a water quality and land use indicator (Harrison *et al.* 2019). This indicator is based on landscape stressor data and water quality data collected from each aquatic plant morphotype (Figure 22).



PUBLIC ACCESS WEBSITE

The Coastal Wetlands Monitoring Program (CWMP) website provides efficient access to program information and summary results for coastal managers, agency personnel, and the interested public (Figure 23). As previously noted, the CWMP website was redeveloped and upgraded by LimnoTech and transitioned from an NRRI server to a permanent web hosting environment at Central Michigan University in spring 2016. The official launch of the new CWMP website occurred on April 26, 2016, including the public components of the website and data management tools for CWMP principal investigators and collaborators. Since that time, coastal managers and agency personnel have used the website's account management system

to request and obtain accounts that provide access to the wetland site mapping tool, which includes reporting of Index of Biotic Integrity (IBI) scores. CWMP researchers have also obtained user accounts that provide access to data upload, entry, editing, download, and mapping tools. LimnoTech is providing ongoing maintenance and support for the website, including modifying and enhancing the site as required to meet CWMP and GLNPO needs, as well as other end user needs.




Figure 23. Front page of the Great Lakes Coastal Wetland Monitoring public website, www.greatlakeswetlands.org.

The CWMP website provides a suite of interrelated webpages and associated tools that allow varying levels of access to results generated by the CWMP, depending on the user's data needs and affiliation. Webpages available on the site allow potential users to request an account and for site administrators to approve and manage access levels for individual accounts. Specific levels of access for the website are as follows:

- **Public** – this level of access does not require a user account and includes access to a basic version of the wetland mapping tool, as well as links to CWMP documents and contact information;

- **Site metrics (level 1)** – provides access to index of biological integrity (IBI) scores by wetland site via the coastal wetland mapping tool;
- **Agency/manager-basic (level 2)** - access to IBI scores and full species lists by wetland site via mapping tool;
- **CWMP scientists (level 4)** - access to data entry/editing tools (+ Level 3 capabilities); and
- **Admin** - access to all information and data included on the website plus administrative tools. A small team of CWMP principal investigators have been given “Admin” access and will handle approval of account requests and assignment of an access level (1-4).

The following sub-sections briefly describe the general site pages that are made available to all users (“Public” level) and the coastal wetland mapping tool features available to “Level 1” and “Level 2” users. User requests for CWMP datasets are handled through a formal process which involves the requestor submitting a letter detailing the request and providing assurances regarding maintaining the publication rights of the CWMP team. Additional pages and tools available to “Level 4”, and “Admin” users for exporting raw monitoring data, entering and editing raw data, and performing administrative tasks are not documented in detail in this report.

COASTAL WETLAND MAPPING TOOL

The enhanced CWMP website provides a new and updated version of the coastal wetland site mapping tool described in previous reports (<http://www.greatlakeswetlands.org/Map>). The basic version of the mapping tool, which is available at the “Public” access level, provides the following features and capabilities (Figure 24):

- Map navigation tools (panning, general zooming, zooming to a specific site etc.);
- Basemap layer control (selection of aerial vs. “ocean” basemaps);
- Display of centroids and polygons representing coastal wetlands that have been monitored thus far under the CWMP;
- Capability to style/symbolize wetland centroids based on: 1) geomorphic type (default view; Figure 24), or 2) year sampled (Figure 25); and
- Reporting of basic site attributes (site name, geomorphic type, latitude, longitude, and sampling years) and general monitoring observations for the site (e.g., hydrology, habitat, disturbances).

In addition to the features made available at the “Public” access level, users with “Level 1” (*Site Metrics*) access to the website can currently obtain information regarding IBI or IBC and a *Water Quality and Land Use Index*.




Figure 24. Coastal Wetland Mapping Tool – Public Version (geomorphic type view).




Figure 25. Coastal Wetland Mapping Tool – Public Version (sampling year view)

Wetland centroids can be symbolized based on IBI scores for a specific biological community, as well as based on geomorphic type and year sampled. For example, vegetation IBI scores calculated for individual sites can be displayed by selecting the “Vegetation IBI” option available in the “Style by:” pull-down menu (Figure 26). In addition, the actual IBI scores can be viewed by clicking on an individual wetland centroid.




Figure 26. Coastal Wetland Mapping Tool with IBI scores displayed.

Users with “Level 2” (Agency/Manager (basic)) access to the website are provided with the same visualization options described above for the “Public” and “Level 1” access levels, but also have the capability of viewing a complete listing of species observed at individual wetland sites. Species lists can be generated by clicking on the “Species List” link provided at the bottom of the “pop-up” summary of site attributes (Figure 27), and the information can then be viewed and copied and pasted to another document, if desired.

“Level 1” and “Level 2” users may also access the following tools that are available in the site mapping tool:

- **Wetland Site Report** – a tool that provides monitoring design information, monitoring observations, and the entire matrix of IBI/IEC/SumRank scores on an individual site basis.
- **Wetland Site Photos** – a photo viewer that allows users to review CWMP-approved digital photos taken during site sampling events.
- **Wetland Site Comparison** – a tool that allows users to select a geographic area of interest on the map and then generate a matrix comparing characteristics and IBI/IEC/SumRank scores across the selected sites.




Figure 27. Coastal Wetland Mapping Tool with wetland macrophyte IBI scores and species list displayed.

OUTREACH TO MANAGERS

There have been many improvements to the website which assist external users with accessing and understanding the results, in particular the site reports and photos. Michigan Department of Environment, Great Lakes and Energy (EGLE) and Central Michigan University hosted a workshop at the Michigan Wetlands Association annual meeting in Kalamazoo on September

12, 2023. The workshop focused on data collection methodology, data access, and data applications and was attended by 22 wetland management professionals.

In 2021, EGLE hired a new Wetland Monitoring and Coastal Wetland Analyst to fill the vacancy left by Anne Garwood. In transitioning into the position, Katie Fairchild met with many of the partners of the GLCWMP. Training included virtual meetings, introduction to the website and Coastal Wetlands Decision Support Tool, and a 2-day GLCWMP field training hosted by CMU. Katie will be leading the outreach efforts for EGLE going forward, including meeting planning, webinar scheduling and facilitation, and convening PIs and restoration partners to encourage application of the monitoring data in wetland restoration projects.

EGLE has also been encouraging restoration practitioners to use the GLCWMP data in project planning, goal setting, and development of adaptive management plans through Michigan's interagency Voluntary Wetland Restoration (VWR) Program. In the past year there have been multiple VWR projects undergoing regulatory review by EGLE where we requested that the practitioners identify if/how the GLCWMP data were used in planning or design of the project, and whether or not the project would be monitored as a benchmark site. Although there is still some uncertainty in how practitioners can or should use these data in project planning, there is momentum in the VWR Program to increase awareness and application of these results.

In 2019, a one-hour documentary on the GLCWMP was released on PBS. The documentary aired across the U.S. "Linking Land and Lakes: Protecting the Great Lakes' Coastal Wetlands" chronicled the work of all 15 universities and government agencies documenting our scientists collecting data to help restore and protect these ecosystems. The WCMU production team traveled the entire Great Lakes basin over 18 months covering 5,000 miles in Michigan, Wisconsin, Indiana, Illinois, New York, Ohio, Pennsylvania, and Ontario, Canada. More than 40 coastal wetland scientists shared their expertise in the documentary. This documentary aired on 275 PBS stations in 46 states, the Virgin Islands, and Washington D.C. beginning in July of 2020. It can be viewed at <https://www.pbs.org/video/linking-land-and-lakes-hdo22u/>

TEAM REPORTS

WESTERN BASIN BIRD/ANURAN TEAM AT THE NATURAL RESOURCES RESEARCH INSTITUTE, UNIVERSITY OF MINNESOTA DULUTH

Team Members

Team Members

- Dr. Annie Bracey, co-PI, avian ecologist (since 2012 as crew lead; 5 years since 2020 as co-PI)
- Josh Bednar, crew leader, wildlife ecologist (13 years since 2012)
- Isabel Dunn, summer field technician, UMD Water Resources graduate student (new)
- Amanda Tveite, summer field technician, UMD Integrated Biosciences graduate student (new)

Training

Annual training for anuran surveys will be virtual on April 15, 2025 and the bird survey training will be held at the Natural Resources Research Institute May 19-23, 2025. Training involves instructing crews on how to conduct standardized field surveys, on basic travel procedures, and on appropriate field safety measures. Individuals are trained to proficiently complete field sheets and audio testing is also completed to insure that their hearing is within the normal range. Rules for site verification, safety issues including caution regarding insects (e.g., Lyme's disease), GPS and compass use, and record keeping are also included in field training to insure that the guidelines in the QAPP are being followed. All individuals involved in conducting bird and anuran surveys will have taken and passed each of the following tests on 1) anuran calls, 2) bird vocalization, and 3) bird visual identification that are based on an online system established by the University of Wisconsin, Green Bay, prior to conducting surveys – see <http://www.birdercertification.org/GreatLakesCoastal>.

Challenges and Lessons Learned

There were no significant challenges that our team encountered this field season. Travel to and from Canada was allowed, so there were no issues with border crossing which we had experienced during Covid travel restrictions. Our primary challenge was site accessibility, whether sites were too far from one another to justify surveying (e.g., a single site >2+ hrs drive from all other sites) or island sites where access is challenging or restricted to daylight use (e.g. ferry service).

Site Visit List

In 2025, 46 sites were initially selected to be surveyed for birds and anurans by the western regional team. Most of these sites have been sampled in previous years for at least one taxonomic group. All of the sites selected for sampling were reviewed to determine whether they were deemed safe and accessible to field crews. Of these 46 sites, 27 will be surveyed for anurans and 30 will be surveyed for birds. The three sites that are only being surveyed for birds and not anurans are due to safety issues associated with accessing sites at night, primarily for sites that require boat access. The remaining sites that were excluded ($n = 16$) were primarily excluded due to access issues which included remote sites on islands which could not safely be reached by the bird and anuran team and other sites where access was not possible primarily due to private property or First Nations or Indigenous lands. Three sites were resamples from 2024 and seven sites are non-panel site (i.e. benchmark sites) occurring in the St. Louis River Estuary in Duluth, MN. These sites are being surveyed as a post-restoration assessment.

The location of the sites that are scheduled to be surveyed in 2025 by the bird and anuran western team stretches from the Duluth-Superior harbor area and extend north along Lake Superior to Thunder Bay, ON and Nipigon, ON as well as eastward along the south shore of Lake Superior as far as northeastern Lake Huron.

Panel Survey Results (from Fall report)

The data collected in 2024 by the western basin bird and anuran team were entered and error checked in the online data entry system and completed in September 2024.

Anurans: In 2024, nine species of anurans were recorded throughout our study sites, with 577 individuals and 52 full choruses counted (Table 1). The average number of species detected per wetland was four, with a minimum of two and a maximum of six. Spring peepers were the most abundant species detected in all wetlands sampled, accounting for 33% of the anuran observations and the majority of full chorus observations (Table 1). There were also large numbers of Green frog and Gray treefrog detections (Table 1). There was only one Chorus Frog detection which was similar to 2023. However, 24 Mink Frogs were detected, whereas there were none detected in 2023..

Table 16. List of anurans recorded during 2024 surveys. The number of individuals counted and the number of full choruses observed (i.e., number of individuals cannot be estimated) are provided for each species.

Species	Number of Individuals	Number of Observations (Full Chorus)
American toad (<i>Anaxyrus americanus</i>)	55	1
Blanchard's cricket frog (<i>Acris blanchardi</i>)	0	0
Bullfrog (<i>Lithobates catesbeianus</i>)	0	0
Chorus frog (western/ boreal – <i>Pseudoacris triseriata</i> & <i>P. maculatas</i>)	1	0
Green frog (<i>Lithobates clamitans</i>)	131	8
Gray treefrog (<i>Hyla versicolor</i>)	85	10
Mink frog (<i>Lithobates septentrionalis</i>)	24	0
Northern leopard frog (<i>Lithobates pipiens</i>)	41	1
Spring peeper (<i>Pseudoacris crucifer</i>)	192	32
Wood frog (<i>Lithobates sylvatica</i>)	47	0
Total	577	52

Birds: Birds were surveyed twice at each site between June 03 and July 09, 2024. A total of 86 identifiable species observations and 2,442 individual birds were recorded. The five most abundant species observed accounted for approximately 47% of all observations. These species, in order of decreasing abundance, were Red-winged Blackbird, Ring-billed Gull, Common Yellowthroat, Song Sparrow and Yellow Warbler.

Interesting bird observations: In the Western Great Lakes region there have been many observations of birds of special concern in the vicinity of the wetlands or using the wetland complexes in 2024 (Table 2). At least one bird of special concern was observed in 16 of the 20 wetland sites surveyed in 2024. There were relatively low numbers of detections for both Virginia and Sora rails which seems to be consistent with lower observations in recent years.

Table 17. List of birds of special interest recorded during 2024 surveys. The number of individuals observed is listed for each species.

Species	Number of Individuals
Sandhill Crane (<i>Grus canadensis</i>)	16
Pied-billed Grebe (<i>Podilymbus podiceps</i>)	4
American Bittern (<i>Botaurus lentiginosus</i>)	2
Virginia Rail (<i>Rallus limicola</i>)	5
Bald Eagle (<i>Haliaeetus leucocephalus</i>)	12
Common Loon (<i>Gavia immer</i>)	1
Sora Rail (<i>Porzana carolina</i>)	4
Great Blue Heron (<i>Ardea herodias</i>)	10
Green Heron (<i>Butorides virescens</i>)	5
Belted Kingfisher (<i>Megaceryle alcyon</i>)	5

Wetland Condition Observations and Results

The western basin bird and anuran team does not have any noteworthy observations to report regarding wetland condition of sites sampled in 2024.

Data Processing

Data entry and QAQC were completed by September 2024. All of the GPS coordinates associated with 2024 field sampling have been uploaded to the CWMP database. The physical data sheets from the point-count level vegetation surveys will be mailed to Doug Tozer at Bird Studies Canada for processing by November 2024.

All new and returning field observers will review the current QAPP and SOPs and any new observers will complete the online certification requirements (see above) prior to conducting field surveys. The supervising PI will conduct mid-season checks by visiting survey locations and verifying that proper protocols are being implemented. All data entry and QA for bird and anuran records will be completed (100%) by September 2025.

Mid-season QC Check Findings

In-person mid-season QC checks were conducted in 2024 to ensure protocols were being followed. The surveyors also reported to the PI daily during fieldwork. Surveyors also took pictures of sites where habitat was suspected to be inappropriate. These photos were then sent to the PI to verify whether the sites in question met sampling criteria or not. Surveyors also

described general field conditions and any issues associated with accessing sites. Data sheets were scanned and sent to the PI periodically throughout the field season to identify any potential issues with an individual's data collection methods. Surveyors were able to effectively communicate with the PI throughout the field season and therefore there were no QC issues that arose or needed to be addressed.

Additional Funding and Projects

Nothing to report

Other Collaboration Activities

Nothing to report

Other Data Requests

The NRRI avian ecology lab was awarded a grant through the USFWS Great Lakes Fish and Wildlife Restoration Act, where we will be documenting avian community responses to remediation and restoration activities in the St. Louis River Estuary (MN/WI) using historical and contemporary bird monitoring data to assess the current ecological health of restoration sites based on breeding bird communities. As part of this project, we will be utilizing some of the CWMP bird data that has been collected in the SLRE since 2011.

Related Student Research

There is one graduate student working on the bird and anuran team at NRRI. Isabel Dunn, who is a Water Resources graduate student at the University of Minnesota-Duluth. Isabel will use the bird data to evaluate avian biodiversity change at restored Great Lakes coastal wetland sites and determine how that relates to the amount of restoration on site, using both taxonomic and functional metrics. This includes determining the extent to which ecosystem services associated with bird communities changes at restored sites.

WESTERN BASIN FISH, INVERTEBRATE AND WATER QUALITY TEAM AT THE NATURAL RESOURCES RESEARCH INSTITUTE, UNIVERSITY OF MINNESOTA DULUTH

Team Members

- Dr. Valerie Brady, PI, aquatic invertebrate ecologist, QC manager (since 2011)
- Dr. Chris Filstrup, co-PI, limnologist (since 2019)
- Kristi Nixon, GIS specialist (since 2016)
- Kari Pierce, crew leader, fish, invertebrate, and water quality sampling (since 2014)
- Bob Hell, aquatic invertebrate taxonomist (since 2011)
- Holly Wellard Kelly, aquatic invertebrate taxonomist (since 2015)
- Paul Jeffrey, permanent field and lab crew member (since 2022)
- Two summer field techs, 2 returning (since 2023)

Training

The NRRI fish/invert/wq team will hold in-person safety and classroom project training in late May/early June 2024, as well as hands-on training for new summer technicians during their first site visit in Green Bay, WI (mid-June, 2025). The entire NRRI team (typically 6-8 participants) will be in attendance during training modules presented by permanent staff who have been working on the Coastal Wetland Monitoring Program for >5 years. Topics covered will include: field safety from environmental hazards, safe boating practices, approved scientific collection permits and responsibilities of the field teams to give prior notification to local fisheries managers and conservation officers before collecting fish from a wetland, Coastal Wetland Monitoring Program overview and introduction to Standard Operating Procedures and datasheets, GPS use and annual QC check, uploading GPS files to the program website, fish collection methods and identification, proper euthanasia and preservation methods for retained fish, water quality data and sample collection, post-collection processing of water samples (filtration and titration), daily calibration of water quality multiparameter instruments, invertebrate collection and field picking of samples, and vegetation identification and habitat quadrats. Hands-on field sampling and safety training will be led by experienced crew chiefs Kari Pierce, Bob Hell, Paul Jeffrey and Holly Wellard-Kelly who have all worked on GLCWMP for more than 3 years. During hands-on training the experienced NRRI crew chiefs (n=4) will guide summer technicians (n=2) on fish identification (with real fish rather than pictures), how to determine vegetation zones, vegetation identification, setting and pulling fyke nets, and which invertebrates to pick from trays (e.g., don't pick terrestrial insects, spiders, or large zooplankton).

Challenges and Lessons Learned (from summer 2024)

The 2024 field season was atypical because we dropped several sites that were selected for sampling. In total, we dropped five sites due to various reasons such as low water levels, no aquatic vegetation present, and access permission issues. Since we started with a slightly lower number of sites than normal years (n=18) and also had to drop multiple sites, PI Valerie Brady added two sites from the “Too Many” line for the crew to sample. Another challenge the crew faced was having a winch strap break on NRRI’s big Jon boat trailer. This occurred when pulling the boat out of the water on a steep and bumpy boat ramp in Superior, WI. The combination of tremendous strain on the strap from a heavy boat, the boat bouncing from a large bump on the boat ramp, and the boat being at an angle when it hit the bump likely contributed to the break. The crew was able to secure the boat to the trailer with two heavy duty ratchet straps until they returned to NRRI the same day. The winch strap was then replaced with a new one. The crew will prevent this from happening again by pulling the boat out of the water very slowly and ensuring the winch boat strap is not too tight.

Site Visit List

The 2025 CWMP field season for the NRRI team consists of 27 sites. Of these 27 sites, there are 17 regular sites, 3 pre-sample sites, 4 resample sites, and 3 benchmark sites. The sites are located in Michigan counties Delta (5), Baraga(1); Wisconsin counties Bayfield (4), Brown (2), Door(3), Douglas (4), Kewaunee (1), and Marinette (2); Minnesota counties Cook (1), St. Louis (1); and Thunder Bay District in Ontario, Canada.

Site 1680, Rowleys Bay Area Wetland, is located on the Mink River State Estuary and will require permission and coordination with the Nature Conservancy.

Site 974, Sand Point Wetland, will require special permission and coordination with the Keweenaw Bay Indian Community to access.

Site 1035, Chequamegon Wetland #2, will require special permission and coordination with the Bad River Band of Lake Superior Chippewa to access.

There are 3 Benchmark sites with 2 in Wisconsin and 1 in Minnesota. Site 1079, Hog Island Area Wetland, is a riverine wetland in Superior, WI. It was restored in 2005 as part of the AOC delisting effort, but seems to still have issues with sediment smelling like diesel fuel when disturbed during sampling in 2020. Site 1194, Gouge Park Pickle Ponds, is a barrier (protected) wetland in Superior, WI. The site was restored in 2023 as part of the AOC delisting effort. Site

7063, Spirit Lake, is a riverine wetland near Duluth, MN. It was restored in 2023 as part of the AOC delisting effort.

Panel Survey Results (from summer 2024)

Regular Panel Sites:

945 – First sampled on 7-17-2013 by the NRRI team. We re-sampled the site this year, with the last visit on 7-9-2024. Sampled SAV and Lily zones for fish, invertebrates, and water quality. Nets at this site (n=6) captured Rock Bass, Three-spined Stickleback, Blackchin Shiner, Brown Bullhead, Pumpkinseed, Bluntnose Minnow, Johnny Darter, and Golden Shiner. No invasive fish were detected. There were 3 native crayfish and 23 Painted Turtles as bycatch in fyke nets.

985 – First sampled on 7-14-2014 by the NRRI team. Last visit on 7-11-2024 and sampled Lily and Sparse Bulrush zones for fish, invertebrates, and water quality. Crew leader Holly Wellard-Kelly noted construction on the nearby shore, held back by retaining fabric. Muskrats and Beavers were also noted as a disturbance at this site. Nets at this site (n=6) captured Brown Bullhead, Bluegill, Pumpkinseed, Rock Bass, Blacknose Shiner, Yellow Perch, Golden Shiner, Johnny Darter, Smallmouth Bass, and Greater Redhorse. Invasive fish captured were Alewife (n=30). There were 6 Painted Turtles as bycatch in fyke nets.

1034 – First sampled on 9-23-2014 by the NRRI team. The NRRI team could not sample this site this year. This site and the boat launch to access this site are on Bad River Tribal lands. We tried to contact the Bad River Natural Resources Department multiple times for access permits to this site but never got a response. Without access permits we were not comfortable sampling this site.

1040 – First sampled on 7-17-2014 by the NRRI team. Last visit on 7-18-2024 and sampled a SAV zone for fish, invertebrates, and water quality, as well as a Typha zone for invertebrates and water quality. Crew leader Kari Pierce noted that the Typha zone was too shallow for fyke nets to be set. She also noted that a nearby homeowner added riprap to their shoreline, likely to help prevent erosion. Nets at this site (n=3) captured Pumpkinseed, White Sucker, Yellow Perch, Brown Bullhead, Rock Bass, Bluegill, Common Shiner, Johnny Darter, Spottail Shiner, Bluntnose Minnow, Smallmouth Bass, Fathead Minnow, Green Sunfish, Golden Shiner, and Hornyhead Chub. No invasive fish were detected.

1068 – First sampled on 7-18-2014 by the NRRI team. Last visit on 7-19-2024 and sampled a SAV zone for invertebrates and water quality. Crew leader Bob Hell noted that nearly the entire edge of the site is a floating bog, and the bottom consists of very deep, bottomless muck. The

site was too unsafe for the crew to exit the boat, and the only sampleable zone, SAV, was too deep and mucky to set nets. The team collected invertebrates and water quality from the boat.

1069 – First sampled on 7-19-2014 by the NRRI team. Last visit on 7-20-2024 and sampled a SAV zone for fish, invertebrates, and water quality. Crew leaders Bob Hell and Kari Pierce noted an extensive floating bog at this site that initially looked like Wet Meadow. It was too dangerous to walk in and mixed with woody shrubs, so the crew did not sample this zone. They also saw an otter at this site. Nets at this site (n=3) captured Pumpkinseed, Bluegill, Northern Pike, Yellow Perch, Blacknose Shiner, Brown Bullhead, Spottail Shiner, Golden Shiner, and Black Bullhead. Invasive fish captured were Three-spined Stickleback (n=6). There were 8 native crayfish and 5 Painted Turtles as bycatch in fyke nets.

1078 – First sampled on 7-23-2014 by the NRRI team. Last visit on 7-24-2024 and sampled Typha and SAV zones for fish, invertebrates, and water quality. The crew noted that they could only sample a portion of the site and were unable to sample within the river because there were no vegetation zones meeting the sampling criteria. They also observed taconite pellets on the bottom of the SAV zone they sampled, likely from the nearby shipping dock. Nets at this site (n=6) captured Rock Bass, Black Bullhead, Pumpkinseed, Bluegill, Johnny Darter, Northern Pike, Black Crappie, Yellow Perch, Mimic Shiner, Spottail Shiner, Common Shiner, Tadpole Madtom, Golden Shiner, and Bluntnose Minnow. Invasive fish captured were Round Goby (n=24), Tubenose Goby (n=4), and Eurasian Ruffe (n=2). Invasive Crayfish captured were Rusty Crayfish (n=1). There were 4 Painted Turtles and 1 native crayfish as bycatch in fyke nets.

1089 – First sampled on 8-21-2013 by the NRRI team. Last visit on 7-29-2024 and sampled SAV for fish, invertebrates, and water quality, as well as a Typha zone for invertebrates and water quality. Crew leader Kari Pierce noted that the Typha zone was too shallow for fyke nets and quickly turned into dry land as they moved towards the shore. Nets at this site (n=3) captured Walleye, Black Crappie, Spottail Shiner, Rock Bass, White Sucker, Yellow Perch, Troutperch, Black or Brown Bullhead YOY, Logperch, Emerald Shiner, Shorthead Redhorse, Smallmouth Bass, Johnny Darter, Tadpole Madtom, Northern Pike, Bluegill, and Pumpkinseed. Invasive fish captured were Eurasian Ruffe (n=2), Round Goby (n=39), and Tubenose Goby (n=7).

1114 – No Fish/Bug crew has sampled this site before, and the reason remains unknown, as there are no notes explaining why it wasn't sampled. The NRRI crew successfully sampled the site this year on 8-5-2024, sampling SAV for invertebrates and water quality. The crew couldn't access the site by boat because large boulders completely blocked boat access to the creek. Additionally, the site was too far (~1 mile) to carry a boat by hand. Crew leader Paul Jeffrey had

the crew walk-in to sample invertebrates and water quality since the creek was wadable. The rest of the site consisted of a sand beach, which did not meet the criteria for sampling.

1187 – First sampled on 7-21-2023 by the NRRI team. This year this site was a re-sample, last visit on 7-19-2024. Sampled a SAV zone for fish, invertebrates, and water quality, as well as a Typha zone for invertebrates and water quality. Crew leader Kari Pierce noted that the Typha zone formed a long, very thin line that became mixed with woody vegetation as it approached the shore, making it too narrow to fit fyke nets. Nets at this site (n=3) captured Northern Pike, Brown Bullhead, Yellow Perch, Emerald Shiner, White Sucker, Pumpkinseed, Golden Shiner, Bluegill, Bluntnose Minnow, and Rock Bass. No invasive fish were detected. There were 2 native crayfish as bycatch in fyke nets.

1192 – First sampled on 7-28-2014 by the NRRI team. Last visit on 7-24-2024 and sampled SAV for fish, invertebrates, and water quality, as well as a Lily zone for invertebrates and water quality. Crew leader Paul Jeffrey noted that the Lily zone did not meet the size criteria for setting fyke nets, so they only collected invertebrates and water quality samples. Nets at this site (n=3) captured Pumpkinseed, Golden Shiner, Black Crappie, Yellow Perch, Bluegill, Rock Bass, Black or Brown Bullhead YOY, Spottail Shiner, Tadpole Madtom, Silver Redhorse, Northern Pike, Logperch, Emerald Shiner, Johnny Darter, Walleye, Brown Bullhead, Bluntnose Minnow, and Mottled Sculpin. Invasive fish captured were White Perch (n=15), Tubenose Goby (n=6), Round Goby (n=32), and Eurasian Ruffe (n=2).

1382 – First sampled on 6-30-2014 by the NRRI team. Last visit on 6-21-2024 and sampled SAV for fish, invertebrates, and water quality. Crew leader Bob Hell noted that the water level was significantly lower than in previous visits. Nets at this site (n=3) captured Bowfin, Banded Killifish, Brook Stickleback, Northern Pike, Pumpkinseed, Bluegill, Brown Bullhead, Bluntnose Minnow, and Emerald Shiner. Invasive fish captured were Threespine Stickleback (n=13), and Round Goby (n=8). There were 5 native crayfish and 1 Painted Turtle as bycatch in fyke nets. Due to storms in the area and heavy rainfall throughout the day when pulling nets, PI Valerie Brady decided to only count the fish captured in fyke nets and not measure them to speed up the process for crew safety.

1436 – First sampled on 6-30-2014 by the NRRI team. During this year's visit on 6-21-2024, crew leader Valerie Brady noted that the entire site was too shallow to sample and had very little vegetation. No vegetation zones meeting sampling criteria were present at this site this year.

1443 – First sampled on 6-29-2014 by NRRI team. During this visit on 6-22-2024, crew leader Valerie Brady noted that this site has significantly changed since the previous visit, with evidence of commercially removed mowed shrubs. She also observed that the maximum depth

of the site was 25-30 cm, with only a little SAV present. No vegetation zones meeting sampling criteria were present at this site this year.

1478 – First sampled on 7-9-2014 by NRRI team. During this visit on 6-23-2024, crew leader Paul Jeffrey noted that the entire site was a sand beach with no vegetation. There were no vegetation zones meeting sampling criteria at this site this year.

1488 – First sampled on 7-11-2014 by NRRI team. Last visit on 6-24-2024 and sampled Sparse Bulrush for fish, invertebrates, and water quality, as well as SAV for invertebrates and water quality. Crew leader Paul Jeffrey noted that the SAV zone was too deep to set fyke nets and the bottom was very soft substrate. He also observed a significant seiche while pulling nets, which made the water levels even deeper. Nets at this site (n=3) captured Rock Bass, Pumpkinseed, Yellow Perch, Bluntnose Minnow, White Sucker, Johnny Darter, Smallmouth Bass, Bowfin, Brown Bullhead, Black Bullhead, Common Shiner, Golden Shiner, and Bluegill. Invasive fish captured were Round Goby (n=54) and Rainbow Smelt (n=1). There was 1 Common Snapping Turtle and 1 Rusty Crayfish as bycatch in fyke nets.

1707 – First sampled on 7-9-2014 by NRRI team. Last visit on 6-23-2024 but crew leader Paul Jeffrey noted that this site was completely sand beach. There were no vegetation zones that met sampling criteria at this site this year.

Benchmark sites

1191 – First sampled on 7-22-2014 by NRRI team. Last visit on 8/27/2024 and sampled SAV and Lily zones for fish, invertebrates, and water quality. A Typha zone was also sampled for invertebrates and water quality. Crew leader Kari Pierce noted that large logs inundated the Typha zone, which became dry quickly, preventing the crew from setting nets. Nets at this site (n=6) captured Yellow Perch, Walleye, Black Crappie, Bluegill, Pumpkinseed, Freshwater Drum, Golden Shiner, Spottail Shiner, Emerald Shiner, White Sucker, Logperch, Mimic Shiner, Silver Redhorse, Shorthead Redhorse, Rock Bass, and Johnny Darter. Invasive fish captured were Round Goby (n=8), Tubenose Goby (n=13), and White Perch (n=1). There were 6 Painted Turtles as bycatch in fyke nets.

7076 – First sampled on 8-9-2021 by NRRI team. Last visit on 7-31-2024 and sampled Lily for fish, invertebrates, and water quality. Crew leader Kari Pierce noted that SAV was present but too deep to sample (over 1.5 m). She also noted that a new box culvert was added under the roadway at this site in either 2022 or 2023 to improve fish movement and connectivity to the main channel. Only two culverts connect this site to the main channel under a roadway: the new box culvert and a smaller pipe culvert. Nets at this site (n=3) captured Black Crappie, Largemouth Bass, Bluegill, Pumpkinseed, Brown Bullhead, Yellow Perch, Black Bullhead, and

Tadpole Madtom. Invasive fish captured were Tubenose Goby (n=5). There were 24 Painted Turtles and 3 Common Snapping Turtles as bycatch in fyke nets.

7080 – This year, PI Valerie Brady created a new site to sample more wetlands upriver on the Wisconsin side of the St. Louis River Estuary. Last visit on 8-21-2024 and sampled SAV for fish, invertebrates, and water quality, as well as a Lily zone for invertebrates and water quality. Crew leader Kari Pierce noted that the Lily zone patches were not large enough to fit three fyke nets, so the crew only collected invertebrates and water quality samples. They also observed large sunken logs at this site, and the little Typha present was clogged with wood debris. Nets at this site (n=3) captured Black Crappie, Pumpkinseed, Bluegill, Spottail Shiner, Yellow Perch, Rock Bass, White Sucker, and Golden Shiner. Invasive fish captured were Tubenose Goby (n=6), Round Goby (n=3), and Eurasian Ruffe (n=1). There were 6 Painted Turtles and 1 native crayfish as bycatch in fyke nets.

Extra Sites and Data (from summer 2024)

There were no special request sites this year.

Extra Data: In collaboration with Amanda Suchy at Central Michigan University, NRRI crews collected greenhouse gas and pore water samples while conducting their standard CWMP water quality sampling. We sent all samples to Amanda for future laboratory analyses. We collected samples at sites: 985, 945, 1488, 1040, and 1192. In total we collected 67 samples including 15 air, 44 gas, and 8 pore water samples.

Wetland Condition Observations and Results (from summer 2024)

This summer in Lake Michigan, we had to drop several sites due to the lack of vegetation. Lake water level swings from year to year could be affecting these wetlands. In addition, vegetation zones that are more common closer to the shore in shallower water (e.g., Typha or Phragmites) were often too dry or not large enough to sample based on CWMP sampling protocols.

Data Processing

As of April 1st, 2025, the NRRI fish/invert/wq team has completed 100% of the invertebrate ID from the 78 samples collected in the 2024 field season. Field datasheets have all been digitized. All fish, habitat, invertebrate, and water quality data are entered into the database and quality control is complete.

Mid-season QC Check Findings (from summer 2024)

Primary long-time crew leaders Kari Pierce and Bob Hell administered mid-season QC check of fish identification with crew members. In 2024 the NRRI fish/invert/wq team surveyed sites as

one 3-person crew or two 4 person crews. New crew members were always working directly with experienced crew members, so the training and evaluation of new crew members was continuous. No issues were noted.

Audit and QC Report and Findings

None. QC of invertebrate samples between team labs has not occurred yet, but the NRRI team will likely swap samples with LSSU soon. We will complete the QC of invertebrate samples by the end of April.

Additional Funding and Projects

None.

Other Collaboration Activities (from summer 2023)

PI Valerie Brady continues to collaborate with MPCA, MNDNR, and WDNR on restoration planning and evaluation for sites in the St. Louis River Estuary. She also presented research on the CWM program at the Wisconsin Wetland Science conference in February 2025.

Other Data Requests

None.

Related Student Research (from summer 2024)

PI Brady's graduate student, Adam Frankiewicz, completed his work on updating a key to the sphaeriid (fingernail) clams of the Great Lakes region. He used CWMP samples to help with this effort and CWMP field crews collected clams for him. He successfully defended his thesis on April 11, 2024.

TEAM NAME: WESTERN BASIN VEGETATION TEAM AT UNIVERSITY OF WISCONSIN-SUPERIOR

Team Members

- Dr. Nicholas Danz, PI, wetland plant ecologist (since 2011)
- Ryne Rutherford, co-crew leader, botanist (since 2014)
- August Camp, botanist (since 2023)

Training (from Fall 2024 report)

This year crew-leader Rutherford took on surveying all the sites in the western basin team and was assisted by botanist August Camp in these efforts. Rutherford and Camp surveyed all sites together. Rutherford provided training to Camp consistent with the project SOP. In all field work, Rutherford and Camp were paired to ensure sampling protocols are followed correctly, and to assist identifying vegetation to species level.

Challenges and Lessons Learned (from Fall 2024 report)

The field season was routine this past year, with water levels about average allowing sites to be easily accessible and no problems encountered during field surveys.

Site Visit List

The 2025 CWMP field season for the NRRI team consists of 27 sites. Of these 27 sites, there are 17 regular sites, 3 pre-sample sites, 4 resample sites, and 3 benchmark sites. The sites are located in Michigan counties Delta (5), Baraga (1); Wisconsin counties Bayfield (4), Brown (2), Door(3), Douglas (4), Kewaunee (1), and Marinette (2); Minnesota counties Cook (1), St. Louis (1); and Thunder Bay District in Ontario, Canada.

Panel Survey Results

N/A

Extra Sites and Data

N/A

Wetland Condition Observations and Results (from 2023)

Across the western Great Lakes sites, the most notable overall observation was the rebound of vegetation at many Lake Michigan sites after a few years of relatively average and stable water levels. The crew observed greater plant density and diversity in the submergent and wet meadow zones. The emergent zones have been slower to recover, although *Zizania* seems to be rebounding in some areas. The increase in diversity in the wet meadow and submergent zones

is well illustrated at site 1488 where diversity was very high. There were also less ruderal species present at Lake MI sites than the previous several years as the more characteristic and dominant species re-established. The crew did not observe any new invasive species this year and anecdotally *Lythrum salicaria* seemed to be decreasing. In terms of species with high conservation value, the crews noted *Potamogeton vaseyii* at site #985 and site #1192.

Data Processing

All vegetation data have been entered into the CWM database. QC checks were completed by Ryne Rutherford in Fall 2024.

Mid-season QC Check Findings

N/A

Audit and QC Report and Findings

Our team continued to prioritize efforts to import GPS coordinates into the database and input our vegetation data immediately following the end of field sampling. Currently, we are completing QA/QC fixes from prior years of the projects and intend for that work to be completed in March 2025.

Additional Funding and Projects

In addition to CWM work in 2024, we surveyed over 400 points in the St. Louis River Estuary for aquatic macrophytes for the Minnesota Pollution Control Agency (MPCA). The MPCA is currently developing a comprehensive, long-term plan to delist the St. Louis River Area of Concern (AOC) through restoration efforts under a grant from USEPA and other project partners. The monitoring and assessment of aquatic macrophytes and soil at several sites in the estuary at various pre- and post-restoration stages will be used in the AOC delisting process. We will continue to collaborate with MCPA through the coming year and plan to use some historical CWM data to help with the efforts to develop indicators of vegetation quality.

Other Collaboration Activities

We continue to collaborate with former project lead Dr. Jeremy Hartsock to summarize patterns of coastal wetlands and aquatic vegetation in the St. Louis River estuary.

PI Danz took the lead to complete the SOGL Coastal Wetland Vegetation indicator report finished in Fall 2024.

Other Data Requests

N/A

Related Student Research

N/A

US CENTRAL BASIN BIRD & ANURAN TEAM AT THE COFRIN CENTER FOR BIODIVERSITY, UNIVERSITY OF WISCONSIN-GREEN BAY

Team Members

- Erin Giese, PI, bird/anuran ecologist (since 2011)
- Dr. Robert Howe, project advisor, bird/anuran ecologist, emeritus (since 2011)
- One full-time summer field tech (since 2022)
- One part-time spring field tech (since 2022)
- Three full-time summer field tech (new summer 2025)

Training

Field technicians from 2024 have either graduated or shifted to different projects. Therefore, Giese hired three new student technicians for the 2025 field season: One graduate student will conduct anuran, bird, and habitat surveys, one graduate student will conduct bird surveys, and one undergraduate (soon to be master's student) will conduct anuran and habitat surveys. We are fortunate to have one returning graduate student anuran technician for their final field season with us to ensure the new crew is trained properly and information is transferred in preparation for the coming years of this project (if funded again). Lastly, we have one returning bird student technician who will work part time to help on board the new bird technicians with the project through mid-late May and then will shift to other projects.

Giese has started training the new incoming field technicians on bird and anuran identification and will continue through May 2025.

Challenges and Lessons Learned

This year was not a particularly challenging year given Great Lakes water levels have lowered; however, we were assigned several sites located on inaccessible private property or sites that were impossible to access due to private roads, unimproved roads, or remote islands. We were able to access 5 points via kayak.

In response to our team's financial struggles, University of Minnesota Duluth PI Dr. Annie Bracey graciously agreed to transfer \$25,000 of their bird/anuran funding to our team so that we may operate normally and in 2025. UW-Green Bay's Grants Office is working with Central Michigan University and UM Duluth on this financial transfer. Dr. Valerie Brady was instrumental in the initiation of this funding transfer.

Site Visit List (from fall report)

Our team was assigned 45 total wetland sites: 1 in Illinois, 9 in Wisconsin, and 35 in Michigan. Of our 45 assigned sites, there were 3 pre-sample (“P”) sites, 2 re-sample (“R”) sites, and 2 benchmark (“B”) sites. We surveyed 27 sites and dropped the remaining ones that were not accessible (e.g., remote islands, lack of roads) and ones that we could not afford to sample. For example, we did not attempt to survey site 1372 in northern Illinois due to the expense of sampling there and did not rent motorized boats in Upper Michigan to reduce costs.

Panel Survey Results (from fall report)

Our first anuran surveys of the 2024 season took place on April 14, 2024 at sites 1382, 1393, 1435, and 1686 near Sturgeon Bay and Baileys Harbor, Wisconsin. Our last surveys occurred in Green Bay, Algoma, and Suamico, Wisconsin on July 9, 2024 at sites 1443, 1451, and 1670. Cumulatively across all sites and samples, we recorded seven anuran species: American toad, spring peeper, gray treefrog, green frog, northern leopard frog, wood frog, and bullfrog, which are each relatively common and expected species in Great Lakes coastal wetlands. We did not detect any uncommon, unusual, or listed anuran species, and we did not detect chorus frog, which we last recorded by our team in 2021. At 8 of our 96 total anuran point count surveys (96 = 32 point count locations × 3 rounds), we did not detect any anurans calling.

Our first bird surveys of the 2024 season took place on May 29, 2024 at sites 1478, 1487, and 1488 near Escanaba, Michigan. Our last surveys occurred in Green Bay, Algoma, and Suamico on July 9, 2024 at sites 1443, 1451, and 1670. Cumulatively across all sites and samples, we recorded 98 bird species, including many target, marsh-obligate bird species: rails (Virginia Rail only, not Sora), American Coot, Common Gallinule, bitterns (American and Least Bitterns), wrens (Marsh and Sedge), Pied-billed Grebe, terns (Black and Forster’s Terns), Swamp Sparrow, Yellow-headed Blackbird, Wilson’s Snipe, Blue-winged Teal, and Sandhill Crane.

- Listed Bird Species:
 - American Bittern: Imperiled–Vulnerable in Wisconsin (S2S3B) during breeding
 - Site 799
 - Black Tern: Endangered in the state of Wisconsin, threatened in the state of Michigan
 - Site 792
 - Common Gallinule: Threatened in the state of Michigan
 - Sites 792, 1436, 1451
 - Common Loon: Threatened in the state of Michigan
 - Sites 608, 642, 736
 - Caspian Tern: Endangered in the state of Wisconsin, threatened in the state of Michigan
 - Sites 608, 770, 1393, 1478

- Common Tern: U.S. Species of Concern, endangered in the state of Wisconsin, threatened in the state of Michigan
 - Sites 799, 834, 1488
- Forster's Tern: Endangered in the state of Wisconsin, threatened in the state of Michigan
 - Site 608
- Great Egret: Threatened in the state of Wisconsin
 - Sites 608, 616, 636, 1488
- Least Bittern: Imperiled–Vulnerable in Wisconsin (S2S3B) during breeding, threatened in the state of Michigan
 - Site 792
- Purple Martin: Imperiled–Vulnerable in Wisconsin (S2S3B) during breeding
 - Sites 1435, 1436, 1451, 1461
- Yellow-headed Blackbird: Critically Imperiled–Imperiled in Wisconsin (S1S2B) during breeding
 - Site 1451
- Invasive Bird Species:
 - European Starling: sites 1443, 1461
 - Mute Swan: site 635

Extra Sites and Data (from fall report)

Like we have done for the last several years, we collected local habitat variables at every point count location following methods outlined by Birds Canada. These data are not entered into the online CWMP DMS. Instead, hard copies are mailed to Dr. Doug Tozer with Birds Canada who then scans the data forms and conducts OCR so they may be automatically and digitally entered into a database.

Wetland Condition Observations and Results (from fall report)

Unlike 2018–2022, our team hardly had any issues pertaining to high water levels this season since Great Lakes levels have been dropping over the last few years. Only one wetland site that our team sampled was described as “drowned” (i.e., did not have any emergent plants), which was site 1393 in northern Door County, Wisconsin in Baileys Harbor (Figure 28). In terms of wetland quality, sites 608, 616, 636, 792, and 799 in the eastern UP produced high quality bird species, such as American Bittern, Least Bittern, Virginia Rail, Common Gallinule, Black Tern, and Forster's Tern. Many of these sites consisted of few invasive plant species and instead contained native sedges, grasses, rushes, bulrushes, and cattails. Sites 792 and 799 contain some of the highest quality coastal wetlands in the Laurentian Great Lakes system (Figure 29). These extensive wetlands are a part of the Munuscong River complex near the rivermouth in the far

eastern UP and are breeding hotspots for Black Terns, Pied-billed Grebes, Marsh and Sedge Wrens, Swamp Sparrows, and other bird species that use coastal marshes.



Figure 28. Site 1393 in Baileys Harbor, Wisconsin was “drowned out” this field season (2024) and did not have any emergent vegetation.

Data Processing

Summer anuran and bird field technicians have completed double data entry for all 2024 anuran and bird point counts and conducted QA/QC such that all double entries match.



Figure 29. High quality site 799 in the Munuscong River wetland complex in eastern Upper Michigan.

Mid-season QC Check Findings (from fall report)

Two field technicians who helped with this project since 2022 and one technician who helped since 2023 returned for the 2024 field season, though Erin Giese and Field Crew Leaders, Mabel Kirst and Haley Spargur, ensured the crew collected data correctly. Erin Giese also regularly checked bird and anuran observations reported by all team members and addressed any issues as needed. However, because one of our team's bird technicians was new to marsh bird surveys, Erin spent >40 hours training them on bird visual and auditory identification and anuran auditory identification both online and in the field. They already had experience conducting forest bird point count surveys and simply needed additional training on marsh birds and anurans.

Audit and QC Report and Findings

We have finished conducting the latest requested QA/QC checks in the Data Verification Interface portal of the CWMP for the years 2016–2021 and 2022 for bird and anuran data.

Additional Funding and Projects

Nothing to report.

Other Collaboration Activities

We received funding from the WI Department of Natural Resources through EPA's AOC GLRI program to monitor wildlife at Ken Evers Nature Area in lower Green Bay in spring and summer 2025 in an effort to evaluate the success of enhancement and restoration efforts. We will monitor breeding anurans, marsh birds, and coastal birds, migratory waterfowl, muskrats, and coastal wetland mustelids. The "health" or "condition" of the site will be evaluated in a number of ways, one of which involves calculating Howe et al.'s (2024) Indices of Biotic Condition (IBC) of breeding anurans and breeding marsh birds.

Other Data Requests (from fall report)

Audubon Great Lakes (AGL) requested CWMP bird data to assist them with assessing the impact of National Fish and Wildlife Foundation's Sustain Our Great Lakes (SOGL) Program on waterbirds, which includes breeding marsh birds. To assess the impact of SOGL funding, they will implement a Before-After-Control-Impact study design, which includes compiling bird survey data that were collected prior to the onset of SOGL funding. CWMP marsh bird data are being used as part of the "before" treatment data set. AGL implemented surveys in 2021 and 2022 (representing the "after" treatment), which will be compared to CWMP survey data collected prior to 2021. CWMP PIs unanimously agreed to proceed with sharing their data;

therefore, Erin Giese led the collective writing of a data sharing agreement with AGL and CWMP PIs. Todd Redder provided AGL with the CWMP breeding marshbird data set.

Species lists were provided to the Wisconsin Department of Natural Resources in agreement for allowing our team to survey on State Natural Areas.

Related Student Research (from fall report)

UW-Green Bay graduate student Whitney Tank is working with Erin Giese and Dr. Dhanamalee Bandara on developing statistical models of habitat associations with marsh-obligate breeding bird species using CWMP data. They are using local habitat point count data collected for this project to develop these bird-habitat associations, which could be used for informing land management decisions and their effects on breeding marsh birds.

US CENTRAL BASIN, CENTRAL MICHIGAN UNIVERSITY (CMU), BIRD/ANURAN TEAM

Team Members

- Dr. Thomas Gehring, PI (since 2011)
- Bridget Wheelock, team leader (since 2013)
- Mary Benjamin, survey lead, graduate student (since 2023)
- Brendan Jankowski, survey lead, undergraduate, prior field tech (since 2023)
- Samuel Rimatzki, new survey lead (2025)
- Audrey George, field technician (new spring 2025)
- Cole Zumbrunnen, field technician (new spring 2025)

Training

The annual training meeting to coordinate US central basin bird and anuran survey leads on survey protocols and QAPP standards will occur by the end of March at Central Michigan University's Biosciences building under the supervision of the team PI Gehring. B. Wheelock, M. Benjamin, and B. Jankowski have previously obtained the bird and anuran identification certification. S. Rimatzki will complete certification prior to data collection. Incoming field technicians (2) will be trained in data recording, speaker broadcasting, and the functions of handheld GPS units on or prior to data collection.

Challenges and Lessons Learned (from fall report)

No major challenges.

Site Visit List

The central basin bird anuran team has been assigned 47 sites. Four sites are re-samples from last year's list (426, 572, 1279, 1915). Three sites are scheduled as benchmarks (515, 1598, 7061). One site is scheduled as pre-sample for next year (1305). The remaining 39 sites are on their regularly scheduled cycle. Five sites have a web reject status because they are islands requiring significant water navigation to reach the site and will not be sampled by bird and anuran crews (424, 427, 533, 551, 553). One site has a web-reject status due to a lack of emergent wetlands (433). Six sites are currently awaiting permission from landowners and will only be surveyed if a roadside point is found or if permission is provided (491, 556, 564, 1568, 1858). Five sites are potentially inaccessible but require further investigation (437, 508, 682, 1656, 7039).

515 & 7061- benchmarked by Don Uzarski every year to generate a long-term data set for sites representing high (515) and low (7061) extremes of the disturbance gradient.

1598- benchmarked to continue monitoring for response to the oil pipeline leak under the straits of Mackinaw in 2018.

Panel Survey Results (from fall report)

Anurans: First sample date – 9 April 2024; Last sample date 13 July 2024

Table 18. Anurans – 8 species

American Toad (*Anaxyrus americanus*)

Bullfrog (*Lithobates catesbeiana*)

Chorus Frog (Western/Boreal) (*Pseudacris triseriata*/*Pseudacris maculata*)

Gray Treefrog (*Hyla versicolor*)

Green Frog (*Lithobates clamitans*)

Northern Leopard Frog (*Lithobates pipiens*)

Spring Peeper (*Pseudacris crucifer*)

Birds: First sample date – 20 May 2024; Last sample date 12 July 2024

Table 19. Birds-118+ species	Code
Alder Flycatcher	ALFL
American Bittern	AMBI
American Crow	AMCR
American Goldfinch	AMGO
American Kestrel	AMKE
American Redstart	AMRE
American Robin	AMRO
American White Pelican	AWPE
Bald Eagle	BAEA
Baltimore Oriole	BAOR
Bank Swallow	BANS
Barn Swallow	BARS
Barred Owl	BADO
Belted Kingfisher	BEKI
Black Tern	BLTE
Black-and-white Warbler	BAWW
Black-bellied Plover	BBPL
Black-capped Chickadee	BCCH
Black-crowned Night Heron	BCNH
Black-necked Stilt	BNST
Black-throated Green Warbler	BTNW
Blackpoll Warbler	BLPW
Blue Jay	BLJA
Blue-gray Gnatcatcher	BGGN
Blue-winged Teal	BWTE
Brown Thrasher	BRTH
Brown-headed Cowbird	BHCO
Canada Goose	CANG
Carolina Wren	CARW
Caspian Tern	CATE
Cedar Waxwing	CEDW
Chimney Swift	CHSW
Chipping Sparrow	CHSP
Common Gallinule	COGA
Common Grackle	COGR

Common Merganser	COME
Common Nighthawk	CONI
Common Raven	CORA
Common Tern	COTE
Common Yellowthroat	COYE
Double-crested Cormorant	DCCO
Downy Woodpecker	DOWO
Eastern Bluebird	EABL
Eastern Kingbird	EAKI
Eastern Phoebe	EAPH
Eastern Wood-Pewee	EAWP
European Starling	EUST
Forster's Tern	FOTE
Gray Catbird	GRCA
Great Blue Heron	GBHE
Great Crested Flycatcher	GCFL
Great Egret	GREG
Greater Yellowlegs	GRYE
Green Heron	GRHE
Hairy Woodpecker	HAWO
Hermit Thrush	HETH
Herring Gull	HERG
House Finch	HOFI
House Sparrow	HOSP
House Wren	HOWR
Indigo Bunting	INBU
Killdeer	KILL
Least Bittern	LEBI
Least Sandpiper	LESA
Lesser Yellowlegs	LEYE
Mallard	MALL
Marsh Wren	MAWR
Merlin	MERL
Mourning Dove	MODO
Mute Swan	MUSW
N. Rough-winged Swallow	NRWS
Nashville Warbler	NAWA
Northern Cardinal	NOCA

Northern Flicker	NOFL
Northern Waterthrush	NOWA
Osprey	OSPR
Pied-billed Grebe	PBGR
Pileated Woodpecker	PIWO
Prothonotary Warbler	PROW
Purple Martin	PUMA
Red-bellied Woodpecker	RBWO
Red-breasted Nuthatch	RBNU
Red-eyed Vireo	REVI
Red-headed Woodpecker	RHWO
Red-shouldered Hawk	RSHA
Red-winged Blackbird	RWBL
Ring-billed Gull	RBGU
Rose-breasted Grosbeak	RBGR
Ruby-throated Hummingbird	RTHU
Sandhill Crane	SACR
Sedge Wren	SEWR
Semipalmated Sandpiper	SESA
Song Sparrow	SOSP
Sora	SORA
Spotted Sandpiper	SPSA
Swamp Sparrow	SWSP
Tennessee Warbler	TEWA
Tree Swallow	TRES
Trumpeter Swan	TRUS
Tufted Titmouse	TUTI
Turkey Vulture	TUVU
Unidentified blackbird	UBLB
Unidentified duck	UDUC
Unidentified flycatcher	UFLY
Unidentified gull	UGUL
Unidentified large bird	ULBD
Unidentified medium bird	UMBD
Unidentified passerine	UPBD
Unidentified Raptor	URAP
Unidentified shorebird	USHO
Unidentified small bird	USBD

Unidentified sparrow	USPA
Unidentified swallow	USWA
Unidentified Tern	UTER
Unidentified woodpecker	UWPR
Unknown swan	USWN
Veery	VEER
Virginia Rail	VIRA
Warbling Vireo	WAVI
Whimbrel	WHIM
White-breasted Nuthatch	WBNU
White-throated Sparrow	WTSP
Willow Flycatcher	WIFL
Wilson's Snipe	WISN
Winter Wren	WIWR
Wood Duck	WODU
Wood Thrush	WOTH
Yellow Warbler	YEWA
Yellow-bellied Flycatcher	YBFL
Yellow-bellied Sapsucker	YBSA
Yellow-headed Blackbird	YHBL
Yellow-rumped Warbler	YRWA
Yellow-throated Vireo	YTVI

Extra Sites and Data (from fall report)

Site 1598 was requested as a benchmark by Dr. Don Uzarski because of its proximity to the Line 5 pipeline in the Straits of Mackinac. It could be used as a baseline in the event of an oil spill. Sites 1651, and 1652 were benchmarked this year because they were being treated with an herbicide for *Phragmites* control. Sites 515 and 7061 were benchmarked by Dr. Don Uzarski because they represent low and high extremes, respectively, along the disturbance gradient and have long term data sets. These data will be used for developing and improving our indices of biotic integrity and indices of environmental condition. Site 7075 was requested as a benchmark for bird only surveys by Mary Benjamin to continue her research on secretive marsh birds and autonomous recording units. No additional data is collected at any of these benchmarked sites. Site 7079 was added as a benchmark in 2023, which was maintained this season. The request was made by NOAA who has plans to undergo hydrological re-connectivity restoration work. Monitoring began now to have baseline data of pre-restoration conditions.

Wetland Condition Observations and Results (from fall report)

Finding appropriate survey points for Site 7079 was difficult in 2024 as well as 2023 due to inaccessibility to private land and a lack of emergent vegetation at accessible sites. Site 7079.3 was requested for quarantine due to a lack of emergent vegetation. Additionally, Site 7079.3 and 7079.4M were located within 500 meters of each other, which broke the anuran survey protocol. If Site 7079.3 is removed, Site 7079.4M should be included as there was emergent vegetation at this location. Additionally, Site 1851.1 was considered for quarantine due to obstruction of the wetland by woody vegetation, however, it was determined that the surveyor was close enough to the wetland to account for the obstruction. Finally, Site 1279.1 was also considered for quarantine due to significant obstruction of the wetland by thick woody vegetation. Based on satellite information, Site 1279.1 is approximately 90 meters from the wetland, however, marsh birds were detected at this location despite the distance and thick vegetation.

Data Processing

All 2024 data has been entered and undergone QC checks. Data Verification system flags for the years 2016-present have all been addressed and resolved. The 2016-2021 batch of DVI records have been addressed.

Mid-season QC Check Findings (from fall report)

On 20 June 2024, mid-year QC checks were completed for each team lead/data collector (Mary Benjamin, Brendan Jankowski, Kimberly Schraitle) at 2 sites each for anurans and birds this year. Data collectors were 100% proficient in the performance criteria including: 1) correct location of sampling points; 2) accuracy of species-level identification; 3) accuracy of abundance category estimates; 4) correct criteria and techniques used for identification of rare species; and 5) correct use of field survey forms.

Audit and QC Report and Findings (from fall report)

As of 26 September 2024 all data have been QA'd with no flags. All GPS coordinates are confirmed or excellent. All data 2011-present has been QA'd in the Data Verification interface.

Additional Funding and Projects

N/A

Other Collaboration Activities

The site list for this year will require collaboration with land management organizations and private land owners to access and conduct surveys. Management organizations will receive

data in exchange for permission to conduct surveys on their properties. Permission acquisition is ongoing.

Other Data Requests (from fall report)

Data has been requested by and sent to seven landowner organizations as a condition of accessing their lands. This includes data collected at 9 sites. Additionally, the fall report was sent to the Michigan DNR for surveys conducted at 10 sites. Site 1849 data went to the Ohio DNR Department of Natural Areas & Preserves. Site 1855 data went to Ohio's Erie Metroparks. Sites 1864 and 1888 partial data went to Ohio DNR Division of Wildlife. Site 1888 and partial 1883 data went to Ottawa National Wildlife Refuge. Site 589 data went to Little Traverse Conservancy. Site 7075 data went to Shiawassee National Wildlife Refuge. Michigan DNR received data for sites 432, 488, 515, 571, 573, 760, 1275, 1651, 1896, and 7061.

Related Student Research

Kylie McElrath is planning to defend her M.S. thesis during spring 2025. Her thesis research examined factors influencing muskrat abundance in Great Lakes coastal wetlands and changes in muskrat spatial distribution patterns over time.

Megan Bos is currently writing her M.S. thesis examining the influence of muskrat houses on water chemistry and plant communities in Great Lakes coastal wetlands.

Megan Casler continues research generating a hierarchical, multi-season occupancy modeling of Rallidae species using basin-wide bird, invertebrate, and vegetation data from the years 2011-2022. Analysis completion and defense anticipated May 2025.

Mary Benjamin is currently working on analyzing her ARU data collected from 2023 and 2024 sites 515, 523, 571, 589, 591, 1273, 7061, and 7075 by using BirdNet and RavenPro.

US CENTRAL BASIN FISH, INVERTEBRATE AND WATER QUALITY TEAM

Team Members

The US Central Basin Fish, Invertebrate and Water Quality Team consists of PIs and members from the following universities:

Central Michigan University (CMU) crew:

- Dr. Donald G. Uzarski, PI (since 2011)

- Bridget Wheelock, Uzarski lab manager, team leader (since 2018)
- Molly Gordon, lead invertebrate taxonomist (since 2011)
- Matthew Sand, water quality technician (since 2020)
- Kimberly Schraitle, lab manager (since 2022)
- Morgan Noffsinger, crew leader (since 2023)
- Howard Mitchell, crew lead (since 2024)
- Three summer field technicians (new 2025)

Grand Valley State University (GVSU) crew:

- Dr. Carl Ruetz III, PI (since 2011)
- Dr. Matthew Cooper, PI (since 2011)
- John Lawrence, team leader (since 2023)
- Emily Eberly, graduate research assistant (since 2024)
- Caden Shannon, summer technician (since 2024)
- John Gargas, summer technician (since 2024)

University of Notre Dame (UND) crew:

- Dr. Gary Lamberti, PI (since 2011)
- Sarah Klepinger, Lamberti lab manager, team leader (since 2019)
- Amaryllis Adey, graduate student (since 2023)
- Caitlin Day, summer technician (since 2024)

Lake Superior State University (LSSU) crew:

- Dr. Ashley Moerke, PI (since 2011)
- Connor Arnold, crew lead (since 2023)
- Sam Ritmatski, undergraduate technician (since 2024)
- Alana Schofield, undergraduate technician (since 2024)
- Nikki Perigo, research technician (since 2024)
- Silas Dunn, macroinvertebrate taxonomist (since 2022)

Training (from fall report)

Central Michigan University hosted the Central Basin training at site 515 in Saginaw Bay on 17 June 2024 and 18 June 2024, attended by GVSU and LSSU. The training was led by Bridget Wheelock who has been a part of the CWMP since 2012. The topics covered included GPS waypoint collection and navigation, water quality collection, invertebrate sampling and picking, fyke net setting/retrieval, and fish handling and identification. Each team used their own

equipment to familiarize themselves with their equipment. Teams conducted additional water quality processing training and certification on their own to familiarize themselves with their equipment.

Additional training for the CMU crew was completed at site 515 in Saginaw Bay on 12 June 2024 as well as in Mount Pleasant and Littlefield Lake from 13 May 2024 to 14 June 2024. The topics covered included lab and field safety, boater safety, IACUC, water quality collection, titration, filtering, in situ data collection, GPS navigation, invertebrate sampling and picking, fyke net setting/retrieval/repair, fish identification, boat operation, and trailering. Mid-season checks were provided by Bridget Wheelock at site 1605 on 24 and 25 of July 2024 to ensure protocols were being followed. Taxonomist Molly Gordon is currently working on training undergraduate students Ace McClelland and Olivia Klein and graduate student Morgan Noffsinger to identify macroinvertebrates in the lab.

University of Notre Dame training occurred on 10 June 2024 at North Chain Lake in South Bend, IN. Proper technique for water and macroinvertebrate collection was demonstrated and practiced, as well as how to set a fyke net and launch a boat.

Fish ID training was provided for the LSSU crew by LSSU PI (Moerke) and CFRE Research Technician (N. Perigo) at the Barch Center for Freshwater Research and Education using the center's preserved specimens. All three crew members identified at least 95% of fish correctly. GPS training also occurred before field season began. Initial field training was provided by LSSU crew chief (Arnold) at Ashmun Bay where the crew went through equipment deployment and sample collection process, and then reviewed lab protocols with the water quality lab manager. Mid-season checks were provided by Nikki Perigo to ensure protocols were being followed.

Update: Central Michigan University will be hosting the 2025 Central Basin fish/invertebrate/water quality training at site 515 in Saginaw Bay on June 16th and 17th. The training will be led by CMU crew leader Bridget Wheelock who has been part of the CWMP since 2012. The topics covered will include water quality collection (in situ data collection, filtering, and titration), GPS navigation, site/zone selection, invertebrate sampling and picking, setting and retrieving fyke nets, and fish handling/identification. The GVSU and LSSU crews will supply and use their own equipment to familiarize themselves with the equipment. The training will be attended by LSSU, GVSU and CMU.

The UND crew will not attend the annual training because they have three experienced team members who can train any new technicians. Instead, the crew will go out on a local lake and practice setting fyke nets, capturing invertebrates, and taking water samples.

Sarah Klepinger, Caitlynn Day and Amaryllis Adey of UND are trained and experienced in CWMP protocols. Any undergraduate technicians will be trained immediately after they are hired.

Interviews for LSSU's summer crews are under way now and expect to be complete by end of March. Field safety training will begin the first week of May at LSSU.

Challenges and Lessons Learned (from fall report)

CMU encountered challenges finding inundated vegetation zones as water levels were lower than 2019. Multiple sites visited throughout this summer no longer had inundated vegetation zones compared to 2019 (see Figure 30).



Figure 30. Wet meadow zone at site 1273 in 2019 (left) and 2024 (right). Photo credit: CMU.

One of the main challenges faced by the GVSU crew this season was navigating around rocky shallow waters, as the water levels this year were low. With many of the sample sites being in areas of high winds and regular storm patterns, the crew had to constantly monitor the weather radar throughout the season and adjust sampling times as needed.

The season went remarkably smoothly for UND with only a few equipment hiccups. UND's pH meter broke near the end of the season, and it is unclear whether it was due to wear or user error. Next year, more time will be put into teaching and emphasizing proper instrument care.

Update: For UND, the 2024 field season went fairly smoothly, likely because so many of our crew were experienced in the CWMP protocols. Sarah Klepinger did learn a lesson about kayak safety, but there were no lasting consequences.

Site Visit List (from fall report)

The US Central Basin was assigned 47 sites (21 CMU, 11 GVSU, 8 LSSU, 7 UND), seven of which were benchmarks (515,792, 1598,1651,1652,7079,7061), and 6 of which were re-sample sites

(436, 444, 591, 635, 736, 922). Sites 1651 and 1652 were requested as a benchmark by Denny Albert because herbicides are being applied for Phragmites. Sites 515 and 7061 were benchmarked because they represent low and high extremes, respectively, along the disturbance gradient and have long term data sets. Site 792 was requested as a benchmark because it represents sites in the mildly impacted category, and we are trying to sample more sites on the high and low ends of the environmental gradient. Site 1598 is close to the line 5 oil pipeline in the Mackinac Straits and was requested as a benchmark to gather historical data in the event of an oil spill. Site 7079 was requested as a benchmark by Dr. Alan Steinman of GVSU to document the restoration of the Mona Lake celery flats.

CMU sampled 11 sites, visit rejected 6 sites, and could not sample 1 site due to access issues (757). The 6 rejected sites (572, 755, 1266, 1273, 1563, and 1783) were all due to lack of wetland vegetation or low water levels.

GVSU sampled 9 sites and rejected 2 sites. Sites 444 and 446 were not sampled because there was no wetland vegetation to sample. Both sites were surrounded by large boulders halfway exposed out of the water, and any vegetation that may have been possible to sample in previous years was distinctly landbound and would be seemingly difficult to sample even in high water. Site 1279 was not accessible by motorboat and was only sampled for WQ and inverts with permission from the landowner to access the water via their property. This landowner was friendly and loaned the crew his kayaks to access the polygon. At site 1306 the crew was approached by an unfriendly property owner who believed the survey was being conducted on his property. He does not actually have any ownership or riparian rights to this wetland, but he requested that the next time samples are done that he be contacted so he knows when to expect nets in the water.

The University of Notre Dame (UND) was assigned seven sites to access for 2024. Of the seven sites assigned to UND, one (1660) was rejected without visiting due to reports from other teams that the wetland was no longer present at that site. This decision was approved by Val Brady. Of the other six sites, four were along the Detroit River (421, 426, 1915 and 1917). Two more were sampled along the Kalamazoo River near Saugatuck, MI (1651 and 1652). Within these six sites, eleven zones were sampled for water and macroinvertebrates, and five of those zones were also sampled for fish.

LSSU was assigned eight sites to sample, three of which (sites 910, 918, and 5357) were rejected for sampling due to lack of or mixed vegetation. An additional 3 (635, 642, and 719) sites in the Les Cheneaux region were picked up from CMU's crew.

Update: A total of 44 sites have been assigned to the Central Basin fish/invertebrate/water quality crews for the 2025 field season. Of those sites 31 are panel year sites, 4 are pre-sample sites, 6 are resample sites from the 2024 field season and 3 are benchmark sites. The benchmark sites are East Saginaw Bay Coastal Wetland #5 (515), Point St. Ignace Wetland (1598), and Indian Harbor Wetland (7061). Site 1598 was requested by Don Uzarski and is being monitored due to potential environmental changes in the Straits of Mackinac. Sites 515 and 7061 were benchmarked because they represent high and low extremes, respectively, along the disturbance gradient and have long term data sets.

CMU's crew was assigned 17 sites for the 2025 season (614, 627, 630, 651, 658, 696, 700, 718, 719, 726, 753, 781, 857, 868, 1568, 1598, and 7061). The GVSU crew was assigned 11 sites for the 2025 season (424, 450, 453, 508, 515, 524, 539, 572, 1279, 1305, 1310). Their plan is to assess and sample all assigned sites. The LSSU crew was assigned 9 sites for the 2025 season (805, 811, 827, 833, 873, 903, 5046, 5757, and 7036). The UND crew has been assigned seven sites for the summer of 2025 (510, 426, 1918, 1919, 1915, 1656, and 1653).

Panel Survey Results (from fall report)

Sampling started on 4 June 2024 and ended for the season on 14 August 2024. The following tables list zones sampled for each site, non-native species by site, and reptile and amphibian species captured in fyke nets, respectively.

Vegetation Zones by Site

Site	Vegetation Zone
421	<i>Phragmites</i>
	SAV
426	<i>Phragmites</i>
436	Lily
	<i>Typha</i>
454	<i>Phragmites</i>
493	<i>Typha</i>
	Dense Bulrush
499	PSP
	<i>Phragmites</i>
	<i>Typha</i>
515	<i>Phragmites</i>
	<i>Typha</i>
	Sparse Bullrush
591	<i>Typha</i>

	Sparse Bulrush
616	<i>Typha</i>
	Sparse Bulrush
617	<i>Typha</i>
629	<i>Typha</i>
	Sparse Bulrush
632	<i>Typha</i>
	Sparse Bulrush
635	Lily
	<i>Typha</i>
642	Sparse Bulrush
	<i>Typha</i>
719	Sparse Bulrush
736	Dense Bulrush
770	<i>Typha</i>
	Dense Bulrush
	SAV
792	Dense Bulrush
	Sparse Bulrush
	Lily
	<i>Typha</i>
795	Sparse Bulrush
	<i>Typha</i>
	<i>Phragmites</i>
817	Lily
857	Dense Bulrush
900	Lily
922	Lily
1279	<i>Typha</i>
1306	Lily
1598	<i>Phragmites</i>
	<i>Typha</i>
	Sparse Bulrush
1605	<i>Typha</i>
1626	<i>Typha</i>
	Lily
	Sparse Bulrush
1651	PSP
	<i>Typha</i>
1652	PSP

1915	Lily
	<i>Typha</i>
1917	<i>Phragmites</i>
	Lily
	<i>Typha</i>
7061	Dense Bulrush
7079	Open Water

Non-native Species by Site

Site	Common Name	Taxa Name
436	Round Goby	<i>Neogobius melanostomus</i>
	Freshwater Tubenose Goby	<i>Proterorhinus semilunaris</i>
454	Round Goby	<i>Neogobius melanostomus</i>
493	Round Goby	<i>Neogobius melanostomus</i>
	Goldfish	<i>Carassius auratus</i>
499	Round Goby	<i>Neogobius melanostomus</i>
	Goldfish	<i>Carassius auratus</i>
	Common Carp	<i>Cyprinus carpio</i>
515	Round Goby	<i>Neogobius melanostomus</i>
591	Round Goby	<i>Neogobius melanostomus</i>
	Freshwater Tubenose Goby	<i>Proterorhinus semilunaris</i>
617	Common Carp (YOY)	<i>Cyprinus carpio</i>
629	Round Goby	<i>Neogobius melanostomus</i>
632	Round Goby	<i>Neogobius melanostomus</i>
635	Round Goby	<i>Neogobius melanostomus</i>
642	Round Goby	<i>Neogobius melanostomus</i>
719	Round Goby	<i>Neogobius melanostomus</i>
736	Round Goby	<i>Neogobius melanostomus</i>
770	Round Goby	<i>Neogobius melanostomus</i>
1598	Round Goby	<i>Neogobius melanostomus</i>
	Freshwater Tubenose Goby	<i>Proterorhinus semilunaris</i>

1605	Freshwater Tubenose Goby	<i>Proterorhinus semilunaris</i>
1626	Round Goby	<i>Neogobius melanostomus</i>
1651	Common Carp	<i>Cyprinus carpio</i>
	Round Goby	<i>Neogobius melanostomus</i>
1917	Common Carp (YOY)	<i>Cyprinus carpio</i>
7061	Round Goby	<i>Neogobius melanostomus</i>
7079	Goldfish	<i>Carassius auratus</i>

Reptile and Amphibian Species Captured in Fyke Nets

Site	Common Name	Taxa Name
421	Common Snapping Turtle	<i>Chelydra serpentina</i>
	Painted Turtle	<i>Chrysemys picta</i>
	Map Turtle	<i>Graptemys picta</i>
426	Painted Turtle	<i>Chrysemys picta</i>
436	Painted Turtle	<i>Chrysemys picta</i>
	Musk Turtle	<i>Sternotherus odoratus</i>
	Map Turtle	<i>Graptemys picta</i>
493	Painted Turtle	<i>Chrysemys picta</i>
	Common Snapping Turtle	<i>Chelydra serpentina</i>
499	Painted Turtle	<i>Chrysemys picta</i>
515	Common Snapping Turtle	<i>Chelydra serpentina</i>
591	Painted Turtle	<i>Chrysemys picta</i>
617	Common Snapping Turtle	<i>Chelydra serpentina</i>
629	Painted Turtle	<i>Chrysemys picta</i>
632	Painted Turtle	<i>Chrysemys picta</i>
635	Painted Turtle	<i>Chrysemys picta</i>
736	Painted Turtle	<i>Chrysemys picta</i>
770	Painted Turtle	<i>Chrysemys picta</i>
817	Painted Turtle	<i>Chrysemys picta</i>
857	Painted Turtle	<i>Chrysemys picta</i>

900	Painted Turtle	<i>Chrysemys picta</i>
1598	Painted Turtle	<i>Chrysemys picta</i>
1626	Painted Turtle	<i>Chrysemys picta</i>
	Common Snapping Turtle	<i>Chelydra serpentina</i>
1651	Map Turtle	<i>Graptemys picta</i>
	Painted Turtle	<i>Chrysemys picta</i>
1917	Painted Turtle	<i>Chrysemys picta</i>
	Common Snapping Turtle	<i>Chelydra serpentina</i>
7079	Painted Turtle	<i>Chrysemys picta</i>
	Common Snapping Turtle	<i>Chelydra serpentina</i>
	Map Turtle	<i>Graptemys picta</i>

Extra Sites and Data (from fall report)

Site 7079 was requested as a benchmark by Dr. Alan Steinman of GVSU to document the restoration of the Mona Lake celery flats. Sites 515 and 7061 were benchmarked by Dr. Don Uzarski because they represent low and high extremes, respectively, along the disturbance gradient and have long term data sets. Site 792 was also benchmarked because it represents somewhat high (mildly impacted) values along the disturbance gradient. These data will be used for developing and improving our indices of biotic integrity and indices of environmental condition. Site 1598 was requested as a benchmark to gather historical data in the event of an oil spill since it is close to the line 5 oil pipeline in the Mackinac Straits.

CMU collected extra soil cores, water samples (pore and surface) and air samples for Dr. Amanda Suchy who is leading the greenhouse gas project at CMU. GVSU also collected extra water samples and air samples at most sampled sites for dissolved greenhouse gas analyses. Water samples were not collected at some GVSU sites due to lack of refrigerator space midway through the season. These samples were delivered to Morgan Noffsinger at CMU for Dr. Amanda Suchy. These data are not entered into the CWMP data management system and are stored on drives and hard copies at the CMU Wetland Ecology Lab.

Wetland Condition Observations and Results (from fall report)

The CMU team noticed that water levels were low to the point where some zones could not be sampled (predominantly Wet Meadow zones). Overall, most of the wetlands sampled by the GVSU team were accessible by boat and intact. Wetland sediment ranged from deep organic materials to gravel bottoms, influencing a wide array of sampling experiences. Water levels were low, making navigation difficult at some sites but not impossible. Only two of the eleven GVSU sites no longer had existing wetland habitat. For the LSSU team, water levels remained relatively high during the sampling season. Zebra/Quagga mussels were observed at site 719 in northern Lake Huron.

Data Processing (from fall report)

Central basin teams are still in the process of entering habitat, fyke, and in situ field data. These data are expected to be entered and QC'd within the next month. All field data for both UND and LSSU has been entered and QC'd.

Nine macroinvertebrate samples were given to Bridget Wheelock from Central Michigan University on 17 June 2024 at the training at site 515 from GVSU. On 19 August 2024, 57 water samples (raw, filtered/dissolved nutrients, and dissolved filtered ions) and 43 macroinvertebrate samples were delivered to Morgan Noffsinger at Central Michigan University by Emily Eberly of Grand Valley State University. On 18 September 2024, 24 water samples were delivered to Morgan Noffsinger at CMU from Notre Dame. Water samples from LSSU will be shipped to CMU by the end of September. Macroinvertebrate identification is underway in the CMU and UND labs. Macroinvertebrate identification has not been completed yet for LSSU's summer 2024 samples, but will begin in November and be led by Silas Dunn, who has been identifying LSSU samples for the past several years.

Nineteen chlorophyll-a samples were mailed to the Lamberti Lab at the University of Notre Dame on 20 August 2024 for processing from GVSU. The Chlorophyll-a samples were received by the lab on 21 August 2024 and are being processed with results expected in early 2025. LSSU and CMU will be shipping their chlorophyll-a samples by the end of September. Once these samples arrive, they will likely be analyzed in December.

Update: For CMU, all habitat and fyke data have been entered into the CWMP database and QC'd. Chlorophyll-a results have been received and will be entered into the database and QC'd by the end of March. All water quality processing is finished, and data has been entered and QC'd in the CWMP database. All CMU and GVSU invertebrates have been identified by the CMU crew. Samples are still in the process of being QC'd and entered in the CWMP database,

however, we are planning on having this finished by April 1, 2025. We are planning on trading randomly selected samples with another lab within CWMP for secondary QA. All flagged errors within the DVI have been addressed for 2016-2021 and we are working on 2011-2015 flagged errors.

For GVSU, all data entry for fish/invert/WQ was completed between September 9, 2024, and October 15, 2024. Data QC was completed between October 15, 2024, and November 15, 2024. All WQ laboratory results were entered into the database and QC'd in March 2025.

For ND, all 2024 habitat and fish data have been entered into the CWMP database and QC'd. Chlorophyll-a analysis is about 60% complete, and we are planning on having it finished and sent out by March 21, 2025. Macroinvertebrate identification is about 65% complete, and we are planning to have it finished and entered into the database before April 1, 2025. We are planning on trading samples with our collaborating QA laboratory within the next month.

For LSSU, Macroinvertebrate ID is complete, and data entry and QC are complete. QC samples are being sent to NRRI for the final check this week. Chlorophyll-a analyses were sent to ND in October, but they have not been received to date. Water quality data were received from CMU and entered into the database and QC'd. We believe we worked through all data verification reports to correct past data entry or collection errors.

Mid-season QC Check Findings (from fall report)

Bridget Wheelock provided the mid-season QC check for the CMU crew and observed sampling in accordance with the SOP at site 1605 on 23 July 2024 and 24 July 2024 with no issues reported. The crew correctly located sampling points, collected data, and identified fish species.

The mid-season QC check did not occur for GVSU this season as Dr. Carl Ruetz was unable to participate in the mid-season QC check. Crew leader John Lawrence was with the field crew during all stages of sampling and observed that sampling occurred in accordance with the SOP.

University of Notre Dame's mid-season QC check occurred on 17 July 2024 and 18 July 2024 by ND's PI, Gary Lamberti. He was largely satisfied with their performance, and only made small adjustments in regard to efficiency (they were spending more time than necessary enthusiastically talking about the fish they captured). However, he was impressed with everyone's fish identification skills.

Research technician Nikki Perigo provided the mid-season QC for LSSU and observed that sampling occurred in accordance with the SOP.

Audit and QC Report and Findings (from fall report)

All field data have been entered into the database and QC'd for University of Notre Dame and Lake Superior State University.

Additional Funding and Projects (from fall report)

Dr. Amanda Suchy from Central Michigan University is leading a project funded by CIGLR where they are investigating spatial and temporal drivers of dissolved greenhouse gases (GHGs) in coastal wetlands of the Great Lakes. By leveraging the sampling done by the CWMP they are able to collect dissolved gas samples across a large spatial scale which would not be possible with one sampling crew alone. They will examine how dissolved GHGs are affected by water chemistry, vegetation cover, and surrounding land use. With a few measurements of this kind, this study will provide baseline data for emissions of GHGs from coastal wetlands of the Great Lakes, which can inform future investigations and climate models for the region. She is also investigating patterns of microplastic deposition in coastal wetlands of the Great Lakes and whether wetland connectivity to the open water environment, vegetation cover, or land use are predictive of microplastic concentrations. For this project, sediments are collected at a subset of coastal wetlands sampled by the CWMP and microplastics are quantified using density separation. Preliminary results suggest that wetland connectivity and vegetation cover are more predictive of microplastic concentrations than nearby land use.

Other Collaboration Activities

None

Other Data Requests

None

Related Student Research

Morgan Noffsinger, a CMU graduate student and crew leader, is using GLCWMP data for her thesis regarding the effects of changing water levels on fish communities within Great Lakes coastal wetlands. CMU undergraduate student Julia Shablin is looking at microplastics composition in soil and additional water samples collected by central basin teams at all sampled 2024 sites. CMU undergraduate student Marta Kendzioriski (former vegetation and fish/invertebrate/water quality crew member) is using CWMP data to look at relationships between focal bird species presence and invertebrate community data.

Emily Eberly, a graduate student under the advising of Dr. Matt Cooper, is using data collected through the GLCWMP for a thesis project analyzing multiple indicators as an assessment of wetland health. Emily presented a poster at the Michigan chapter of AFS annual meeting using data from the GLCWMP in February 2025.

Eberly, E.A., Brady, V. J., Lamberti, G. A., Ruetz III, C. R., Uzarski, D. G., & Cooper, M. J. (2025, February 19-21). Fish assemblages in Lake Michigan coastal wetlands [Poster Presentation]. Michigan chapter of the American Fisheries Society, Marquette, MI, United States.

LSSU undergraduate student Clayton Robertson presented his senior thesis on use of coastal wetlands in the St. Marys River by black bass at the Michigan American Fisheries Society conference last March and Clayton was awarded the best student poster award. Connor Arnold, an LSSU undergraduate student, is conducting his senior research on the extent and effects of hypoxia on coastal wetland fishes in the St. Marys River. Connor collected DO and fish data during the day and night at two wetlands that experience hypoxia and two that do not. He will present his research at the senior symposium in December.

US CENTRAL BASIN VEGETATION TEAM

Team Members

- Dr. Dennis Albert, PI, wetland vegetation ecologist/botanist (since 2011)
- Matthew Sand, crew leader, wetland plants and water chemistry (since 2017)
- Kim Schraitle, CMU lab manager, wetland plants (since 2022)
- Katlyn Groulx, CMU crew leader, wetland plants (since 2023)
- Two CMU summer field technicians (2025)

Training

Matthew Sand (5 years of crew leader experience) refreshed Emma Waatti (second year graduate student and crew leader) and trained Katlyn Groulx (second year on crew, first year crew leader) the week of June 17th–June 20th. This included SOP training and sampling logistics as a crew leader. On June 18th, Matt Sand refreshed/trained Emma, Katlyn and the two new technicians in Mount Pleasant wetlands. Topics covered included: identification of common Michigan coastal wetland macrophytes, proper use of GPS for taking waypoints, using a compass to set transect bearings, percent cover estimation, collection of plants for expert ID, and completion of datasheets.

Matt Sand refreshed/trained Emma Waatti, Katlyn Groulx, Kim Schraitle, and the two new summer field technicians on in-situ vegetation protocols at Nayanquing Point Wildlife Area Wetland (493) in Linwood, MI on June 19th, 2024. They also calibrated individual percent cover estimates.

On June 14th, 2024, the crews and lab managers met with Dr. Dennis Albert via Webex to discuss the upcoming sample year and ask questions about macrophyte identification and sampling protocols. Following the meeting, crew members were tested on a subset of specimens covered in training PowerPoints and collected from Mount Pleasant wetlands. Crew leaders/lab managers, Emma Waatti, Katlyn Groulx, Matthew Sand, and Kim Schraitle all correctly identified at least 90% of the specimens.

Update: Matt Sand will lead a plant identification and sampling protocol training in mid-June, and Dr. Dennis Albert will lead a virtual plant identification training in late June for the 2025 field season.

Challenges and Lessons Learned (from fall report)

Due to water levels dropping from the previous high-water years in 2020 and 2021, the patterns of vegetation zonation still continue to be in-flux within the wet meadow, emergent, and submergent vegetation zones. This made it difficult to judge the start waypoint in a few central basin wetlands. Some start waypoints were placed at small tree lines that are now standing dead; in these cases, the starting waypoint was pushed back to the new, well-developed tree line. When samplers were unclear on how to treat zones, they consulted one of the more experienced crew leaders, Matthew Sand, for confirmation and additionally added information about zonation in the notes section on the data sheets.

At some sites, crews also experienced difficult or dangerous sampling conditions due to the drop in the water levels. One site in particular, Bar Lake Wetland #2 (1279), the sample crew experienced deep organic matter in the inundated areas of the wetland (Figure 31). They could only sample this wetland by slowly canoeing through the dense floating vegetation and muck. Additionally, at Swan Creek Wetland (1915), there was no way to access the start waypoint other than walking through several meters of invasive *Phragmites australis* that was in deep organic matter. Private property bumped up to the start point, therefore, no start waypoint was taken at the site, and was instead taken via Google Earth later in the lab.



Figure 31. Dense vegetation with extreme depths of organic matter found at Bar Lake Wetland #2 (1279) in 2024.

Site Visit List

The Central Basin vegetation crews sampled 43 sites: 24 panel sites from 2024, 5 resampled panel sites from 2023, 8 benchmark sites, and 6 pre-sample panel sites from 2025.

Goulais River Wetland (5357) was the only Canadian panel site sampled by the Central Basin crew with all others being in the United States.

Crews could not access 2 sites due to either landowner permissions or access issues. Landowner permission and access could not be obtained for Palmers Point Area Wetland (918) and Bass Cove Area Wetland (757), so these sites were rejected from 2024 sampling.

Update: 45 sites have been assigned the Central Basin vegetation crew for the 2025 field season. Of those sites, 35 are panel year sites, 6 are resample sites from the 2024 field season, and 4 are benchmark sites. The benchmark sites are East Saginaw Bay Coastal Wetland #5 (515), Mackinac Creek Wetland (616), Point St. Ignace Wetland (1598), and Indian Harbor Wetland (7061). All were requested by Dr. Don Uzarski and Dr. Denny Albert. Site 515

represents high disturbance, 616 is one of the most protected sites on Lake Huron and represents low disturbance, 1598 is being monitored due to potential environmental changes in the Straits of Mackinac, and 7061 is one of the most pristine sites on Lake Michigan and represents low disturbance.

Panel Survey Results (from 2024 sampling)

In the US Central Basin, the first day of vegetation sampling took place on June 19th, 2024, and the last day of sampling took place on September 6th, 2024. Sampling was conducted in order from Southern Sites (Lake Erie and SE Lake Michigan) to Northern Sites (Lake Superior) to catch vegetation when most fertile and identifiable.

Most anthropogenically disturbed sites (i.e. Swan Creek Wetland – 1915, Mouillee Marsh – 1917, West Saginaw Bay #1 – 499) lacked the presence of wet meadow zones due to invasive *Typha* and *Phragmites*, these sites generally had less species observations as well. Several island sites of little disturbance (Hog Island Wetland #9 – 1266, Jensen Harbor Wetland – 1279, and Temperance Island Wetland #1 – 1605) had very diverse wet meadow zones but sparse emergent vegetation and almost no submergent vegetation due to wave energy.

In general, we noted a few expansions of invasive species and a few new sites for newly collected species. Species that were later identified by Dr. Dennis Albert as new observations, were sent to Todd Redder to be placed on the Michigan Coastal Wetland Vegetation master list.

Iris lacustris (Dwarf Lake Iris) was observed near the sample transects at Thompsons Harbor Wetland #1 (572) but not in any quadrats. This species is well established in this area of the state, but its continued presence is still noteworthy. *Rorippa aquatica* (Lake Cress) was observed with small percent cover in a quadrat at Munuscong Lake Wetland #5 (795). This is a unique submergent species of Special Concern status, which often loses all of its leaves when collected, but our collection arrived intact, although it promptly lost a few leaves to assist in identification! This observation will be sent to Michigan Natural Features Inventory as it has not been recorded in this county before.

Update: The three panel sites from 2024 with the highest species richness are Nayanquing Point Wildlife Area Wetland (493) with 92 taxa, Duck Island Area Wetland (736) with 87 taxa, and Cheboygan Area Wetland #3 (591) with 84 taxa.

Six sites were sampled for the first time in 2024: Trenton Channel Wetland (426), Thompsons Harbor Wetland #1 (572), McKay Bay Wetland (719), Jensen Harbor Wetland (1273), Bar Lake Wetland #2 (1279), and Goulais River Wetland (5357).

Invasive species recorded in 2024 include:

Invasive Species	Sites
<i>Acorus calamus</i>	817, 922 [†]
<i>Agrostis gigantea</i>	1273 ⁻ , 1605 [†] , 1783 [†]
<i>Butomus umbellatus</i>	436, 1917 [†]
<i>Centaurea stoebe</i>	572 ⁻ , 1605 [†]
<i>Cirsium arvense</i>	792 ^{*†} , 795 [†] , 900 [†] , 1915 [†]
<i>Frangula alnus</i>	1306 [†]
<i>Hydrocharis morsus-ranae</i>	421, 436, 493, 515 [*] , 591, 792 [*] , 795, 1915, 1917
<i>Linaria vulgaris</i>	454 [†] , 591 [†] , 616 ^{*†}
<i>Lysimachia nummularia</i>	421 [†] , 616 ^{*†}
<i>Lythrum salicaria</i>	436, 444, 493 [†] , 499 [†] , 515 [*] , 591, 616 [*] , 632, 736 [†] , 770, 792 ^{*†} , 795 [†] , 817, 922, 1306, 1563, 1626 [†] , 1651 [*] , 1652, 7079 [*]
<i>Myriophyllum spicatum</i>	719 ⁻ , 736, 922 [†] , 1306, 1651 [*] , 1652, 1915, 1917, 7061 [*]
<i>Nitellopsis obtusa</i>	493 [†] , 1306 [†] , 1651 ^{*†}
<i>Pastinaca sativa</i>	736 [†] , 1563 [†]
<i>Persicaria lapathifolia</i>	795 [†] , 1563 [†]
<i>Persicaria maculosa</i>	736, 770 [†] , 857 [†] , 1598 [*] , 1651 ^{*†}
<i>Phalaris arundinacea</i>	436 [†] , 444, 493, 499, 591, 616 [*] , 617, 632, 642, 719 ⁻ , 736, 770, 792 [*] , 817, 900, 922, 1273 ⁻ , 1563, 1626, 1783, 7061 [*] , 7079 ^{*†}
<i>Phragmites australis</i>	421, 426 ⁻ , 436, 444, 493, 499, 515 [*] , 817 [†] , 1279 ⁻ , 1598 [*] , 1651 [*] , 1652 [†] , 1915, 1917, 7079 ^{*†}
<i>Poa compressa</i>	1273 ⁻ , 1605, 7061 [*]
<i>Potamogeton crispus</i>	421, 426 ⁻ , 493, 515 [*] , 1306, 1651 [*] , 1652, 1915
<i>Typha angustifolia</i>	493, 591, 616 [*] , 617, 629 [†] , 632, 635, 642, 770 [†] , 792 [*] , 1279 ⁻ , 1306 [†] , 1563 [†] , 1605 [†] , 1626, 1651 [*] , 1652 [†]
<i>Typha glauca</i>	436, 444 [†] , 499, 616 [*] , 617, 635, 792 [*] , 795 [†] , 1279 ⁻ , 1306, 1598 [*] , 1626, 1651 [*] , 1652, 1917, 7079 [*]

*Extra/Benchmark site

[†]First observation occurred in 2024.

⁻First time sampling occurred in 2024.

Michigan Species of Special Concern recorded in 2024 include:

Species of Special Concern	Sites
<i>Myriophyllum alterniflorum</i>	635†, 857†
<i>Nelumbo lutea</i>	1915, 1917
<i>Ranunculus gmelinii</i>	591†
<i>Rorippa aquatica</i>	795†

†First observation occurred in 2024.

Pinguicula vulgaris (butterwort) was observed at Jensen Harbor Wetland (1273) but not in any quadrats.

Extra Sites and Data (from 2024 sampling)

Benchmark site East Saginaw Bay Coastal Wetland #5 (515) was sampled on August 8th, 2024. This site was selected again as a benchmark to track long-term trends at a site that was highly degraded throughout earlier long-term sampling. This season, apparent herbicidal treatment of invasive *Phragmites australis* resulted in low species cover and relatively high percentages of standing dead *Phragmites australis* (Figure 32).

Mackinac Creek Wetland (616) was sampled on July 29th–July 30th, 2024. This is a high-quality, well-protected site that has a long history of sampling prior to the beginning of the CWMP, as well as having been regularly sampled as part of CWMP.

Munuscong Lake Wetland #2, #3, Munuscong River Delta (792) was sampled on August 27th, 2024. This site was sampled as part of a restoration project looking at the response of invasive *Typha x glauca* (hybrid cattail) and *Hydrocharis morsus-ranae* (frogbit) to various harvest and treatment approaches.

Point St. Ignace Wetland (1598) was sampled on July 29th–July 30th, 2024, to track the potential environmental changes in the Straits of Mackinac. The only notable change that was observed is that there are several more open pools/patches of *Typha x glauca* and invasive *Phragmites australis*.

Kalamazoo River Wetland (1651) was sampled on June 24th–25th 2024, and Douglas Bayou Wetland (1652) was sampled on June 25h–26th 2024. These sites are being monitored post-herbicide treatment of invasive *Phragmites*.

The pristine Indian Harbor Wetland (7061) benchmark was sampled on July 17th, 2024. The presence of *Myriophyllum spicatum* was first noticed at this site in 2022 and it was observed

again in 2024. When sampled in 2023, there was notation taken on the presence of *Typha x glauca* which was not noted when sampled in 2024.



Figure 32: Comparison between invasive *Phragmites australis* at East Saginaw Bay Coastal Wetland #5 (515B) 2023 (Top) and 2024 (bottom)

Mona Lake - Celery Flats Wetland (7079) was sampled on September 6th, 2024. This wetland was sampled to monitor the reconnection to the Great Lakes and the re-establishment of wetland vegetation to a previously cultivated celery pond, although restoration for this site has not commenced. The crew noted a lack of zonation again this year at this site. Two transects had only an emergent zone and one transect had only a meadow and emergent zone. One of the transects sampled had evidence of mowing invasive *Phragmites australis* and invasive *Lythrum salicaria*. The substrate at this site was coated by a layer of loose sludge and the water clarity was very low. Hardly any submergent plants were

documented at this site.

The data for all these sites will be entered into GreatLakesWetlands.org web site.

New to the sampling program this year was data taken at the start point of each transect. Kendalyn Town, a graduate student at SUNY Brockport under Rachel Schultz is looking at the vegetation composition of the upland edge of the wetland for her thesis project. Crews were given sampling sheets to obtain information on plant species in a quadrat at the start point of each transect. Crews would then collect unidentified plants from each site and transect to be sent to SUNY Brockport for identification by Kendalyn Town. An example quadrat from Kendalyn's upland edge sampling is seen in Figure 33 below.



Figure 33: Upland edge quadrat for Suny Brockport student, Kendalyn Town at Duck Island Area Wetland (736) in 2024

Update:

The three benchmark sites sampled in 2024 with the highest species richness include Indian Harbor Wetland (7061) with 83 taxa, Mackinac Creek Wetland (616) with 63 taxa, and East Saginaw Bay Coastal Wetland #5 (515) with 55 taxa.

Wetland Condition Observations and Results (from 2024 sampling)

The most visible trend noted by sampling crews is that the vegetation zonation was impacted by receding water levels. In many sites, the remains of dead woody plants or *Typha* sp. persist in the wet meadow and upper emergent zones. Some sites however, appear to be establishing

more distinct zonation following the years of fluctuating water levels. For example, at Point aux Chenes Wetland (436), the Central Basin crews noted that the upper emergent zone of Transect 3 had higher coverage values than in 2023 (Figure 34).



Figure 34: Comparison between sample point 6 of transect 3 at Point aux Chenes Wetland (436) in 2023 (Top) and 2024 (Bottom)

Another observation from Dr. Albert while doing plant identification was that there were far more fertile (flowering) plants collected in the meadow zone than in recent high-water years, representing a broad diversity of plants including sedges, rushes, spike rushes, St. John's wort, aster, goldenrods, etc. In contrast, the number of submergent plants collected dropped significantly from recent high-water years.

Data Processing

Dr. Dennis Albert has finished the last of the plant identifications, and data entry will begin in October 2024. All data was entered and quality-checked by February 2025.

Update: All 2024 vegetation data was entered into the GreatLakesWetlands.org database.

Mid-season QC Check Findings (from fall report)

Matthew Sand completed mid-season QC checks. Matt Sand, Emma Waatti, Katlyn Groulx, and their respective crews completed Jensen Harbor Wetland (1273) on July 18th 2024. Each crew completed their own transect at this site. Matt Sand quality-checked Emma's and Katlyn's transects by re-sampling the first quadrat to verify location, percent cover estimates, and plant identification. No corrections to transects were needed for both sampling crews.

Audit and QC Report and Findings

A CMU technician began entering 2024 vegetation data to GreatLakesWetlands.org in October. When the entry was completed, data was quality-checked by Emma Waatti, Matt Sand, and Kim Schraitle. Finally, data was reviewed by Dr. Dennis Albert. There were no data entry issues noted for the Spring 2025 report.

Update: All entered vegetation data was quality-checked by Emma Waatti or Katlyn Groulx, final-checked by Matt Sand or Kim Schraitle, and reviewed by Dr. Dennis Albert.

Additional Funding and Projects

None

Other Collaboration Activities

There are no external collaboration activities to report for the 2024 field season other than the internal collaboration with Kendalyn Town from SUNY Brockport under Rachel Schultz.

CWMP vegetation data and IBIs from 2011-2022 was used to develop Great Lakes status and trends for The State of the Great Lakes 2025 Wetland Vegetation report.

Other Data Requests

Per agreement, a list of species encountered/observed in 2024 will be sent to outside entities to be able to have site access on their conservancies and properties. This includes Detroit River International Wildlife Refuge for Swan Creek Wetland (1915), Little Traverse Conservancy for Mackinac Bay Wetland (617), Duck Island Area Wetland (736), and Mackinac Creek Wetland (616B), and GI Nature and Land Conservancy for Trenton Channel Wetland (426).

Related Student Research

Data from 2023 will be used in a student's research project. Graduate student and crew leader, Emma Waatti, will be using 2023 CWMP vegetation data to assess the effects of road salt on wetland vegetation for her master's thesis research.

CANADIAN CENTRAL/EASTERN BASIN BIRD/ANURAN TEAM AT BIRDS CANADA, PORT ROWAN/LONG POINT, ONTARIO

Team Members

- Dr. Doug Tozer, PI, waterbird and anuran ecologist (since 2011)
- Jeremy Bensette, bird and anuran field crew (since 2014)
- Tim Arthur, bird and anuran field crew (since 2017)
- Tyler Hoar, bird and anuran contractor (since 2011)
- Nadine Litwin, bird and anuran contractor (since 2011)

Training

All four field crew members / contractors will receive training refreshers via Zoom in early April 2025. Topics will include site selection and station placement; anuran and bird survey field protocols; reporting; safety procedures; data entry; and GPS procedures. All members previously showed comprehension of the topics through written and practical in-person tests and successfully completed the online anuran and bird identification tests.

Challenges and Lessons Learned

Field work in 2024 went smoothly with no noteworthy challenges. With all team members having many years of experience working on the project, sampling progressed as planned.

Field work in 2025 looks straightforward for us. We foresee no issues and many of the sites we have visited in multiple previous years.

Site Visit List

We considered 58 sites for sampling in 2024, which consisted of 2 benchmark sites, 7 resample sites, 6 pre-sample sites, and 43 panel sites. We surveyed 42 of the 58 sites for anurans and/or birds. We were unable to survey 16 of the sites due to issues with obtaining landowner access or safety.

We are considering 59 sites for sampling in 2025, which consist of 51 panel, 7 resample, and 1 special-request benchmark sites. We will attempt to survey 47 of these 57 sites. We are unable to survey 10 of the sites due to the following:

- issues with obtaining landowner access (4 sites)
- safety (6 sites)

Panel Survey Results (from fall report)

Sampling for anurans occurred from 14 March until 5 July 2024 and sampling for birds occurred from 20 May to 9 July 2024. Of note were 133 point occurrences of 8 Ontario bird species at risk or of conservation concern.

Species	ON-ESA/SARA Status*	No. Occurrences	
		<u>2023</u> (n = 45 sites)	<u>2024</u> (n = 42 sites)
Bald Eagle	Special concern	14	20
Bank Swallow	Threatened	12	20
Barn Swallow	Threatened	49	43
Black Tern	Special concern	—	12
Bobolink	Threatened	1	—
Chimney Swift	Threatened	7	6
Common Nighthawk	Threatened	2	2
Least Bittern	Threatened	30	28
Red-headed Woodpecker	Endangered	1	2
Total		116	133

*Status is the assessment of greatest concern based on Ontario's Endangered Species Act (ON-ESA) or Canada's Species at Risk Act (SARA).

Also of note were 9 occurrences of Chorus Frog, some populations of which are listed as threatened in Canada (we logged 11 occurrences in 2023).

Extra Sites and Data (from fall report)

We sampled 1 benchmark site in 2024: Hillman Marsh (5422) in Lake Erie.

We collected additional habitat data at each bird and anuran sample point following a slightly modified version of Birds Canada's Great Lakes Marsh Monitoring Program habitat sampling protocol. These data are being collected to augment species-habitat relationship models, especially for certain marsh bird species, some of which are strongly influenced by local vegetation characteristics (i.e., within a few hundred meters of the sampling point), and are stored in an Access database on Birds Canada's secure servers in Port Rowan, Ontario.

Wetland Condition Observations and Results (from fall report)

We sensed that the abundance of secretive marsh birds continued to be relatively high at survey stations with suitable emergent vegetation in 2024 compared to most previous years since 2011. By contrast, these species were absent or at lower abundance at some stations with especially high water where emergent vegetation was relatively sparse. Our observations are reflected in the relationship between lake levels and the number of secretive marsh bird detections over the years, with higher lake levels generally yielding higher abundance of secretive marsh birds (see Figure x). Our observations are also reflected in the results reported by Homan et al. (2021) and Tozer et al. (2024), who used CWMP bird data from throughout the Great Lakes across several years to show that higher water levels generally yield higher wetland bird occurrence and abundance.

Data Processing

All of our data have been entered into and checked in the CWMP database.

Mid-season QC Check Findings

Mid-season checks will be performed in mid-June.

Audit and QC Report and Findings

No issues to report.

Additional Funding and Projects

We received additional funding to augment the bird and anuran team's capacity to complete a 10-year trend analysis for birds, as well as for anurans, using all of the CWMP data from Canada and the US. These projects are described further in the next section.

Other Collaboration Activities

The CWMP bird and anuran team is collaborating with Danielle Ethier, Bird Population Scientist at Birds Canada in Port Rowan, Ontario, to calculate bird and frog trends in coastal wetlands throughout Canada and the U.S. based on CWMP data. The bird paper is now published: <https://doi.org/10.1093/ornithapp/duad062> and the frog paper is in press with Ecosphere. The abstract for the frog paper is included at the end of this section.

The CWMP bird and anuran team is also collaborating with the other CWMP teams on a book entitled “Limnology of Coastal Wetlands Associated with Large Freshwater Lakes.” We are co-authoring the “Wildlife” chapter in the book, which will include various information based on CWMP data. The draft abstract for the chapter is included at the end of this section.

Birds Canada started working with Parks Canada to see if the Index of Biotic Condition (IBC, Howe et al. 2023) for birds and frogs might meet their wetland monitoring and reporting needs at 1) Lake Superior National Marine Conservation Area, 2) Point Pelee National Park (Lake Erie), and 3) Thousand Islands National Park (Lake Ontario). Initial meetings are promising. The ongoing, standardized IBCs for birds and frogs from throughout the Great Lakes based on CWMP data are especially attractive to Parks Canada because they provide an ideal reference within which to place results from these 3 Canadian parks.

Other Data Requests

Nothing to report.

Related Student Research

Nothing to report.




Figure 35. Higher lake levels on lakes Huron, Erie, and Ontario generally yield higher numbers of detections of secretive marsh birds of conservation concern (American Bittern, American Coot, Common Gallinule, Least Bittern, Pied-billed Grebe, Sora, Virginia Rail) by the Canadian Central/Eastern basin bird survey team. This is because rising and higher lake levels inundate emergent vegetation and break up dense stands of emergent vegetation, which is preferred by most of these species. Standardized lake levels were calculated by subtracting the long-term mean for each lake from the annual mean for each lake and dividing by the standard deviation, given the reference value is the same for all lakes (International Great Lakes Datum 1985). Secretive marsh bird detections per station were calculated by summing the number of individuals observed on any of the visits to a station in each year, and dividing by the number of stations surveyed in each year.

Occurrence patterns and trends of frogs in coastal wetlands of the Great Lakes call for further habitat restoration

Douglas C. Tozer^{1*#}, Annie M. Bracey², Valerie J. Brady², Michael F. Chislock³, Jan J. H. Ciborowski⁴, Matthew J. Cooper⁵, Giuseppe E. Fiorino⁶, Thomas M. Gehring⁷, Erin E. Gnass Giese⁸, Greg P. Grabas⁶, Anna M. Harrison⁷, Robert W. Howe⁸, Gary A. Lamberti⁹, Gregory J. Lawrence³, Gerald J. Niemi², Donald G. Uzarski⁷, Bridget A. Wheelock⁷, Danielle M. Ethier^{1#}

Countless wetlands have been lost and degraded globally making amphibians the most threatened vertebrate class. However, despite facing extensive threats and stressors, coastal wetlands of the Laurentian Great Lakes of North America (lakes Superior, Michigan, Huron, Erie, and Ontario) still support sizable populations of frogs (order Anura, including toads). We used data from the Great Lakes Coastal Wetland Monitoring Program to quantify the first-ever annual occurrence probabilities and trends (2011–2023) of eight marsh-breeding frog species, or groups of species, at 1,550 point count locations in 747 coastal wetlands throughout the Great Lakes, and to assess 11 potential drivers of occurrence. Sampled wetlands were marshes greater than 4 ha in area with a permanent or periodic surface-water connection to an adjacent Great Lake or their connecting river systems. Across our study area, green frog (*Lithobates clamitans*) occurrence increased by 8% per year, whereas chorus frog (*Pseudacris maculata*, *P. triseriata*) occurrence decreased by 14% per year. We found more positive or stable trends in occurrence among lakes and species (85%) than negative trends (15%). Occurrence of all species was negatively associated with one or two indicators of poor water quality: specific conductance, ammonium nitrogen, nitrate nitrogen, and urban and agricultural land cover in the surrounding watershed (median area: 12 km²). Occurrence of multiple species was positively associated with high lake levels and surrounding wetland (< 250 m) and forest (< 2.5 km) land cover and negatively associated with surrounding road density (< 2.5 km). Even though occurrence of most species was increasing or stable and was relatively high (> 50%), all will likely benefit from conservation actions. Fifty to 90% of Great Lakes coastal wetlands have been lost and converted to anthropogenic land uses leaving frog populations at a fraction of their former, original sizes. Therefore, extra precaution is critical to help ensure their growth and persistence. Improving water quality, increasing natural forest and wetland land cover within 2.5 km, and reducing roads within 2.5 km of Great Lakes coastal wetlands will help conserve these important indicator species in this globally-recognized but imperiled ecosystem.

Status: in press, Ecosphere.

Limnology of Coastal Wetlands Associated with Large Freshwater Lakes **Wildlife**

Douglas C. Tozer, Annie M. Bracey, Giuseppe E. Fiorino, Erin E. Gnass Giese, Robert W. Howe, Gregory J. Lawrence, Bridget A. Wheelock, Thomas M. Gehring

Abstract Coastal wetlands of large lakes (CWLLs) are used by numerous wildlife species, including birds, mammals, reptiles, and amphibians. Our main purpose in this chapter is to explore what makes CWLLs unique among wetlands with respect to wildlife use and conservation. We pursue the following objectives for wildlife in CWLLs: 1) describe diversity; 2) overview influential factors; 3) explore a

selection of conservation solutions; and 4) discuss areas for further research and a call to action. We highlight that high-quality wildlife habitat is created, maintained, and influenced in CWLLs by 1) waves, seiches, and ice scour; 2) fluctuating water levels; 3) littoral sediment transport; and 4) unique hydrogeomorphic characteristics—all of which are stronger in CWLLs, or only occur in CWLLs, compared to other wetlands. Given their high value for supporting healthy wildlife populations, we stress that CWLLs should be given priority for permanent protection, restoration, and conservation, including the implementation of conservation measures and actions that facilitate resilient and adaptable wetlands in the face of future extremes due to climate change.

CANADIAN CENTRAL BASIN FISH, INVERTEBRATE AND WATER QUALITY TEAM AT THE UNIVERSITY OF WINDSOR AND UNIVERSITY OF WISCONSIN RIVER FALLS

Team Members

- Dr. Jan Ciborowski (UW), PI, aquatic ecologist, (since 2011)
- Dr. Joseph Gathman (UWRF), co-PI, aquatic ecologist, team leader (since 2011)
- Li Wang (UW), GIS specialist, data/QC manager (since 2011)
- Michelle Dobrin (UW), lead invertebrate taxonomist (since 2011)
- Stephanie Johnson (UW), crew leader and lab team member (since 2016)
- Emilee Mancini (UW), field and lab crew member (since 2020)
- Julia Santini (UW), field and lab crew member (since 2023)

Training

All crew members in 2025 will have at least two years of experience on the project. Most have many years of experience. Refresher training will be carried out at University of Windsor in early June under the supervision of PI Joseph Gathman and experienced crew member Stephanie Johnson. All field crew members will review updates to the QAPP and SOP documents, and receive instruction in GPS use, assessment of site-suitability criteria (open water connection to lake, presence of a wetland, safe access), identification of vegetation zones to be sampled, water quality-sample collection, preprocessing and shipping samples to water quality labs, calibrating and reading field instruments and meters, setting, removing, cleaning and transporting fyke nets, cleaning and transporting boats and other sampling equipment, and protocols for collecting and preserving macroinvertebrates. All field-crew members have been certified for identifying common fishes and Species-at-Risk through the Royal Ontario Museum's course in fish identification in 2023 or earlier.

The crew leader in 2025 will be co-PI Joseph Gathman, who has been a PI since the beginning of the CWM program and has been identifying fish and macroinvertebrates for three decades. He will lead the crew in the field at almost all sites, except at a few local (close to Windsor) sites where the field crew might be led by Stephanie Johnson, who has many years of field-crew supervision experience in the CWM project. Gathman will also prescreen the suitability of sample sites, coordinate all logistics, secure accommodations, and obtain sampling permissions where necessary.

Challenges and Lessons Learned

We have nothing new to report here based on our 2024 experiences. As was the case in 2023 and 2024, water levels in Lake Erie and Lake Huron are anticipated to continue their multi-year decline, so we do not expect to face the high-water challenges of earlier years, meaning that it is unlikely that we will have to employ the deep-water sampling procedures that were followed during the recent high-water years.

Site Visit List

We anticipate sampling 30 sites during the 2025 field season, one of which is a benchmark site (5422 - Hillman Marsh). Four sites will be re-sample sites that were sampled in 2024, and five will be pre-sample sites (sites that will be re-sampled in 2026). Three sites will require permits to be issued prior to sampling: site 5702 is an Ontario Provincial park and will require a scientific-collection permit; site 5422 is a conservation area requiring a permit from the Essex Region Conservation Authority; and site 6001 is a private hunt club requiring special permission for entry. A few sites assigned to us are sites that have been inaccessible in the past, so in lieu of these we will sample a few Lake Ontario sites that were assigned to the Canadian Wildlife Service based in Toronto, as we have done several times in previous years. As in previous years, we will sample Hillman Marsh as a benchmark site to support the monitoring work of the Essex Region Conservation Authority, which has designated this site as a Conservation Area.

Panel Survey Results (from fall report)

Field sampling began on June 23 at site 5585, MacGregor Point Wetland 2, and was finished up in the first week of September at site 5422, Hillman Marsh.

Fishes collected over the summer largely consisted of the usual species. We caught two fish Species-At-Risk (SAR) in Canada: one spotted gar (*Lepisosteus oculatus*), collected at site 5304, Flat Creek Wetland, and one lake chubsucker (*Erimyzon sucetta*) at site 5831, Rondeau Provincial Park (both sites were on the Canadian shore of Lake Erie. Regarding non-native species, we caught relatively few common carp (*Cyprinus carpio*) or goldfish (*Carassius auratus*) compared to most previous years, but found round gobies (*Neogobius melanostomus*) and

tubenose gobies (*Proterorhinus semilunaris*) at several sites. Reptiles observed included many painted turtles (*Chrysemys picta*), a few snapping turtles (*Chelydra serpentina*), and several musk turtles (*Sternotherus odoratus*), plus one northern map turtle (*Graptemys geographica*). Northern water snakes (*Nerodia sipedon*) were observed at multiple sites.

As in 2023, as compared to many earlier years we had an unusually large number of samples with relatively few invertebrates, i.e., fewer samples of 150 individuals could be collected within the prescribed time than is usually the case. Non-native invertebrates collected included a number of zebra mussels and the snail, *Bithynia tentaculata*, at several wetlands and one occurrence of the amphipod, *Echinogammarus ischnus*, (at site 5574, Lypps Beach), on the Canadian shore of Lake Erie. The relatively large number of zebra mussels collected in dip-net samples may have been a result of the earlier high-water years which may have allowed the mussels to colonize the wetlands.

Extra Sites and Data (from fall report)

Point Pelee Marsh 2 (5762), in Point Pelee National Park (Ontario, Canada), was sampled for the sixth time in 2024 as a benchmark site in cooperation with Parks Canada, which is conducting a multi-year restoration project to increase the amount of open-water area at Point Pelee. In 2018, the barrier beach that protected the marsh broke open during a series of strong storms/seiche events, and the breach had remained open until 2022. Park personnel are interested in our monitoring changes as they proceed with their restoration work as well as changes resulting from the breach and recreation of the barrier beach. The water in the marsh has been too deep for fish sampling for the last six years, so we collected only water quality samples/data and invertebrate samples.

The other benchmark site in 2024, Hillman Marsh, is a large wetland near Point Pelee, which is overseen by the Essex Region Conservation Authority (ERCA). We have frequently sampled this wetland in collaboration with ERCA personnel to assist them in tracking changes over the years. In 2024, we sampled macroinvertebrates and water quality at the marsh. Fishes were not sampled because the water was not deep enough to allow us to set fyke nets.

Wetland Condition Observations and Results (from fall report)

We observed little year-to-year change in Lake Huron and Lake Erie water levels in 2024. Wet meadow conditions were similar to 2023 and most were too shallow/unflooded to allow us to sample them for fish. Also, many previously devegetated areas (resulting from high water in earlier years) had still not yet recovered their vegetation, making them unsuitable for sampling.

Data Processing

All data have been entered into the database and QC checked and all data-verification requests have been completed. QC swaps of invertebrate samples with other teams have also been completed.

Mid-season QC Check Findings

This will be conducted mid-season in 2025.

Audit and QC Report and Findings

Nothing new to report.

Additional Funding and Projects

None to report.

Other Collaboration Activities

The Ciborowski lab at U. Windsor will continue its collaboration with the Essex Regional Conservation Authority in monitoring the changing conditions at Hillman Marsh (site 5422), Essex County, Ontario, as restoration work continues there. We will continue to interact with staff at Point Pelee National Park as opportunities arise.

Four other University of Windsor faculty, Drs. Ken Drouillard, Mike McKay, Chris Weisener, and Catherine Febria, will begin collaborations with the Ciborowski lab. We anticipate that these interactions will broaden the scope of our lab's work and allow individuals from our lab and theirs to work together on CWM and other projects.

Other Data Requests

Nothing to report.

Related Student Research

Nothing to report.

CANADIAN CENTRAL BASIN VEGETATION TEAM AT THE UNIVERSITY OF WINDSOR AND UNIVERSITY OF WISCONSIN RIVER FALLS

Team Members

- Dr. Jan Ciborowski (UW), PI, aquatic ecologist (since 2011)
- Dr. Joseph Gathman (UWRF), co-PI, aquatic ecologist, team leader (since 2011)
- Carla Huebert (UW), crew leader, plant taxonomist (since 2013)
- Li Wang (UW), GIS specialist, data/QC manager (since 2011)

Training

Carla Huebert has conducted vegetation sampling for the project since 2013 so only a review of protocols will be needed as outlined in the QAPP. The review in May of 2025 will include instruction in GPS use, assessment of site selection criteria (open water connection to lake, presence of a wetland, safe access), and identification of vegetation zones to be sampled. Carla will also receive refresher training and review in field data and lab entry to become familiar with changes to the database. Joseph Gathman and Carla will preview sites to be sampled and agree on sampling strategies for each site, as well as ensuring that they are up to date on any new non-native species that may be encountered.

Challenges and Lessons Learned

We have nothing new to report here based on our 2024 experiences. As was the case in 2023 and 2024, water levels in Lake Erie and Lake Huron are anticipated to continue their multi-year decline, so we do not expect to face the high-water challenges of earlier years. Further, we expect that vegetation will continue to fill in areas that had been cleared of vegetation by high lake levels, so we are likely to return to more “normal” conditions and procedures.

Site Visit List

We anticipate sampling 30 sites during the 2025 field season, one of which is a benchmark site (5422 - Hillman Marsh). Four sites will be re-sample sites that were sampled in 2024, and five will be pre-sample sites (sites that will be re-sampled in 2026). Three sites will require permits to be issued prior to sampling: site 5702 is an Ontario provincial park and will require a scientific-collection permit; site 5422 is a conservation area requiring a permit from the Essex Region Conservation Authority; and site 6001 is a private hunt club requiring special permission for entry. A few sites assigned to us are sites that have been inaccessible in the past, so in lieu of these we will sample a few Lake Ontario sites that were assigned to the Canadian Wildlife Service based in Toronto, as we have done several times in previous years. As in previous years,

we will sample Hillman Marsh as a benchmark site to support the monitoring and restoration work of the Essex Region Conservation Authority, which has designated this site as a Conservation Area.

Panel Survey Results (from fall report)

Water levels continued their downward trend in 2024 in the basin areas sampled by the team, including in Lake Huron's Bruce Peninsula, where the lower water levels revealed several rare coastal alvar meadows at two of our sites, Fishing Islands 11 (5286), and Sadler Creek Wetland 1 (5844). When these wetlands were last visited during the previous cycle (2019), these alvars were underwater, and had little to no vegetation growing at that time. With several years now of drier, exposed land, these alvars have had a chance to regenerate their unique plant communities, and many uncommon and rare species were surveyed, including dwarf lake iris (*Iris lacustris*), small yellow lady's slipper (*Cypripedium parviflorum*), bulrush sedge (*Carex scirpoidea*), Canada bluets (*Houstonia canadensis*), and Arctic sweet coltsfoot (*Petasites frigidus*).

Another highlight to report for 2024 was the Wainfleet Wetland (6016), located in Lake Erie's eastern basin. The site is unusual in the Lake Erie basin for having exposed Niagara escarpment limestone substrate throughout the wet meadow zone, which has in turn, created a unique alvar meadow, something rather rare for our team to find within a Lake Erie wetland. Species that our team usually only observes at our rockier Lake Huron sites, such as golden sedge (*Carex aurea*), Crawe's sedge (*Carex crawei*), and upland white goldenrod (*Solidago ptarmicoides*), were observed at the Wainfleet wetland. Other uncommon species found here include Torrey's rush (*Juncus torreyi*), gray goldenrod (*Solidago nemoralis*), and false dragon-head (*Physostegia virginiana*).

An interesting observation noted by the crew from the Wainfleet wetland wet meadow is how this unique alvar formation has prevented invasive *Phragmites australis* from overtaking the meadow, as it most often will do in a sediment-rich wetland. It appears that, because there is so little organic matter on top of the limestone substrate at Wainfleet, *Phragmites* cannot gain a foothold there. Also, since the entire meadow and site are both surrounded by a thick wall of *Phragmites* that has not been treated or sprayed since the site was first sampled by our team ten years ago in 2014, it seems highly plausible that by 2024, it would have easily overtaken this diminutive wet meadow, had it been able to do so.

Another surprisingly diverse site came from another Lake Erie wetland, Wheatley East Two Creeks (6054), located in the lower central basin. The year 2024 marked the first year that a CWMP crew has visited this site. While there, the veg team encountered several species that they have either seldom or never encountered before in their 12 years of CWMP experience, including hop sedge (*Carex lupulina*), ditch stonecrop (*Penthorum sedoides*), and Lake Ontario aster (*Symphyotrichum ontarionis*). All of these species were found growing in typical Lake Erie mucky clay substrate.

Invasive species:

Creeping Water Primrose (*Ludwigia peploides*): Was first discovered by the UWindsor team in 2023 at two of our western Lake Erie sites. One of those sites, Lypps Beach Wetland (5574) was chosen as a resample site in 2024 to monitor any expansion of this aggressive new species. When the crew returned to the site in 2024, they noted that creeping water primrose had not only successfully overwintered but had also rapidly spread throughout the wetland. In 2023, this species was found in large clusters, primarily on the western side of the wetland. However, in 2024, it was found covering the majority of the wetland edge margins and beyond. It was also observed to have increased in density, with its creeping stems forming a mat so dense that one had to walk upon it rather than through it.

Elecampane (*Inula helenium*): Elecampane (Figure 3) was found for the first time by our team at one of our eastern Lake Ontario sites, Hay Bay Marsh 7 (5406). It was found growing in the upper wet meadow zone in several locations, with some plants there reaching over two meters in height. While it has not been observed at any of our other sampled wetlands, it seemed to be relatively well established in that area of Lake Ontario, with the crew observing it growing in ditches and lower lying areas whilst travelling through the area by vehicle.

European Frog-bit (*Hydrocharis morsus-ranae*): In addition to being recorded at several Lake Erie sites where Frog Bit is a usual occurrence, it was found again at a St. Mary's River site along Lake Huron's north channel, Everen's Point (5276). It had been found there in the previous cycle year of 2019, and it was present again 2024. However, the positive news is that the plant has not invaded any new areas of the wetland, and in 2024 is still confined to a small, shallow pocket near the boat launch.

Species at risk:

American water-willow (*Justicia americana*) was observed again in 2024 at our benchmark site, Point Pelee Marsh 2 (5762). It had also been found in several areas of the site throughout the six years the site has been sampled, beginning in 2019. (Federal COSEWIC Status: Threatened).

Dwarf lake iris (*Iris lacustris*) was found for the first time at one of our Lake Huron Bruce Peninsula sites, Sadler Creek Wetland 1 (5844). It was observed growing in several locations on the alvar meadow and was observed in three quadrats at the site. (Federal COSEWIC Status: Special Concern).

Swamp rose mallow (*Hibiscus moscheutos*) was found at four Canadian sites in 2024: Flat Creek Wetland (5304), Rondeau Provincial Park Wetland (5831), Hillman Marsh (5422), and Point Pelee Marsh (5762). (Federal COSEWIC Status: Special Concern).

Extra Sites and Data

We did not sample any extra sites in 2024.

Wetland Condition Observations and Results (from fall report)

The high-water years from 2018 to 2021 had a negative effect on the forest edge and understory of many wetlands, with the sustained high waters drowning out the wet meadow zone, and fully flooding the forest edge habitat. Depending on the location, this sustained flooding pushed into the forest and woody areas by 75 meters or more.

However, an interesting phenomenon was observed by our team at several wet meadows sampled throughout the basin in 2024. After several years, the lower-elevation trees and understory shrubs have finally fully succumbed to the sustained flooding, including species of birch (*Betula sp.*), alder (*Alnus sp.*), cedar (*Thuja sp.*), and maple (*Acer sp.*). The flooding, in combination with the Emerald Ash Borer disease having recently reached into the Lake Huron north channel area of Ontario, has also sped up the death of many of the nearshore ash tree species (*Fraxinus sp.*).

With the death of so many trees, the formerly shaded forest canopy has now transitioned into an open, full sun zone with plenty of moist, organic matter. Many of the standing dead tree limbs and trunks have now fallen, exposing this area to even more sun. After water levels began to recede in 2021, wet meadow vegetation has slowly expanded into these new open area zones, increasing the length of the wet meadow zone by up to 75 meters or more. An example of this was at one of our sites on the St. Mary's River, in Lake Huron's north channel, Everen's Point (5276).

If water levels remain low for the next several years, the forest edge and understory will likely begin to regenerate and slowly shade out, and subsequently, push out the wet meadow once again, but for now, the wet meadow zone at several of our sites has dramatically increased in size. Figure 39 shows an example of one of these newly expanded wet meadows.

Data Processing

All vegetation data and GPS waypoints have been entered into the database.

Mid-season QC Check Findings

These will be conducted in mid-season.

Audit and QC Report and Findings

Completed for 2023 data.

Additional Funding and Projects

None to report.

Other Collaboration Activities

Nothing to report.

Other Data Requests

Nothing to report.

Related Student Research

No additional projects to report.

CANADIAN EASTERN BASIN FISH, INVERTEBRATE AND WATER QUALITY TEAM AT CANADIAN WILDLIFE SERVICE

Team Members

- Joe Fiorino, PI, wetland ecologist (since 2016)
- Ian Smith, crew leader, fish sampling, GIS tech (since 2014)
- Hayley Rogers, team leader, vegetation sampling (since 2017)
- Patrick Rivers, team leader, WQ/invert sampling (intermittent since 2014)
- Albert Garofalo, field crew member, vegetation sampling (intermittent since 2018)
- Marissa Zago, field crew member, vegetation/fish/WQ/invert sampling (2 years, 2018, 2023)
- Kyra Riley, summer student field tech, vegetation/fish/WQ/invert sampling (2024)
- Whistler Brown, summer student field tech, vegetation/fish/WQ/invert sampling (2024)

Training

Environment and Climate Change Canada – Canadian Wildlife Service (ECCC-CWS) field crew members were trained by Joe Fiorino, Ian Smith, and Hayley Rogers. The sampling protocol, use of technical equipment, occupational health and safety, and field-based decision-making were covered in detail over multiple days; staff were assessed in the field and lab for proper sample collection, data recording, GPS use, water processing, equipment calibration, and lab sample preparation and storage. A practice session at a nearby wetland and in our lab facility was conducted in July 2024 to provide hands-on training to new staff. An experienced staff member was paired with new personnel to reinforce project protocols and ensure high data quality.

Trainings will be similarly conducted in 2025.

Challenges and Lessons Learned

None

Site Visit List (from fall report)

As in previous years, the number of sites originally assigned to our group exceeded the capacity of the ECCC-CWS field crew, so four sites were given to SUNY-Brockport (5008, 5196, 5635, 5806) and six sites were given to University of Windsor (5131, 5256, 5259, 5402, 5406, 6031).

Ten sites were sampled and one site was a “web reject” because it is not reasonably accessible by boat or canoe (5394). Fish sampling was not possible at two sites. The open water section of Rattray Marsh (5799) is typically accessed through a narrow channel, but it has filled in with cattails in recent years. The crew was able to portage with canoes to access the site for water quality, inverts, and vegetation sampling, but there was no reasonable/safe way to transport fyke nets. Wellers Bay 4 (6037) is a former training site used by the Department of National Defence, and penetrating the substrate (which is required to set fyke nets) is not permitted due to the risk of Unexploded Explosive Ordnance (UXO).

Panel Survey Results (from fall report)

Sampling occurred August 6, 2024 to August 23, 2024. All water quality (except chlorophyll-a; waiting on lab results), fish, and invertebrate data have been received, entered into the database, and QC'ed.

Reptiles:

Painted Turtle (*Chrysemys picta*) were caught at Bayside Wetland 2 (5064; 2 individuals), Colborne Creek (5179; 4 individuals), Salmon River Wetland (5562; 5 individuals), and Wellers Bay Wetland 14 (6034; 1 individual). Musk Turtle (*Sternotherus odoratus*) were caught at Salmon River Wetland (5562; 3 individuals).



Figure 36. Musk Turtle caught at Salmon River Wetland (5562).

Rare species:

Grass Pickerel (*Esox americanus vermiculatus*), a species of Special Concern status in Ontario, was caught at Carrying Place (5161; 1 individual).

Non-native species:

Round Goby (*Neogobius melanostomus*) were caught at Bayside Wetland 2 (5064; 61 individuals), Sawguin Creek Marsh 8 (5876; 105 individuals), Wellers Bay Wetland 14 (6034; 10 individuals), and West Lake Wetland 5 (6048; 4 individuals). Common Carp (*Cyprinus carpio*) were caught at Colborne Creek (5179; 3 individuals).

Other notes: Rattray Marsh (5799) has undergone significant restoration over the last decade. A previously installed carp exclusion fence was recently revitalized and appears to have resulted in improved marsh condition. The section of the marsh to the west of the fence appeared to have lower turbidity and supported a relatively dense population of lilies, whereas the rest of the site supported little aquatic vegetation.

Extra Sites and Data (from fall report)

No benchmark sites were sampled.

Continued to collect data on short-term variation in dissolved oxygen and water levels for Dr. Jan Ciborowski (University of Windsor). These data are managed by Dr. Ciborowski's lab.

Continued to collect dissolved gas samples, air samples, and porewater samples for Amanda Suchy (Central Michigan University). These data are managed by Amanda Suchy.

Wetland Condition Observations and Results (from fall report)

Water levels in August on Lake Ontario were consistent with the long-term average. Nothing else to add beyond what was mentioned in the Panel Survey Results above.

Data Processing

All water quality, fish, and invert data have been entered and QC'ed. Geospatial data were mapped in GIS to verify that waypoints were recorded correctly.

Mid-season QC Check Findings

No difficulties or anomalies were observed during mid-season checks.

Audit and QC Report and Findings

All data entry has been QC'ed by a team member with multiple years of experience working with the data entry system. The majority of QC issues identified between 2016 and 2022 in the Data Verification Interface have been addressed, as well as past point-matching issues.

Additional Funding and Projects

Since 2021, ECCC-CWS has worked with the International Joint Commission to update marsh bird ecological performance indicators used for adaptive management of outflow regulation on Lake Ontario. ECCC-CWS received support from the bird/anuran team in December 2021 to conduct an analysis using CWMP data, and ultimately identified six potential bird-based indicators for consideration by the IJC. This work was published in the Journal of Great Lakes Research in early 2023. Since then, CWS has worked with the Hydrodynamic and Ecohydraulic Section of the National Hydrological Service to utilize spatially-explicit, model-generated data pertaining to water depths, flooding patterns, and habitat extent and structure at wetland sites on Lake Ontario to develop predictive models of marsh bird abundance and richness. These predictive models are currently being used as ecological performance indicators for the ongoing expedited review of the current water-level regulation plan (Plan 2014). The associated manuscript is currently being prepared for publication.

Other Collaboration Activities

ECCC-CWS funded a project on trends in anuran populations in collaboration with Birds Canada and various CWMP PIs and team members. Birds Canada used contemporary statistical techniques to assess trends in anuran populations in the Great Lakes basin. The primary source

of data was the Great Lakes Coastal Wetland Monitoring Program (including data from 2011 to 2022). The associated manuscript has been accepted for publication in Ecosphere.

Other Data Requests

In October 2023, Central Lake Ontario Conservation Authority requested water quality data (temperature, pH, turbidity, conductivity) for all Lake Ontario sites (2011 to 2022). These data will be used to calculate water quality index scores for CWMP sites to provide a reference for comparison to local sites.

Related Student Research

None at this time.

CANADIAN EASTERN BASIN VEGETATION TEAM AT CANADIAN WILDLIFE SERVICE

Team Members

- Joe Fiorino, PI, wetland ecologist (since 2016)
- Ian Smith, crew leader, fish sampling, GIS tech (since 2014)
- Hayley Rogers, team leader, vegetation sampling (since 2017)
- Patrick Rivers, team leader, WQ/invert sampling (intermittent since 2014)
- Albert Garofalo, field crew member, vegetation sampling (intermittent since 2018)
- Kyra Riley, summer student field tech, vegetation/fish/WQ/invert sampling (2024)
- Whistler Brown, summer student field tech, vegetation/fish/WQ/invert sampling (2024)

Training

Environment and Climate Change Canada – Canadian Wildlife Service (ECCC-CWS) field crew members were trained by Joe Fiorino, Ian Smith, and Hayley Rogers. The sampling protocol, use of technical equipment, occupational health and safety, and field-based decision-making were covered in detail over multiple days; staff were assessed in the field for GPS use, measuring and spacing of transects, filling out datasheets properly, ensuring species coverages were recorded correctly and standardized, and collecting and taking notes for unknown plant specimens. A practice session at a nearby wetland and in our lab facility was conducted in July 2024 to provide hands-on training to new staff. An experienced staff member was paired with new

personnel to reinforce project protocols and ensure high data quality. Training for 2025 will be conducted similarly.

Challenges and Lessons Learned (from fall report)

None

Site Visit List (from fall report)

As in previous years, the number of sites originally assigned to our group exceeded the capacity of the ECCC-CWS field crew, so four sites were given to SUNY-Brockport (5008, 5196, 5635, 5806) and six sites were given to University of Windsor (5131, 5256, 5259, 5402, 5406, 6031). Ten sites were sampled and one site was a “web reject” because it is not reasonably accessible by boat or canoe (5394).

Site list and status (Environment and Climate Change Canada – Canadian Wildlife Service).

Site ID	Site Name	Site Status
5064	Bayside 2	Sampled
5161	Carrying Place	Sampled
5179	Colborne Creek	Sampled
5394	Gull Pond	Web reject
5562	Salmon River Wetland	Sampled
5795	Hendrie Valley	Sampled
5799	Rattray Marsh	Sampled
5876	Sawguin Creek Marsh 8	Sampled
6034	Wellers 14	Sampled
6037	Wellers Bay 4	Sampled
6048	West Lake 5	Sampled

Panel Survey Results (from fall report)

Sampling occurred August 6, 2024 to August 23, 2024. All data have been entered into the database and QC'ed.

Non-native species:

Typha x glauca dominates most wetlands on Lake Ontario. Many invasive species are common (e.g., *Hydrocharis morsus-ranae*, *Myriophyllum spicatum*, *Lythrum salicaria*, *Nitellopsis obtusa*, *Phalaris arundinacea*). Less common invasive species that were observed in 2024 included

Impatiens glandulifera (Colbourne Creek 1), *Lysimachia nummularia* (Hendrie Valley) and *Rhamnus cathartica* (Wellers Bay 4).

Other notes: Rattray Marsh (5799) has undergone significant restoration over the last decade. A previously installed carp exclusion fence was recently revitalized and appears to have resulted in improved marsh condition. The section of the marsh to the west of the fence appeared to have lower turbidity and supported a relatively dense population of lilies (Figure 37), whereas the rest of the site supported little aquatic vegetation.



Figure 37. Section of Rattray Marsh (5799) to the west of the carp exclusion fence.

Extra Sites and Data (from fall report)

No benchmark sites were sampled.

An extra quadrat was sampled at the upland edge of most transects for Kendalyn Town (SUNY-Brockport). These data are managed by Kendalyn Town.

Wetland Condition Observations and Results (from fall report)

Water levels on Lake Ontario were consistent with the long-term average. Nothing else to add beyond what was mentioned in the Panel Survey Results above.

Data Processing

All data have been entered and QC'ed. Geospatial data for vegetation transects were mapped in GIS to verify that quadrat waypoints were recorded correctly and that transect direction, spacing, and widths were accurate.

Mid-season QC Check Findings

No difficulties or anomalies were observed during mid-season checks. Checks will be conducted mid-season 2025.

Audit and QC Report and Findings

All data entry has been QC'ed by a team member with multiple years of experience working with the data entry system. QC issues identified between 2016 and 2022 in the Data Verification Interface and past point-matching issues have been addressed.

Additional Funding and Projects

Since 2021, ECCC-CWS has worked with the International Joint Commission to update marsh bird ecological performance indicators used for adaptive management of outflow regulation on Lake Ontario. ECCC-CWS received support from the bird/anuran team in December 2021 to conduct an analysis using CWMP data, and ultimately identified six potential bird-based indicators for consideration by the IJC. This work was published in the Journal of Great Lakes Research in early 2023. Since then, CWS has worked with the Hydrodynamic and Ecohydraulic Section of the National Hydrological Service to utilize spatially-explicit, model-generated data pertaining to water depths, flooding patterns, and habitat extent and structure at wetland sites on Lake Ontario to develop predictive models of marsh bird abundance and richness. These predictive models are currently being used as ecological performance indicators for the ongoing expedited review of the current water-level regulation plan (Plan 2014). The associated manuscript is currently being prepared for publication.

Other Collaboration Activities

ECCC-CWS funded a project on trends in anuran populations in collaboration with Birds Canada and various CWMP PIs and team members. Birds Canada used contemporary statistical techniques to assess trends in anuran populations in the Great Lakes basin. The primary source of data was the Great Lakes Coastal Wetland Monitoring Program (including data from 2011 to 2022). The associated manuscript has been accepted for publication in Ecosphere.

Other Data Requests (from fall report)

In September 2024, Meteorological Services of Canada requested quadrat-level plant data (species, cover, GPS coordinates) for US and Canadian sites for all years and lakes except Michigan (2011 to 2023). Their group will use these data to assist in the calibration of a model that predicts coastal wetland vegetation response to key physical variables (e.g., bathymetry, water depths, currents, wet-dry cycles). This will be used in the context of a study to assess the vulnerability of Great Lakes coastal wetlands to climate change as part of the Great Lakes Freshwater Ecosystem Initiative. They also requested to share these data with a contractor that they are hiring to develop a remote sensing classification algorithm for wetland vegetation.

Related Student Research

A junior staff member is currently working on a project investigating spatial and temporal trends of *Nitellopsis obtusa* in the Great Lakes.

US EASTERN BASIN BIRD AND ANURAN TEAM AT SUNY BROCKPORT

Team Members

- Dr. Kathryn Amatangelo, PI (since 2014)
- Dr. Kristen Malone, bird/anuran PI (2023)
- Matt Silverhart, acting fish PI, project manager, crew leader (since 2020)
- Alexa Lashway, graduate research assistant and bird team lead (since 2024)
- William Sidore, undergraduate research assistant and anuran team lead (since 2023)

Training

Both field technicians (Alexa Lashway and William Sidore) were trained by PI Dr. Kristen Malone and project manager Matthew Silverhart on proper field sampling techniques, data collection and recording, GPS use, and field safety. Both technicians were accompanied by PI Dr. Kristen Mallone and project manager Matthew Silverhart for the first Bird and Anuran samplings of the season to ensure proper sampling techniques and train the technicians. Anuran training and observation took place May 1, 2024 at site 15: Yanty Marsh. Bird training and observation took place June 3, 2024 at site 15: Yanty Marsh. Lastly, both field technicians were trained in data entry and QC checks in the database by project manager Matthew Silverhart. Both field technicians were successfully trained, passed the Bird (Alexa Lashway) and Anuran (William

Sidore) identification exams, and met pre-season training performance criteria described in the project QAPP.

Training for 2025 will be conducted similarly.

Challenges and Lessons Learned (from fall report)

Welcoming a new Project Manager, Matthew Silverhart, welcomed a large range of new challenges. Having not previously been a part of the Bird/Anuran sampling, there was a steep learning curve when it came to understanding how this sampling team works most efficiently. With help from PI Dr. Kristen Malone and previous field technicians, the season was able to be organized well and went smoothly.

Site Visit List (from fall report)

Of the 24 assigned sites for the Bird/Anuran team of SUNY Brockport, 23 were sampled in full and 1 could not be accessed (either due to physical barriers or lack of ability to sample safely. 18 of the assigned sites were panel sites, 2 were resample sites from the previous year, 2 were presample sites for the following season, and 3 were benchmark sites (site 28 was both a benchmark and a resample site). All benchmark sites were requested by SUNY Brockport PIs due to restoration projects either being planned, ongoing, or having previously occurred at the specified sites. This information can be used to better inform and shape restoration efforts.

Panel Survey Results (from fall report)

Sampling of panel sites for anurans began on 1 May 2024 at site 15: Yanty Marsh and concluded on 9 July 2024 at site 137: Ray Bay Marsh. During the anuran sampling of panel sites, six species of anurans were observed. Those species were American Toad, Bullfrog, Chorus Frog, Gray Treefrog, Green Frog, and Spring Peeper. The Chorus Frog was only observed one time during our sampling at site 86: Sterling Creek Wetland. The species observed most frequently were the Bullfrog, Gray Treefrog, and Green Frog.

Sampling of panel sites for birds began on 3 June 2024 at site 15: Yanty Marsh and concluded on 9 July 2024 at site 137: Ray Bay Marsh. Table 4 shows results of bird sampling from panel sites. During the bird sampling of panel sites, 77 species of birds were observed and seven of those species were classified as unidentified. Four of the bird species at panel sites are listed as threatened, one species is listed as a species of special concern, and one species is listed as a high priority species of greatest conservation needed by the New York Department of Environmental Conservation (NYDEC). The threatened species are Bald Eagle, Common Tern, Northern Harrier, and Pied-billed Grebe. The species of special concern is the Osprey and the

high priority species of greatest conservation needed is the Bobolink. At most panel sites, more species of bird were observed in the AM sampling period than the PM sampling period. The most common species observed at panel sites were the Red-winged Blackbird, Yellow Warbler, American Robin, and Swamp Sparrow.

Extra Sites and Data (from fall report)

Sampling of benchmark sites for anurans began on 8 May 2024 at site 7052: Braddock Bay and concluded on 27 June 2024 at site 28: Salmon Creek. During the anuran sampling of benchmark sites, six species of anurans were observed. Those species were American Toad, Bullfrog, Gray Treefrog, Green Frog, Northern Leopard Frog, and Spring Peeper. The Northern Leopard Frog was only observed one time during our sampling at site 7052: Braddock Bay. The species observed most frequently were the Bullfrog, Gray Treefrog, and Green Frog.

Sampling of benchmark sites for birds began on 4 June 2024 at site 7052: Braddock Bay and concluded on 26 June 2024 at site 28: Salmon Creek. Table 6 shows results of bird sampling from benchmark sites. During the bird sampling of benchmark sites, 46 species of birds were observed and four of those species were classified as unidentified. Four of the bird species at benchmark sites are listed as threatened, one species is listed as a species of special concern, and one species is listed as a high priority species of greatest conservation needed by the New York Department of Environmental Conservation (NYDEC). The threatened species observed were Bald Eagle, Common Tern, Northern Harrier, and Pied-billed Grebe. The species observed of special concern was the Osprey and the high priority species of greatest conservation needed observed was the Bobolink. At most benchmark sites, more species of bird were observed in the AM sampling period than the PM sampling period.

Wetland Condition Observations and Results (from fall report)

With most sampling points being accessible from nearby parks and or roads, there were not major impacts from changing wetland conditions.

Data Processing

All data collected during 2024 GLCWMP Bird/Anuran sampling has been entered and checked. The habitat forms were collected during bird sampling and were mailed to Doug Tozer on 25 September 2024. Digital copies have been created for backup.

Mid-season QC Check Findings

Mid-season QC check for the Bird/Anuran team occurred at site 15 on 2 June 2024 with PI Dr. Kristen Malone administering the mid-season QC check. The crew members involved were Alexa Lashway and William Sidore. The crew performed all tasks to satisfaction and there were no issues noted that needed to be addressed.

Mid-season checks will be conducted similarly in 2025.

Audit and QC Report and Findings

Other than minor data entry errors, there were no large-scale errors of note for the Bird/Anuran data entry and QC.

Additional Funding and Projects

None

Other Collaboration Activities

None

Related Student Research

No student research coincided with Bird/Anuran sampling this season.

US EASTERN BASIN FISH, INVERTEBRATE, AND WATER QUALITY TEAM AT SUNY BROCKPORT

Team Members

- Dr. Kathryn Amatangelo, PI (since 2014)
- Dr. Michael Chislock, PI (since 2018)
- Matthew Silverhart, acting fish PI, project manager, field crew leader (since 2020)
- Dillon Vandemortel, graduate research assistant (since 2023)
- Madelynn Edwards, invertebrate laboratory technician (since 2019)
- Brayden Link, graduate research assistant (since 2024)
- Kai Schedel, undergraduate research assistant (since 2024)
- Grace Trebilcock, undergraduate research assistant (since 2024)

Training

All field technicians were trained by Project Manager Matthew Silverhart on proper field sampling techniques, lab data collection and recording, GPS use, boat use and safety, fish identification, fyke net operation, macroinvertebrate collection and storage, and data entry. PI Dr. Michael Chislock and Project Manager Matthew Silverhart trained field technicians on proper water quality sample storage, processing, and analysis. Training took place June 17-21, 2024 at the SUNY Brockport campus and site 7052-Braddock Bay Wetland for field training. All field technicians were successfully trained and met pre-season and mid-season training performance criteria described in the project QAPP. These performance checks were administered by the associated PI and Project Manager.

Training will be conducted similarly for 2025.

Challenges and Lessons Learned (from fall report)

Transitioning to a new Project Manager, Matthew Silverhart, welcomed a large range of new challenges. Lake Ontario wetlands pose many challenges not experienced in sampling coastal wetlands in Lake Michigan and Lake Huron, which are what Matthew Silverhart was accustomed to sampling. At SUNY Brockport, the teams utilize “mud-buddy” motors, which operate differently than a traditional outboard motor, but allow for easier access and navigation of barrier wetlands that are rather commonly found in Lake Ontario.

Additionally, at SUNY Brockport, the Vegetation team travels and samples alongside the Fish/Invert/WQ team, leading to new logistical challenges not experienced before for the new Project Manager. Coordinating 2 crews across two boats and a canoe was a larger logistical challenge than previously encountered.

Site Visit List (from fall report)

Of the 20 assigned sites for the Fish/Invert/WQ team of SUNY Brockport, 11 were sampled in full, 3 were unable to be sampled (due to lack of vegetation that fit GLCWMP SOP criteria), and 6 could not be accessed (either due to physical barriers or lack of ability to sample safely. 12 of the assigned sites were panel sites, 4 were resample sites from the previous year, 2 were presample sites for the following season, and 3 were benchmark sites (site 28 was both a benchmark and a resample site). All benchmark sites were requested by SUNY Brockport PIs due to restoration projects either being planned, ongoing, or having previously occurred at the specified sites. This information can be used to better inform and shape restoration efforts.

Panel Survey Results (from fall report)

38 species of fish (and turtles) were observed during the fyke net sampling of panel sites with three of those species being listed as non-native by the New York Department of Environmental Conservation (NYDEC. The three non-native species caught during panel sampling were Common Carp, Freshwater Tubenose Goby, and Round Goby. Spotted Gar, which are listed as endangered by the NYDEC, were encountered while sampling site 1940, which is Presque Isle State Park in Pennsylvania. Panel sampling began on 26 June 2024 at site 28 and concluded on 19 August 2024 at site 82.



(Left to right) Dillon VanDemortel, Kai Schedel, and Grace Trebilcock performing sweeps with D-nets to collect macroinvertebrates at site 5196: Collin's Creek Wetland. Photo taken by Matthew Silverhart.

Extra Sites and Data (from fall report)

23 species of fish (and turtles) were observed during the fyke net sampling of benchmark sites with two of those species being listed as non-native by the NYDEC. The two non-native species caught during benchmark sampling were Common Carp and Round Goby. Benchmark sampling began on 20 June 2024 at site 7052 and concluded on 27 June 2024 at site 51. Each benchmark site was requested by PIs at SUNY Brockport to continue monitoring of previous restoration efforts conducted at those sites.

Wetland Condition Observations and Results (from fall report)

Many of the barrier wetlands sampled have been separated from the open water of the Great Lakes basin by roads and culverts. While this is not a new occurrence, it makes it increasingly difficult to access barrier wetlands and to evaluate their connectivity. This, coupled with changing water levels around the Great Lakes basin, can have impacts on the seasonal passage for fish both to and from the barrier wetlands.

A vast majority of the wetlands sampled exhibited large mats of floating *Typha spp.* which were nearly impenetrable for sampling for Fish/Invert/WQ. While they are a monodominant vegetation zone, they do not allow for any of the Fish/Invert/WQ teams to sample because there is no water on top of the mat and the water beneath them is inaccessible due to the thick root structures of the vegetation. Even if you can penetrate through the mat, there is only thick muck below.

Data Processing

SUNY Brockport crews have completed 100% data entry and QC checks for fish, field water quality, and field invertebrate data. 100% unknown fish were identified and entered in the database. 100% laboratory water quality analyses, data entry, and QC checks were completed. Laboratory invertebrate processing and identification have been completed, entered, and QC'd.

Mid-season QC Check Findings

Mid-season QC check for the Fish/Invert/WQ team occurred at site 7023 on 22 July 2024 with Matthew Silverhart and Dr. Michael Chislock administering the mid-season QC check. The crew members involved were Dillon VanDemortel, Kai Schedel, and Grace Trebilcock. The crew performed all tasks to satisfaction and there were no issues noted that needed to be addressed.

Mid-season checks will be performed in 2025.

Audit and QC Report and Findings

Other than minor data entry errors, there were no large-scale errors of note for the Fish/Invert/WQ data entry and QC. The QC report was completed for the Invert ID following the completion of that task and entry which took place in January of 2025

Additional Funding and Projects

No additional funding was used for any related projects or additional sampling.

Other Collaboration Activities (from fall report)

None.

Other Data Requests

There were no additional data requests during the 2024 sampling season for the Fish/Invert/WQ sampling team.

Related Student Research

No student research coincided with Fish/Invert/WQ sampling this season.

US EASTERN BASIN VEGETATION TEAM AT SUNY BROCKPORT

Team Members

- Dr. Kathryn Amatangelo, PI (since 2014)
- Dr. Rachel Schultz, PI (since 2019)
- Matthew Silverhart, project manager, field crew leader (since 2020)
- Kendalyn Town, graduate research assistant (since 2022)
- Sophia Maum, undergraduate technician (since 2024)

Training

Both field technicians (Kendalyn Town and Sophia Maum) were trained by PI Dr. Rachel Schultz and project manager Matthew Silverhart on proper field sampling techniques, data collection and recording, GPS use, and canoe use and safety. Both technicians were trained by PI Dr. Rachel Schultz in plant identification and sample preservation and storage. All training took place June 17-20, 2024 at the SUNY Brockport campus and benchmark site 7052-Braddock Bay, for field training. Lastly, both field technicians were trained in data entry and QC checks in the database by project manager Matthew Silverhart. Both field technicians were successfully trained, passed the plant identification quiz, and met pre-season training performance criteria described in the project QAPP.

Training will be conducted in a similar fashion for 2025.

Challenges and Lessons Learned (from fall report)

Transitioning to a new Project Manager, Matthew Silverhart, welcomed a large range of new challenges. Lake Ontario wetlands pose many challenges not experienced in sampling coastal wetlands in Lake Michigan and Lake Huron, which are what Matthew Silverhart was accustomed to sampling. At SUNY Brockport, the Vegetation team travels and samples alongside the Fish/Invert/WQ teams, leading to new logistical challenges not experienced before for the new Project Manager. Coordinating 2 crews across two boats and a canoe was a larger logistical challenge than previously encountered.

Site Visit List (from fall report)

Of the 20 assigned sites for the Vegetation team of SUNY Brockport, 17 were sampled in full and 3 could not be accessed (either due to physical barriers or lack of ability to sample safely. Twelve of the assigned sites were panel sites, 4 were resample sites from the previous year, 2 were presample sites for the following season, and 3 were benchmark sites (site 28 was both a benchmark and a resample site). All benchmark sites were requested by SUNY Brockport PIs due to restoration projects either being planned, ongoing, or having previously occurred at the specified sites. This information can be used to better inform and shape restoration efforts.

Panel Survey Results (from fall report)

During the sampling of panel sites, 31 species of non-native plants were identified and 19 of those species were classified as invasive. *Lobelia cardinalis*, which is listed as an “exploitably vulnerable native plant” by the New York Department of Environmental Conservation, was encountered while sampling site 82: Blind Sodus Bay. Panel sampling began on 26 June 2024 at site 28 and concluded on 19 August 2024 at site 82.

Extra Sites and Data (from fall report)

At the benchmark sites, there were 16 non-native species identified and 14 of those species were listed as invasive species. Benchmark sampling began on 20 June 2024 at site 7052 and concluded on 27 June 2024 at site 51. Each benchmark site was requested by PIs at SUNY Brockport to continue monitoring of previously restoration efforts conducted at those sites.

One additional quadrat at the start point of each transect, along the wetland-upland edge (aside from any transects where the narrow sampling procedure was used in the uppermost vegetation zone) was collected for use in a thesis project by Kendalyn Town. In each quadrat, all plant species were identified, and their percent cover was estimated. This data will be used to answer questions about whether wetland vegetation species are using the wetland-upland edges as refugia.

Wetland Condition Observations and Results (from fall report)

Many of the barrier wetlands encountered have been separated from the open water of the Great Lakes basin by roads and culverts. A vast majority of the wetlands sampled exhibited large mats of floating *Typha spp.* which made it difficult for vegetation crews to access the meadow portion of the transects.



Sophia Maum (front) and Kendalyn Town (back) record visual observations of vegetation in a quadrat at site 82: Blind Sodus Bay. Photo taken by Dr. Rachel Schultz.

Data Processing

SUNY Brockport crews have completed 100% data entry and QC checks for vegetation data. 100% unknown plants were identified and entered in the database.

Mid-season QC Check Findings

Mid-season QC check for the Vegetation team occurred at site 82 on 18 July 2024 with PI Dr. Rachel Schultz administering the mid-season QC check. The crew members involved were

Kendalyn Town and Sophia Maum. The crew performed all tasks to satisfaction and there were no issues noted that needed to be addressed.

Mid-season checks will be performed similarly in 2025.

Audit and QC Report and Findings

Other than minor data entry errors, there were no large-scale errors of note for the vegetation data entry and QC.

Additional Funding and Projects

The only additional project was the aforementioned additional data collection for Kendalyn Town's project. This work required no additional funding for supplies or equipment as all equipment needed for sampling was already present with the crew.

Other Collaboration Activities (from fall report)

Kendalyn Town had requested the additional quadrat data collection be conducted by the other crews that take part in the GLCWMP vegetation sampling. Other crews have been sending over their data to Kendalyn Town as part of this collaboration.

Other Data Requests

Catherine Landis at SUNY ESF has requested Wild Rice data from PI Dr. Rachel Schultz.

Related Student Research

Please see the aforementioned Kendalyn Town thesis project description for this portion.

ASSESSMENT AND OVERSIGHT

The Quality Assurance Project Plan (QAPP) for this program was originally written, signed by all co-PIs, and approved by USEPA in the spring of 2011, prior to beginning any fieldwork. Throughout the first round of the project (2011-2015), five revisions were made to the QAPP. These revisions were necessary to improve methodology, better clarify protocols, and ensure the safety of all personnel. After each revision, all co-PIs and US EPA reviewed and signed the updated document prior to commencing fieldwork. The final QAPP revision for round 1 of the project was signed in March 2015. This 2015 revision (QAPP_r5) served as the basis for the second round of monitoring (2016-2020).

For the second 5-year sampling rotation, no substantial methodological or quality assurance/quality control changes were necessary. The QAPP_r5 document was reviewed by project PIs prior to our February 19, 2016 project meeting. The only changes that were required to QAPP_r5 related to the data management system. Project PIs signed the updated QAPP (QAPP_CWMII_v1) at the February 19, 2016 meeting. In thoroughly reviewing the QAPP and SOPs in early 2018, crews found inconsistencies between the QAPP and SOPs and another handful of minor corrections and clarifications. PIs signed off on these changes at the 2018 PI meeting in Michigan in February. These fixes were incorporated into the QAPP in 2018 and PIs again signed off on the QAPP at the March 1, 2019, meeting in Michigan. The updated QAPP (QAPP_CWMII_rev 1) and SOPs were submitted to EPA in April of 2019.

For the third 5-year sampling rotation, again no substantial methodological or QA/QC changes were necessary. The QAPP was updated to reflect turnover in program personnel, to continue to strive for clarity and understandability by others and to make the QAPP more of a stand-alone document without reference to proposals or reports, and to remove inconsistencies between the QAPP and SOPs. The only substantive change was to update the water chemistry section to better reflect the updated EPA guidance on calculating error and variability in various water chemistry measurements. This QAPP (QAPP_CWMPIII_2021) was signed by PIs in the spring of 2021. The QAPP was updated in spring of 2023 (signed by all PIs) to reflect the re-creation of the Site Management System by Limnotech to be housed at Central Michigan University. We are in the process of again updating the water quality SOP and that section of the QAPP to further clarify a few things and ensure that crews have the guidance they need to avoid confusion. These changes will be finalized and the QAPP signed by PIs in late 2025.

Major QA/QC elements that are on-going for this program:

- Training of all new laboratory staff responsible for macroinvertebrate sample processing: This training is conducted by experienced technicians at each regional lab

and is overseen by the respective co-PI or resident macroinvertebrate expert. Those labs without such an expert sent their new staff to the closest collaborating lab for training. Macroinvertebrate IDers communicate with each other via their own email list and assist each other with difficult identifications and other questions that arise. Every few years, typically when a major identification guide is updated, IDers for all teams meet either in-person or virtually to discuss taxonomic issues and questions.

- Training of all fish, macroinvertebrate, vegetation, bird, anuran and water quality field crew members following the QAPP and SOPs. This included passing tests for procedural competence as well as identification tests for fish, vegetation, birds, and anurans. Training certification documents were archived with the lead PI and QA managers.
- GPS testing: Every GPS unit used during the field season was tested for accuracy and its ability to upload data to a computer. Field staff collected a series of points at locations that could be recognized on a Google Earth image (e.g., sidewalk intersections) then uploaded the points to Google Earth and viewed the points for accuracy. Precision was calculated by using the measurement tool in Google Earth. Results of these tests have been archived and referenced to each GPS receiver by serial number.
- Review of sites rejected after initial site visits: In cases where a site was rejected during a site visit, the reason for rejection was documented by the field crew in the site selection database. The project QA managers (Brady and Cooper) then reviewed these records to ensure consistency among crews. Occasionally, field crew leaders contacted Uzarski, Brady, or Cooper when deciding whether to reject a site. The frequency of these consultations increased in 2018 and 2019 as high water levels made sampling particularly challenging, but had returned to normal by 2020 as crews have become more accustomed to the high water levels and because water levels dropped quite a bit in 2021 and again in 2023 due to drought across the upper Great Lakes. Water levels are expected to be around “normal” for 2025.
- Collection of all training/certification documents and mid-season QA/QC forms from regional labs: These documents will be retained as a permanent record for the project.
- Maintenance, calibration, and documentation for all field meters: All field meters were calibrated and maintained according to manufacturer recommendations. Calibration/maintenance records are being archived at each institution.

- Collection of duplicate field samples: Precision and accuracy of many field-collected variables is being evaluated with duplicate samples. Duplicate water quality samples were collected in conjunction with approximately every 10th WQ sample collected.
- QC checks for all data entered into the data management system (DMS): Every data point that is entered into the DMS is being checked to verify consistency between the primary record (e.g., field data sheet) and the database. QC should be complete for all data by the spring semi-annual report submission each year.
- Linking of GPS points with field database: Inevitably, some errors occur when crew members type in GPS waypoint names and numbers. All non-linking points between these two databases were assessed and corrected in 2014, which took a hundred or more person-hours. We now have a more automated way to link GPS waypoints with data, crews are paying more attention to waypoint name/number accuracy, and the lat/longs for critical locations are being typed directly into the data management system. These three actions have greatly reduced number of GPS waypoints that cannot be linked to data in the DMS system.
- Mid-season QC checks: These were completed by PIs or head field crew leaders for each of the field crews to ensure that there were no sampling issues that developed after training and while crews were sampling on their own.
- Creation/maintenance of specimen reference collections: Reference collections for macroinvertebrates, fish, and plants have either been created or are being maintained and updated by each regional team. Macroinvertebrate reference collections, in particular, were developed or expanded as these samples were processed. Vegetation reference collections are often being kept in collaboration with local herbaria.
- Data Quality Objectives (DQO) for laboratory analyses: Participating water quality laboratories have generated estimates of precision, bias, accuracy, representativeness, completeness, comparability, and sensitivity for all water quality analyses.

DATA VERIFICATION

In 2022-2023 we, in collaboration with GDIT, implemented a data verification protocol that is being used to identify and resolve, or otherwise flag, issues related to data accuracy, consistency, and compliance with the Quality Assurance Project Plan (QAPP) and SOPs established for sampling the various taxa groups. The overall goal of this process is to establish

the usability of each data record to ensure that the CWMP datasets are properly communicated to and applied by end data users. Initially, approximately 120 data verification criteria (rules) were developed by GDIT (USEPA's contractor) to conduct a suite of checks for specific components of the anuran, bird, vegetation, fish, macroinvertebrate, and water quality datasets. Examples of data verification checks include:

- Identifying bird surveys that took place outside the sampling seasonal frame (e.g., after breeding season).
- Identifying fish surveys for which nets did not fish correctly and yet the crew entered data from those nets.
- Identifying vegetation surveys for which some other number of transects than three was sampled.

The data verification checks have been automated by GDIT to run against the semi-annual CWMP database release (MS Access format) that is delivered to GLNPO in May and October of each year. Each record that fails to meet specific verification criteria (such as they listed above) is flagged with an appropriate data qualifier code (e.g., "LINTC" – lack of internal consistency, or "MRV" – missing required value). The results from the automated checks are written to a set of comma-separated variable (CSV) files (i.e., one file per check type), which are delivered by GDIT to LimnoTech for integration into the CWMP DMS. Over the past six months, LimnoTech has incorporated additional tables ("data_rev_*") into the DMS and developed a utility application to ingest the CSV files into those dedicated tables. The enhanced DMS provides the capability to store and manage multiple sets of data verification results, including tracking of issue resolution and the assignment of data usability flags on a record-specific basis. Verification check results are stored in a set of dedicated tables, which are readily linked to any CWMP taxa data table that the results may be associated with. While this approach supports linking the raw data to verification results/flags when needed, it also avoids burdening the raw data tables with the detailed verification information.

Due to the large variety and number of verification checks and results, a dedicated "Data Verification" page was implemented by LimnoTech on the CWMP main website to provide a platform for CWMP team members to efficient review and respond to individual verification results (Figure 38). The tool will allow any "Level 4" CWMP user to efficiently filter for verification results that are pertinent to their specific taxa team, to download the results to an Excel spreadsheet, and then to provide appropriate feedback for each individual result, including documenting the resolution of the issue (if any). Ultimately, each record will be assigned an appropriate data usability flag based on assessment by lead PIs.




Figure 38. CWMP data verification user interface.

The CWMP lead PIs reviewed the verification criteria information provided by GLNPO and GDIT, as well as the data verification tool described above. The tool was introduced and training provided to all taxonomic groups in the fall of 2022.

This effort has initially been focused on addressing a set of DV check results generated and provided by GDIT (EPA contractor) in fall 2022 for the 2016-2021 monitoring datasets. In addition to achieving improved data quality, consistency and documentation, this effort has provided opportunities to “tune” the rules for some DV checks and to plan and implement improvements to QA/QC methods used during data entry and review of annual monitoring datasets prior to the semi-annual database releases to EPA. The DV work is being facilitated by the Data Verification Interface (DVI) a custom tool accessible to CWMP team members through the program website. The DVI provides taxa teams with a streamlined approach for reviewing DV issues, applying corrections to data records (where applicable), documenting the check’s resolution status, and assigning data usability status. To complement the information that taxa teams provide on DV issue cause, resolution, and data usability, the DVI is being enhanced to provide a “post-audit” analysis of the status of individual records. This capability, which is currently being incorporated into the DVI, will allow teams and the lead PIs to identify and address any outstanding data quality gaps following the initial review effort.

As of May 2024, substantial progress had been made in addressing the 2016-2021 DV check results. More than 14,500 issues were originally identified by the DV checks in the 2016-2021 dataset, and greater than 95% of those issues have been reviewed and addressed in some fashion by the taxa teams. In addition, the taxa teams have reviewed and addressed roughly 95% of the 2022 DV check results provided by GDIT last fall. Work on addressing the remaining issues for 2016-2021 and 2022 will be ongoing this year, with a goal of completing data verification work on those datasets this calendar year (2025). Teams are now also beginning work on addressing check results provided by GDIT for the 2011-2015 datasets. It is anticipated that review of those datasets will be ongoing into calendar year 2026.

EXAMPLE WATER QUALITY QC INFORMATION

Laboratory Quality Assurances:

Water quality analyses from 2024 have been completed by the NRRI Central Analytical Laboratory, Central Michigan University's Wetland Ecology Laboratory, Grand Valley State University's Annis Water Resources Institute, Brockport's water quality lab, and Environment Canada's national water quality lab. Laboratory results from 2024 have passed the criteria shown below (Table 20) or were excluded from the database.

Table 20. Data acceptance criteria for water quality analyses.

QA Component	Acceptance Criteria
External Standards (QCCS)	$\pm 10\%$
Standard curve	$r^2 \geq 0.99$
Blanks	$\pm 10\%$
Blank spikes	$\pm 20\%$
Mid-point check standards	$\pm 10\%$
Lab Duplicates	$\pm 15\%$ RPD* for samples above the LOQ**
Matrix spikes	$\pm 20\%$

**Relative Percent Difference (RPD)*: While our standard laboratory convention is to analyze 10% of the samples in duplicate and use %RSD ($100 * CV$) of the duplicates as a guide for accepting or rejecting the data, another measure of the variation of duplicates is RPD: $RPD = ((|x_1 - x_2|) / \text{mean}) * 100$.

** *LOQ = Limit of Quantification*: The LOQ is defined as the value for an analyte great enough to produce <15% RSD for its replication. $LOQ = 10(S.D.)$ where $10(S.D.)$ is 10 times the standard deviation of the gross blank signal and the standard deviation is measured for a set of two replicates (in most cases).

Variability in Field Replicates

An analysis of field duplicate variability for samples collected in 2024 is shown in Table 21. It is important to note that for many constituents, the variability within sample sets is related to the mean concentration, and as concentrations approach the method detection limit (MDL), the variability increases dramatically. A calculation of field replicate variability with values at or near the level of detection will often result in high RPDs. For example, if the chlorophyll measurements on a set of field duplicates are 0.8 µg/L and 0.3 µg/L, mean = 0.6, resulting in a RPD of 91% ($RPD = [abs(rep\ a - rep\ b) / (rep\ a + rep\ b) / 2] * 100$), but since the MDL is $\pm 0.5\ \mu\text{g/L}$, this can be misleading.

The same can occur with analyte lab duplicates, and in these instances the QA officer will determine whether data are acceptable. It is also important to note that RPD on field duplicates incorporates environmental (e.g., spatial) variability, since duplicate samples are collected from adjacent locations, as well as analytical variability (e.g., instrument drift). Therefore, RPD of field duplicates is generally higher than RPD of laboratory duplicates. Table 21 below lists average RPD values for 2024. Higher than expected average RPD values were associated with a preponderance of near detection limit values for ammonium, nitrate, and soluble reactive phosphorus (SRP), and high spatial variability for chlorophyll and turbidity. Other variables, such as Total N, had values that were well above detection limits and low spatial variability; therefore, these values had much lower average RPD. Acceptance of data associated with higher-than-expected RPD was determined by the QA officers. The maximum expected RPD values are based on the MN Pollution Control Agency quality assurance project plan provided for the Event Based Sampling Program (<http://www.pca.state.mn.us/index.php/water/water-types-and-programs/surface-water/surface-water-financial-assistance/event-based-sampling-grants.html#for-grantees>).

Table 21. Field duplicate sample variability for 2024 in relative percent difference for water quality parameters with the acceptance criteria. The maximum expected RPD values are based on the MN Pollution Control Agency quality monitoring requirements for integrated assessments (<https://www.pca.state.mn.us/sites/default/files/wq-s1-15n.pdf>). Average RPD (n) min-max RPD.

Analyte	MDL	Maximum expected RPD	2024
Chlorophyll-a ug/L	0.5 µg/l All Labs 0.01 µg/L Brockport	30	30 (18) 0-131
Total Phosphorus mg/L	0.0008 mg/L Brockport 0.006 mg/L CMU 0.0005 mg/L Env Can 0.006 mg/L GVSU 0.004 mg/L NRRI 0.0005 mg/L U Windsor	30	17 (18) 2-61
*Soluble Reactive Phosphorus mg/L	0.0003 mg/L Brockport 0.005 mg/L CMU 0.0002 mg/L Env Can 0.005 mg/L GVSU 0.003 mg/L NRRI 0.0002 mg/L U Windsor	10	38 (18) 0-181
Total Nitrogen mg/L	0.014 mg/L Brockport 0.027 mg/L CMU 0.015 mg/L Env Can 0.1 mg/L GVSU 0.02 mg/L NRRI 0.015 mg/L U Windsor	30	6 (18) 0.2-27
*NH4-N mg/L	0.002 mg/L Brockport 0.01 mg/L CMU 0.005 mg/L Env Can 0.01 mg/L GVSU 0.008 mg/L NRRI 0.005 mg/L U Windsor	10	28 (18) 0-142
*NO2/NO3-N mg/L	0.002 mg/L Brockport 0.008 mg/L CMU 0.005 mg/L Env Can 0.005 mg/L NRRI 0.005 mg/L U Windsor	10	12 (18) 0-44
True Color pt-co	1 CU Brockport 0.5 CU Env Can 2 CU NRRI 1 CU U Windsor	10	14 (13) 0-63
Chloride mg/L	0.1 mg/L CMU 0.01 mg/L Env Can	20	19 (15) 0-159

Table 21. Field duplicate sample variability for 2024 in relative percent difference for water quality parameters with the acceptance criteria. The maximum expected RPD values are based on the MN Pollution Control Agency quality monitoring requirements for integrated assessments (<https://www.pca.state.mn.us/sites/default/files/wq-s1-15n.pdf>). Average RPD (n) min-max RPD.

Analyte	MDL	Maximum expected RPD	2024
	1.2 mg/L NRRI 0.01 mg/L U Windsor		

Notes:

*The variability between soluble reactive phosphorus, ammonium-N and nitrate/nitrite-N field replicates often exceeded the criteria, however many values for each were < 10 X the MDL

Field duplicates are a second sample taken immediately after an initial sample in the exact same location to assess the site, sampling and possible temporal variability. Duplicate samples are collected in the exactly the same manner as the first sample, including the normal sampling equipment cleaning procedures. The relative percent difference (RPD) between the duplicate samples is calculated with the following equation:

$$RPD = (| \text{Result 1} - \text{Result 2} |) / ((\text{Result 1} + \text{Result 2}) / 2) \times 100$$

COMMUNICATION AMONG PERSONNEL

Regional team leaders and co-PIs continue to maintain close communication as the program enters its fifteenth year (fifth and final year of round 3 sampling). Nearly all program members virtually attended an all-hands Zoom program organizational meeting in February of 2024. The 2025 PI meeting was cancelled due to circumstances beyond our control. Holding these meetings virtually means that PIs can have field and laboratory technicians and grad students attend without worrying about having a travel budget since budgets are very strained now. At these meetings, PIs discuss issues pertaining to the upcoming field season, how we can keep field teams safe, data validation and correction, manuscripts, and report products. Individual taxonomic teams held their meetings virtually just before or after the overall program meeting.

Regional team leaders and co-PIs continue to hold many conference calls and e-mail discussions regarding fieldwork, taxonomic changes, data analysis, indicator refinement, and publications throughout the duration of the project. Typically, most PIs spend the first week of field season in the field with their crews to ensure that all protocols are being followed according to the standards set forth in the QAPP and SOPs and to certify or re-certify crew members. That changed because of Covid-19 (depending on the field crew and PI), but in 2023 the field season returned to normal and we have high hopes that such normality will continue. Again this year

most crews have returning and experienced personal, and the PIs will be in contact and do training and provide advice in the manner that best suits their circumstances, at a minimum via phone calls and webinars. Under all circumstances, PIs keep in close contact with crews via cell phone, text, and email, and the leadership team is also always available via cell phone and text to answer crew questions.

OVERALL

The quality management system developed for this project has been fully implemented and PIs and their respective staff members continue to follow established protocols very closely, relying on the QAPP and SOPs as guiding documents. QA managers were also encouraged by each crew's continued willingness to contact their supervisors or, in many cases, the project management team when questions arise.

Despite the somewhat dangerous nature of this work, injury rates continue to be very low. We are very proud of what our field crews accomplished safely despite a global pandemic. Crews sampled safely, accurately, and without spreading Covid-19. The entire CWM team is relieved that crews continue to maintain an exemplary safety record. This is due to the leadership and safety consciousness of PIs, field crew chiefs, and field team leaders. PIs are not complacent about the lack of injuries and are grateful for the willingness of their crews to work long hours day after day, to successfully sample under often adverse conditions (including a global pandemic), and to conduct that sampling in accordance with strict QA procedures.

LEVERAGED BENEFITS OF PROJECT (2010 – 2023)

This project has generated a number of spin-off projects and serves as a platform for many graduate and undergraduate thesis topics. In addition, project PIs are collaborating with many other groups to assist them in getting data for areas that are or will be restored or that are under consideration for protection. Finally, the project supports or partially supports many jobs (jobs created/retained). All of these are detailed below.

SPIN-OFF PROJECTS (CUMULATIVE SINCE 2010)

Investigating the Use of eDNA to Determine Fish Use of Otherwise Unsampleable Habitats:

Some habitats cannot be sampled using fyke nets because of inappropriate water depth, unstable or unconsolidated bottom sediments or because that habitat is too fragile (e.g. wild rice). CoPI Valerie Brady with NRRI researcher Chan Lan Chun are investigating how well fyke net fish catches agree with fish eDNA collected from nearby benthic sediment to determine if eDNA could be used as a surrogate in situations where fish cannot be physically collected to determine habitat use.

Macroinvertebrate Monitoring for Delisting the Degradation of Benthos Beneficial Use

Impairment in the Muskegon Lake Area of Concern: The West Michigan Shoreline Regional Development Commission, with support from the Michigan Department of Environment, Great Lakes, and Energy funded a project to conduct macroinvertebrate sampling at 2 coastal wetlands in the Muskegon Lake Area of Concern in an effort to evaluate “Degradation of Benthos” BUI in the AOC. Samples were collected in 2021 and 2023 and data from several Lake Michigan reference wetlands were used to compare the AOC restoration sites. Dr. Matt Cooper led this project with students from Muskegon Community College.

Compiling and Assessing IBI and Environmental Stress Data to Assess Habitat Condition in the Detroit River Area of Concern (AOC):

The Detroit River Canadian Clean-up (convened by Environment and Climate Change Canada and the Province of Ontario) is evaluating the weight of evidence with regard to delisting several Beneficial Use Impairments in the Detroit River AOC (Degradation of Fish and Wildlife, Degradation of Benthos, and Loss of Fish and Wildlife Habitat). However, years of monitoring and assessment have failed to demonstrate clear time trends in the condition of biota (aquatic vegetation, aquatic macroinvertebrates, fishes, birds) of the Detroit River’s aquatic and riparian habitats. Attempts to evaluate indices of biotic integrity (IBIs) using the Reference Condition Approach (RCA) have been limited by an inability to achieve consensus on appropriate reference conditions. CoPIs Jan Ciborowski, Greg Grabas and Doug Tozer compiled land-based stressor data at the scale of second-order watersheds for the Detroit River AOC to let us assess how the IBI scores for sites in the Detroit River and adjacent areas (Lake Erie, Lake St. Clair, St. Clair River) vary as a function of environmental stress. We compiled all available biological monitoring datasets relating to aquatic vegetation, macroinvertebrates, fishes and birds within the study region and calculated composite measures of condition (IBIs) for each of the groups of biota and plotted the resulting scores against the stressor measures. We found provisional evidence of environmental stress thresholds for at least one IBI of each of the taxa investigated. Mapping the distribution of

nondegraded vs. degraded watersheds for each of the biological groups will help the DRCC identify whether and where further remediation is necessary to allow delisting of the BUIs.

Minnesota Land Trust Natural Areas Project and Grassy Point Restoration: In 2018, the Minnesota Land Trust contracted a project with the Natural Resources Research Institute in Duluth, MN to conduct bird surveys along the St. Louis River Estuary (SLRE), within nine project areas that were nominated for inclusion in the Duluth Natural Areas Program (DNAP). This program was created in 2002 to manage Duluth's environmentally significant areas to ensure the preservation of services and values such as habitat diversity and water quality. In addition to data collected for this project, we also included breeding bird data collected by the CWMP at benchmark sites located within the SLRE that aligned spatially with the nine DNAP project areas. Collectively these data were used to determine if the proposed land parcels included in the nomination met the criteria of qualifying as an Important Bird Congregation Area (criteria included numeric thresholds for different guilds of species). Use of these data qualified all nine parcels as meeting the Important Bird Congregation Area criteria.

These data were then used in a spin-off project with Minnesota Land Trust, where bird communities were associated with spatially-explicit environmental and habitat variables to help guide conservation and management effort in the SLRE. In this project we were also able to identify habitat availability at the landscape-level to identify specific features that are under-represented in the SLRE but likely important to avian species (specifically wetland-dependent species). These analyses have been used to guide restoration plans at specific locations within the SLRE, including Grassy Point (a wetland located in a heavily industrialized area of the SLRE). Efforts to restore this wetland site are being developed by using the habitat requirements of wetland-dependent marsh bird species as a guide and restoration goal. The plans for Grassy Point are complete and on-the-ground restoration is scheduled to begin in the spring of 2020. NRRI CWMP teams will be involved in post-restoration monitoring of this site as well.

Deriving and Calibrating Environmental and Biological data for Lake Erie in Support of the Great Lakes Water Quality Agreement's Nearshore Framework: As part of the Annex 2 and Annex 7 plans of the revised GLWQA, Environment and Climate Change Canada (ECCC) and GLNPO began work to jointly develop an Integrated Nearshore Framework for the Great Lakes. The goal was to assemble scientific and technical recommendations for nearshore assessment. The assessment was expected to be used to set priorities and design an approach to identify areas of high quality for protection and areas under stress requiring restoration. ECCC and GLNPO convened several workshops beginning in 2014. In 2016, ECCC initiated a pilot project on the Canadian side of Lake Erie to come up with a workable methodology and approach to combining assessments of different condition measures. CWM coPIs Jan Ciborowski and Greg Grabas took part in a series

of workshops and contributed information collected in part from CWM wetland surveys on Lake Erie. The first overall assessment of the nearshore in Lake Erie was reported in 2018. The weight of evidence indicated that there is a strong east to west gradient in nearshore condition with the highest quality habitat and biota observed in the eastern basin, and low quality in the western basin, influenced largely by seasonal occurrences of cyanobacteria. The nearshore of the Detroit River and Lake St. Clair was classified as being of moderate quality. Insufficient data were available to assess the St. Clair River. Assessments of the condition of coastal wetland across the study area were limited by variation in the types of data collected by different programs. A future goal will be to determine how best to align data collected from other programs with information collected using the CWM protocols.

Real-Time Logging of Water Level, DO, Light, and Wind to Assess Hydrological Conditions in

Great Lakes Coastal Wetlands: The University of Windsor is coordinating a project to test the hypothesis that the numbers and species of fishes caught in wetland fyke nets are related to temporal variation in dissolved-oxygen (DO), and that such DO variation is partly driven by seiche activity causing temporary movement of cool, well-oxygenated lakewater into and out of wetlands. This variation in DO may be especially important in the densely vegetated, shoreline-associated wetland zones (usually wet meadow, under high-water conditions). An SOP document was developed in spring 2019 and circulated to all field crews.

Each field team has been encouraged to deploy water level and DO loggers at their fyke net sites over the course of the summer. In addition to providing important basic hydrological information about the condition of coastal wetlands, the resulting Great Lakes-wide dataset will be used to help account for variation in fish catches and ultimately improve the precision of fish IBI estimates. Preliminary data collected over the field season and suggestions for improvement will be discussed at the winter field meeting.

Bathymetry and mapping of wetlands in Point Pelee National Park during a period of

hydrologic change: In 2018 Point Pelee National Park (PPNP) received approval through the Parks Canada Conservation and Restoration Project to begin a 4-year marsh restoration project. The project was focused 1) on increasing open water habitat and interspersions within the marsh and 2) reducing invasive vegetation. Members of the Ciborowski CWM team were asked if they would be able to conduct a preliminary survey of PPNP wetlands to determine the bottom profile and distribution of submerged aquatic vegetation. There was especial interest in the bathymetry of Lake Pond, whose eastern shoreline had been breached by wave action from Lake Erie during the summer as a consequence of the historically high water levels. In fall 2018 and during the 2019 field season, we conducted a benchmark survey of vegetation, aquatic invertebrates and water chemistry. We also assessed water depth, macrophyte distribution and

cover and sediment characteristics throughout the wetland using the remotely-operated ROVER, which was developed for shallow-water data collection in remote locations. Water level and dissolved oxygen loggers set in place in the spring provided a full-season record of the frequency of seiches and associated changes in water quality. CWM researchers are anticipated to be involved as collaborators throughout the restoration project.

Inventory and distribution of zooplankton in coastal wetlands: As part of ongoing interest in assessing the condition of CWM wetlands we began assessing the community composition of zooplankton in the wetlands visited as part of the annual program. Pilot samples were first collected in 2017. In 2018, zooplankton samples were collected at 16 Great Lakes coastal wetlands, situated off Manitoulin Island, northern Lake Huron, the western basin of Lake Erie, the Bruce Peninsula and Georgian Bay. In each wetland, samples were collected at 3 shallow-water points along a dissolved oxygen gradient. Records of water depth, substrate characteristics and vegetation density and composition were also tabulated. The sampling methods were based on techniques proposed by Loughheed and Chow-Fraser (2002) in developing their Zooplankton Quality Index. Seven Lake Huron wetlands were sampled in 2019.

Evaluating Fish and Invertebrate Distribution in Great Lakes Coastal Wetlands - an Occupancy

Modeling Approach: Led by University of Windsor postdoctoral fellow student Martin Jeanmougin, this project involves fish PIs Joseph Gathman, Carl Ruetz, Dennis Higgs and Jan Ciborowski. Occupancy modelling is a statistical approach that allows one to estimate the probability that a taxon is present in an area and the probability that it can be detected by sampling. Applying this approach to the invertebrate and fish CWM data could help us to identify important environmental factors influencing the likelihood that selected taxa occur in particular habitats and to more accurately estimate their distribution across the Great Lakes. Also, an analysis of the detection patterns can provide important information on potential biases in the protocols we use to sample the biota. The previous work done by K. Dykstra of Grand Valley State University (Carl Ruetz's lab) for the thesis on Yellow Perch distribution will be a good starting point for this project.

Genetic Barcodes for Wetland Macroinvertebrates: Surveillance of aquatic macroinvertebrates in the Great Lakes is of utmost importance. However, many organisms, particularly aquatic macroinvertebrates, lack information that can assist in their identification, whether through molecular barcodes or morphological characteristics. We are using previously collected aquatic macroinvertebrate samples from throughout the Great Lakes basins to generate genetic barcodes that will assist in identification of species (MOTUs) and expand the currently available molecular genetic databases. Our work is targeting specific groups to improve morphological

identification to lowest taxonomic levels. Finally, we will be able to use these data to test the usefulness of metabarcoding for Great Lakes surveillance to provide managers with valuable monitoring information.

Assessing Climate Vulnerability in Apostle Islands Coastal Wetlands: Funded by the National Park Service and GLRI, a team from Northland College sampled fish, macroinvertebrates, vegetation, and hydrologic variables in lagoon wetlands throughout the Apostle Islands National Lakeshore to identify species and communities that may be particularly vulnerable to climate change. This work represents an intensification of sampling effort within a sensitive and relatively pristine area of the Great Lakes. Data from this project were analyzed in relation to CWMP data to put Apostle Islands wetlands into a broader Great Lakes context.

Functional Indicators of Coastal Wetland Condition: Funded by the USGS through a Cooperative Ecosystem Studies Unit (CESU), this pilot project ran from fall 2016 through fall of 2019 to better determine functional indicators of Great Lakes coastal wetland usage by Great Lakes fish species. Sampling was done during the spring and fall at about 15 US wetlands already being assessed for CWM indicators during the summer. Data collected focus on fish usage of wetlands and the forage base for those fish, evaluated using macroinvertebrate sampling and examination of fish gut contents. Special emphasis was placed on determining usage of wetlands by young or spawning fish.

Conservation Assessment for Amphibians and Birds of the Great Lakes: Several members of the CWM project team have initiated an effort to examine the role that Great Lakes wetlands play in the conservation of amphibians and birds in North America. The Great Lakes have many large, intact freshwater wetlands in the interior portion of the North American continent. Their unique character, size, and plant composition supports populations of many species of amphibians and birds, many of which have been identified as endangered, threatened, or of special concern in North America. CWM PIs will use the extensive data that have been gathered by USEPA, such as the Great Lakes Environmental Indicators project and the Great Lakes Wetlands Consortium, as well as Bird Studies Canada, as critical input to this assessment. The initial stages in the development of the conservation assessment will be to analyze habitat and landscape characteristics associated with Great Lakes coastal wetlands that are important to wetland-obligate bird species occupying these habitats. By combining breeding bird data from the sources above and incorporating landscape variables, classification trees can be developed to predict presence and relative abundance of these species across the Great Lakes Basin. These methods, outlined in Hannah Panci's thesis; 'Habitat and landscape characteristics that influence Sedge Wren (*Cisthorus platensis*) and Marsh Wren (*C. palustris*) distribution and abundance in Great Lakes Coastal Wetlands' (University of Minnesota Duluth). She compiled data for over 800 wetlands in her analysis, which will provide a basis for analyzing additional wetland-obligate species.

Bird and Anuran Metrics and Indicator Calculations: Avian and anuran responses to landscape stressors can be used to inform land managers about the health of coastal wetlands and the landscape stressors that affect these systems (Howe et. al. 2007). Data that has been entered into the data management system and QC'd are being used to calculate some of the metrics and indicators for these wetlands.

Influence of broadcast timing and survey duration on marsh breeding bird point count results: Several members of the project team, with D. Tozer as lead, examined the importance of survey duration and timing of broadcast playbacks on occurrence and counts of wetland breeding birds. The results of this analysis suggest that 10-min point counts are superior to 15-min counts which have important implications for future monitoring and cost-effectiveness. These findings have been published in the journal of Avian Conservation and Ecology (Tozer et al. 2017).

North Maumee Bay Survey of Diked Wetland vs. Un-Diked Wetland: Erie Marsh Preserve is being studied as a benchmark site for the CWM project. As a benchmark site, Erie Marsh Preserve will serve as a comparison against randomly-selected project sites, and will be surveyed each year of the CWM project. Benchmark sampling began prior to Phase 1 of a planned restoration by The Nature Conservancy, allowing for pre- and post-restoration comparisons. In addition, biota and habitat within the diked wetlands area will be compared to conditions outside of the dike, but still within the preserve. These data will also be used for post-construction comparisons to determine what biotic and abiotic changes will occur once restoration efforts have reconnected the dike to the shallow waters of Lake Erie.

Cattails-to-Methane Biofuels Research: CWM crews collected samples of invasive plants (hybrid cattail) which were analyzed by Kettering University and their Swedish Biogas partner to determine the amount of methane that can be generated from this invasive. These samples was compared to their data set of agricultural crops, sewage sludge, and livestock waste that are currently used to commercially generate methane. Results demonstrated that hybrid cattail and reed canary grass both generated adequate levels of methane for use as feedstocks for biodigestion. The result of this and other CWM data collection are summarized in the Carson *et al.* 2018 journal article. The cattails-to-methane biofuels project is also funded (separately) by GLRI.

Plant IBI Evaluation: A presentation at the 2014 Joint Aquatic Science meeting in Portland, Oregon evaluated Floristic Quality Index and Mean Conservatism score changes over time utilized data collected during the first three years of the GLRI study. Mean C scores showed little change between years from 2011 through 2013 due to stable water levels.

Correlation between Wetland Macrophytes and Wetland Soil Nutrients: CWM vegetation crews collected wetland soil samples and provided corresponding macrophyte data to

substantially increase the number of sites and samples available to the USEPA Mid-Continent Ecology Division. USEPA MED researchers studied wetland macrophyte and wetland soil nutrient correlations. The MED laboratory ran the sediment nutrient analyses and shared the data with CWM PIs.

Comparative study of bulrush growth between Great Lakes coastal wetlands and Pacific Northwest estuaries. This study includes investigation of water level effects on bulrush growth rates in Great Lakes coastal wetlands. With leveraged funding from NSF for the primary project on bulrush ability to withstand wave energy.

Braddock Bay, Lake Ontario, Sedge Meadow and Barrier Beach Restoration: Braddock Bay is being studied as a benchmark site in conjunction with the US Army Corps of Engineers to assess the current extent of, and potential restoration of, sedge meadow and the potential of restoring the eroded barrier beach to reduce wetland loss. CWM crews collected pre-restoration data to help plan and implement restoration activities and will collect post-restoration data to help plan and implement restoration activities and assess results. The results will help build a model for future sedge meadow restoration in Lake Ontario to mitigate the harmful impacts of invasive cattails and provide habitat for fish and wildlife species. Additionally, this project will be expanded, in conjunction with Ducks Unlimited, to four nearby wetlands, pending funding from NOAA.

Thunder Bay AOC, Lake Superior, Wetland Restoration: Nine wetlands around Thunder Bay were sampled for macroinvertebrates, water quality, and aquatic vegetation by CWM crews in 2013 using methods closely related to CWM methods. These data will provide pre-restoration baseline data as part of the AOC delisting process. Wetlands sampled included both wetlands in need of restoration and wetlands being used as a regional reference. All of this sampling was in addition to normal CWM sampling, and was done with funding from Environment Canada.

Common Tern Geolocator Project: In early June 2013, the NRRI CWM bird team volunteered to assist the Wisconsin DNR in deploying geolocator units on Common Terns nesting on Interstate Island. In 2013, 15 birds between the ages of 4-9 yrs old were outfitted with geolocators. Body measurements and blood samples were also taken to determine the sex of each individual. In June of 2014, geolocators were removed from seven birds that returned to nest on the island. Of the seven retrieved geolocators, four were from female birds and three from males. The data collected during the year will be used to better understand the migratory routes of Common Terns nesting on Interstate Island. This is the first time that geolocators have been placed on Common Terns nesting in the Midwest, which is important because this species is listed as threatened in Minnesota and endangered in Wisconsin. Tracking Common Terns throughout their annual cycle will help identify locations that are important during the non-breeding portion of their life cycle. Data are currently being analyzed by researchers at the Natural Resources Research Institute in Duluth MN.

Using Monitoring Results to Improve Management of Michigan's State-Owned Coastal

Wetlands: One year project, 2016-2017, awarded to Central Michigan University by the Michigan Department of Environmental Quality. The project will focus on the prioritization of high-quality and important state-owned coastal wetlands that have been monitored as part of the Great Lakes CWM program, and development of site-specific management plans for these wetlands which address diverse management goals and objectives with a broad focus including biodiversity, ecological services, habitat for fish and wildlife, climate change adaptation, and rare species.

Developing a Decision Support System for Prioritizing Protection and Restoration of

Great Lakes Coastal Wetlands: While a number of large coastal wetland restoration projects have been initiated in the Great Lakes, there remains little regional or basin-scale prioritization of restoration efforts. Until recently we lacked the data necessary for making systematic prioritization decisions for wetland protection and restoration. However, now that basin-wide coastal wetland monitoring data is available, development of a robust prioritization tool is possible and we propose to develop a new Decision Support System (DSS) to prioritize protection and restoration investments. This project, funded by the Upper Midwest and Great Lakes Landscape Conservation Cooperative, the Michigan Office of the Great Lakes, and the US Army Corp. of Engineers, has developed a DSS for wetlands along the US shoreline of the Great Lakes.

Quantifying Coastal Wetland – Nearshore Linkages in Lake Michigan for Sustaining Sport Fishes:

With support from Sea Grant (Illinois-Indiana and Wisconsin programs), personnel from UND and CWM are comparing food webs from coastal wetlands and nearshore areas of Lake Michigan to determine the importance of coastal wetlands in sustaining the Lake Michigan food web. The project emphasis is on identifying sport fish-mediated linkages between wetland and nearshore habitats. Specifically, we are (1) constructing cross-habitat food webs using stable C and N isotope mixing models, (2) estimating coastal wetland habitat use by sport fishes using otolith microchemistry, and (3) building predictive models of both linkage types that account for the major drivers of fish-mediated linkages in multiple Lake Michigan wetland types, including some wetlands sampled by the coastal wetland monitoring project. Collaborators are the University of Wisconsin – Green Bay and Loyola University Chicago.

Clough Island (Duluth/Superior) Preservation and Restoration: The Wisconsin Department of Natural Resources requested (and funded) a special report on sites sampled using CWM protocols around Clough Island within the St. Louis River Area of Concern (AOC). Their interests were to see if CWM data indicated any differences in habitat or species composition/abundances among Clough Island and other St. Louis River sites, and also how Clough Island compared to other nearby Lake Superior coastal wetlands. The 46 page report was submitted to Cherie Hagan of the WDNR in May of 2014. Clough Island was recently acquired by the Nature Conservancy and they are using the data in the report for their development of conservation plans for the area.

Floodwood Pond and Buck Pond South, Lake Ontario, Wetland Pothole Restoration: Open water potholes were established in these two wetlands by The Nature Conservancy to replace openings that had filled with cattail following lake-level regulation. CWM crews collected pre- and post-restoration data as benchmark sites in both wetlands to allow TNC to assess changes.

Buck Pond West and Buttonwood Creek, Lake Ontario, Sedge Meadow Restoration: These two wetlands in the Rochester Embayment AOC are actively being restored by a consortium involving Ducks Unlimited, The College at Brockport, NYS Department of Environmental Conservation, and the Town of Greece. CWM crews collected pre-restoration data as a benchmark site to help plan and implement restoration activities. Post-restoration data collection is underway under CWM to help assess results and help build a model for future sedge meadow restoration in Lake Ontario to mitigate the harmful impacts of invasive cattails and provide habitat for fish and wildlife species.

Salmon/West Creek, Long Pond, and Buck Pond East, Lake Ontario, Emergent Marsh Restoration: These three wetlands in the Rochester Embayment AOC are being studied as benchmark sites by CWM crews to provide the U.S. Fish and Wildlife Service with pre-restoration data for projects currently in the design phase. Future CWM data collection has been requested to assist in post-restoration assessment.

Lower Green Bay and Fox River AOC: Results from the Coastal Wetland Monitoring (CWM) Project and the Great Lakes Environmental Indicators (GLEI) Project are playing a central role in a \$471,000 effort to establish fish and wildlife beneficial use impairment (BUI) removal targets for the Lower Green Bay and Fox River AOC (2015-2017) 1) Protocols for intensive sampling of bird, anurans, and emergent wetland plants in the project area have followed the exact methods used in the CWM project so that results will be directly comparable with sites elsewhere in the Great Lakes. 2) Data from GLEI on diatoms, plants, invertebrates, fish, birds, and anurans and from CWM on birds and anurans have been used to identify sensitive species that are known to occur in the AOC and have shown to be sensitive to environmental stressors elsewhere in the Great Lakes. These species have been compiled into a database of priority conservation targets. 3) Methods of quantifying environmental condition developed and refined in the GLEI and CWM projects are being used to assess current condition of the AOC (as well as specific sites within the AOC) and to set specific targets for the removal of two important BUIs (fish and wildlife populations and fish and wildlife habitats). 4. Application of the Index of Ecological Condition method (e.g., Howe et al. 2007) for measuring the condition of birds, anurans, and other fish and wildlife groups. Follow-up work was funded for 2018-2020 at \$87,000 to continue refining field monitoring methods and metrics of 40 fish and wildlife habitats and populations.

SOGL/SOLEC Indicators: CWM project PIs have developed a set of indicator metrics for the State of the Great Lakes/State of the Lakes Ecosystem Conference (SOLEC). These metrics fill a much-needed gap in quantifying responses of biotic communities to environmental stress throughout the Great Lakes. Sites for all coastal wetlands sampled by the GLEI, CWM, and Marsh Monitoring Program projects have been scored according to several complementary indices that provide information about local and regional condition of existing wetlands.

Roxana Marsh Restoration (Lake Michigan): The University of Notre Dame (UND) team, led by graduate student Katherine O'Reilly and undergraduate Amelia McReynolds under the direction of project co-PI Gary Lamberti, leveraged the GLCWM monitoring project to do an assessment of recently-restored Roxana Marsh along the south shore of Lake Michigan. Roxana Marsh is a 10-ha coastal wetland located along the Grand Calumet River in northwestern Indiana. An EPA-led cleanup of the west branch of the Grand Calumet River AOC including the marsh was completed in 2012 and involved removing approximately 235,000 cubic yards of contaminated sediment and the reestablishment of native plants. Ms. McReynolds obtained a summer 2015 fellowship from the College of Science at UND to study the biological recovery of Roxana Marsh, during which several protocols from the GLCWM project were employed. During summer 2015 sampling of Roxana Marsh, an unexpected inhabitant of the Roxana Marsh was discovered -- the invasive oriental weatherfish (*Misgurnus anguillicaudatus*). Oriental weatherfish are native to southeast Asia and believed to have been introduced to the U.S. via the aquarium trade. Although there have been previous observations of *M. anguillicaudatus* in the river dating back to 2002, it had not been previously recorded in Roxana Marsh, and little information is available on its biological impacts there or elsewhere. We are currently using stable carbon and nitrogen isotopes, along with diet analysis, to determine the role of *M. anguillicaudatus* in the wetland food web and its potential for competition with native fauna for food or habitat resources. This discovery received media attention from the Illinois-Indiana Sea Grant College Program.

Chlorophyll-*a* Modeling: The UND team, in collaboration with Northland College, CMU, and others, is investigating the drivers that influence water column chlorophyll-*a* in coastal wetlands. Our hypothesis is that chlorophyll-*a* will be related to nutrient status of wetlands and degree of development of adjoining land. Along with CWM water data, we are utilizing GIS land use and connectivity data. Specifically, we seek to answer the following questions: (1) What variables best predict chlorophyll-*a* in coastal wetlands across the entire Great Lakes basin? (2) How do these variables change across each basin (i.e., Lake Michigan, Lake Erie, Lake Ontario, Lake Superior, Lake Huron)? (3) Are there differences in predictor variables across sub-basins (e.g., Lake Erie North vs. Lake Erie South)? (4) Does wetland type (lacustrine, riverine, or barrier) change chlorophyll-*a* predictors? (5) How do other potential variables, such as vegetation zone type or year, change chlorophyll-*a* predictors?

Invasion Vulnerability Index: The UND team, in collaboration with other CWM teams, aims to create a usable tool that predicts which aquatic invasive species from a list of 10 Great Lakes

Aquatic Nuisance Species Information System (GLANSIS) watchlist species are of highest concern for prevention and early detection. We will combine Habitat Suitability Indexes (HSIs) made using wetland site-specific physio-chemical measurements and potential pathway data (distance to potential introduction pathways and distance to known established populations). Ultimately, we will produce an interactive, exploratory tool where a wetland can be selected, and a table will appear that shows the breakdown of invasion risk by species as invasion likelihood scores. If more information is desired about how the invasion likelihood score was calculated, an attribute table will display the numerical values for each criterion in the model. One of the main concerns with invasive species is how climate change will alter habitat suitability. To accommodate this concern, we will also include versions with future climate change scenarios using published IPCC environmental conditions. This information will be packaged together in an IVI for Great Lakes wetlands usable by scientists, managers, and the general public.

Green Bay Area Wetlands: Data from the benchmark site Suamico River Area Wetland was requested by and shared with personnel from the Wisconsin Department of Natural Resources and The Nature Conservancy, who are involved in the restoration activities to re-connect a diked area with Green Bay. In 2011 NRRI sampled outside the diked area following CWM methods, and in 2013 we sampled within the diked area as a special request. The data were summarized for fish, invertebrates, water quality, birds, and vegetation and shared with David Halfmann (WDNR) and Nicole Van Helden (TNC).

Hybridizing fish: In 2013 the NRRI field crew encountered gar around the Green Bay area of Lake Michigan which exhibited mixed morphological traits of shortnose and longnose species. At that time, John Lyons at the Wisconsin Department of Natural Resources was working on a project to confirm hybrid individuals in the Fox River watershed (which drains into Green Bay, WI). Josh Dumke at NRRI contributed photos of gar captured in Green Bay during Coastal Wetland Monitoring fish surveys to John Lyons, and those contributions were acknowledged in a recently-published article: (Lyons, J., and J.T. Sipiorski. 2020. Possible large-scale hybridization and introgression between Longnose Gar (*Lepisosteus osseus*) and Shortnose Gar (*Lepisosteus platostomus*) in the Fox River drainage, Wisconsin. *American Midland Naturalist*, 183:105-115). In 2014 and 2015 Coastal Wetland Monitoring fish teams collected gar fin clips across the entire Great Lakes basin for a much more comprehensive look at species distributions and hybridization, but sample processing and analysis of those stored samples is dependent upon securing additional funds.

Management alternatives for hybrid cattail (*Typha x glauca*) 2011- 2014: Differing harvest regimes for hybrid cattail were evaluated at Cheboygan, Cedarville, and Munuscong Bay in northern Michigan with USEPA GLRI funding. At all of these sites plant data was collected by CWM and used as baseline data that was compared to control sites. Analyses demonstrated that during low-water conditions, native plant diversity was increased by harvest of hybrid cattail.

Impacts of hybrid cattail management on European frogbit (*Hydrocharis morsus-ranae*); This study, funded by MI DNR in 2016-2017 for research by Loyola Chicago and Oregon State University studied the response of European frogbit to cattail management, using CWM plant data collected in Munuscong Bay as baseline data. CWM data collected from 2011 to 2015 provided documentation of the expanding range of frogbit into the western Great Lakes. The study found that open, flooded stands of hybrid cattail provided important habitat for European frogbit, but that management to remove cattail was not effective for frogbit control.

Nutrient limitation in Great Lakes coastal wetlands: GLCWMP water quality data indicate that reactive nitrogen concentration is often much lower in wetland habitats than the adjacent Great Lake nearshore. With funding from Illinois-Indiana Sea Grant and the Wisconsin DNR we have evaluated the role of nitrogen limitation on benthic algal growth in wetlands throughout Lakes Michigan, Huron, and Superior.

SUPPORT FOR UN-AFFILIATED PROJECTS (2011 – 2023)

CWM PIs and data managers continue to provide data and support to other research projects around the Great Lakes even though CWM PIs are not collaborators on these projects. Dr. Laura Bourgeau-Chavez at Michigan Tech University mapped the spatial extent of Great Lakes coastal wetlands using GIS and satellite information to help in tracking wetland gains and losses over time (Implementation of the Great Lakes Coastal Wetlands Consortium Mapping Protocol, funded by GLRI). We provided her with vegetation data and sampling locations each year to assist with this effort. Dr. Bourgeau-Chavez was also given funding to assess herbicide effectiveness against *Phragmites* in Green Bay and Saginaw Bay. CWM data are being used to find the best locations, provide baseline data, and provide pointers on site access (from field crew notes) in support of this project.

Reports on new locations of non-native and invasive species: Vegetation sampling crews and PIs have been pro-active over the years in reporting new locations of invasive vegetation. Fish and macroinvertebrate PIs and crews have also realized that they may be discovering new locations of invasive species, particularly invasive macroinvertebrates. To ensure that all new sightings get recorded, we are pulling all records of non-native fish and macroinvertebrates out of the database once per year and sending these records to the Nonindigenous Aquatic Species tracking website maintained by USGS (<http://nas2.er.usgs.gov/>). Wetland vegetation PIs contributed new SOLEC indicator guidelines and reports and continue to participate in the indicator review process.

Wetland Floristic Quality in the St. Louis River Estuary: With support from WI Sea Grant 2014-2017, vegetation PI N. Danz has integrated vegetation surveys from the CWM project with data from 14 other recent projects in the estuary. A new relational database was created that is

being used to assess spatial and temporal patterns in floristic quality and to develop materials to inform and monitor wetland restorations in this AOC.

Coordination and Partnership with National Audubon: Per the agreement to share CWMP bird data with the National Audubon Society, we have provided data and guidance on appropriate use of these data for their project “Prioritizing coastal wetlands for marsh bird conservation in the U.S. Great Lakes”. The resulting manuscript from this project is currently in review with the journal *‘Biological Conservation’* and per the agreement all CWMP bird and anuran co-investigators have had the opportunity to contribute to the manuscript and be included as co-authors. We expect to maintain communications regarding any potential future use of the CWMP data by National Audubon and will continue to provide guidance on appropriate uses in future projects and analyses.

Targeting Invasive Plant Species in Wisconsin Coastal Wetlands: In collaboration with WI Department of Natural Resources and Lake Superior Research Institute, vegetation PIs have summarized patterns of invasive plant occurrence in Wisconsin coastal wetlands. These summaries are being used to develop a more comprehensive invasive plant monitoring strategy throughout the Wisconsin basin.

REQUESTS FOR ASSISTANCE COLLECTING MONITORING DATA (2011 – 2023)

Project PIs provided monitoring data and interpretation of data for many wetlands where restoration activities were being proposed by applicants for “Sustain Our Great Lakes” funding. This program is administered by the National Fish and Wildlife Foundation (NFWF) and includes GLRI funding. Proposal writers made data/information requests via NFWF, who communicated the requests to us. Lead PI Don Uzarski, with assistance from co-PIs, then pulled relevant project data and provided interpretations of IBI scores and water quality data. This information was then communicated to NFWF, who communicated with the applicants. This information sharing reflects the value of having coastal wetland monitoring data to inform restoration and protection decisions. We anticipate similar information sharing in the coming years as additional restoration and protection opportunities arise.

In addition to the NFWF program, CWM PIs have received many requests to sample particular wetlands of interest to various agencies and groups. In some instances the wetlands are scheduled for restoration and it is hoped that our project can provide pre-restoration data, and perhaps also provide post-restoration data to show the beginnings of site condition improvement, depending on the timing. Such requests have come from the St. Louis River (Lake Superior), Maumee Bay (Lake Erie), and Rochester (Lake Ontario) Area of Concern delisting groups, the Great Lakes National Park Service, the Nature Conservancy (sites across lakes Michigan and Huron for both groups), as well as state natural resource departments. Several requests involve restorations specifically targeted to create habitat for biota that are being sampled by CWM. Examples include: a NOAA-led restoration of wetlands bordering the Little Rapids of the St. Marys River to restore critical spawning habitat for many native freshwater

fishes and provide important nursery and rearing habitat in backwater areas; TNC-led restoration of pike spawning habitats on Lake Ontario and in Green Bay; a US Army Corps of Engineers project in Green Bay to create protective barrier islands and restore many acres of aquatic and wetland vegetation; a USACE project to improve wetland fish and vegetation habitat in Braddock Bay, Lake Ontario; a New York state project to increase nesting habitat for state-endangered black tern; and projects in Wisconsin to restore degraded coastal wetlands on the Lake Superior shore. Many of these restoration activities are being funded through GLRI, so through collaboration we increase efficiency and effectiveness of restoration efforts across the Great Lakes basin.

At some sites, restoration is still in the planning stages and restoration committees are interested in the data CWM can provide to help them create a restoration plan. This is happening in the St. Louis River AOC, in Sodus Bay, Lake Ontario, for the Rochester NY AOC, wetlands along Wisconsin's Lake Superior shoreline, and for the St. Marys River restoration in 2015 by tribal biologists at Sault Ste Marie.

Other groups have requested help sampling sites that are believed to be in very good condition (at least for their geographic location), or are among the last examples of their kind, and are on lists to be protected. These requests have come from The Nature Conservancy for Green Bay sites (they are developing a regional conservation strategy and attempting to protect the best remaining sites); the St. Louis River AOC delisting committee to provide target data for restoration work (i.e., what should a restored site "look" like); and the Wisconsin DNR Natural Heritage Inventory has requested assistance in looking for rare, endangered, and threatened species and habitats in all of the coastal wetlands along Wisconsin's Lake Superior coastline. Southern Lake Michigan wetlands have mostly been lost, and only three remain that are truly coastal wetlands. CWM PIs are working with Illinois agencies and conservation groups to collaboratively and thoroughly sample one of these sites, and the results will be used to help manage all 3 sites.

Other managers have also requested data to help them better manage wetland areas. For example, the Michigan Clean Water Corps requested CWM data to better understand and manage Stony Lake, Michigan. Staff of a coal-fired power plant abutting a CWM site requested our fish data to help them better understand and manage the effects of their outfalls on the resident fish community. The Michigan Natural Features Inventory is requesting our data as part of a GLRI-funded invasive species mapping project. The US Fish and Wildlife Service requested all data possible from wetlands located within the Rochester, NY, Area of Concern as they assess trends in the wetlands and compare data to designated delisting criteria. The NERR on Lake Erie (Old Woman Creek) has requested our monitoring data to add to their own. The University of Wisconsin Green Bay will use our data to monitor control of *Phragmites* in one of their wetlands, and hope to show habitat restoration. Thunder Bay National Marine Sanctuary (Lake Huron) has requested our data to facilitate protection and management of coastal resources within the Sanctuary. The Wisconsin DNR has requested data for the Fish Creek

Wetland as part of an Environmental Impact Assessment related to a proposed Confined Animal Feeding Operation upstream of the wetland.

We have received a request from the USFWS for data to support development of a black tern distribution/habitat model for the Great Lakes region. The initial effort will focus on Lakes Huron, Erie and their connecting channels. Various FWS programs (e.g., Migratory Bird, Joint Venture, and Landscape Conservation Cooperatives) are interested in this model as an input to conservation planning for Great Lakes wetlands.

The College at Brockport has been notifying an invasive species rapid-response team led by The Nature Conservancy after each new sighting of water chestnut. Coupling the monitoring efforts of this project with a rapid-response team helped to eradicate small infestations of this new invasive before it became a more established infestation.

We are also now receiving requests to do methods comparison studies. For example, USGS and Five Fathom National Marine Park have both requested data and sampling to compare with their own sampling data.

Overall, CWM PIs have had many requests to sample specific wetlands. It has been challenging to accommodate all requests within our statistical sampling design and our sampling capacities.

STUDENT RESEARCH SUPPORT (2011 – 2023)

Graduate Research with Leveraged Funding:

- Using advanced morphometrics to improve identification of Sphaeriidae (fingernail clams) of the Great lakes as informed by DNA analyses (University of Minnesota Duluth; other field crews providing specimens).
- Importance of coastal wetlands to offshore fishes of the Great Lakes: Dietary support and habitat utilization (Central Michigan University; with additional funding from several small University grants and the US Fish and Wildlife Service).
- Spatial variation in macroinvertebrate communities within two emergent plant zones in Great Lakes coastal wetlands (Central Michigan University; with additional funding from CMU).
- Invertebrate co-occurrence patterns in coastal wetlands of the Great Lakes: Community assembly rules (Central Michigan University; additional funding from CMU)
- Functional indicators of Great Lakes coastal wetland health (University of Notre Dame; additional funding by Illinois-Indiana Sea Grant).

- Evaluating environmental DNA detection alongside standard fish sampling in Great Lakes coastal wetland monitoring (University of Notre Dame; additional funding by Illinois-Indiana Sea Grant).
- Nutrient-limitation in Great Lakes coastal wetlands (University of Notre Dame; additional funding by the UND College of Science).
- A summary of snapping turtle (*Chelydra serpentina*) by-catch records in Lake Ontario coastal wetlands (with additional funding by University of Toronto).
- Evaluating a zoobenthic indicator of Great Lakes wetland condition (with additional funding from University of Windsor).
- Testing and comparing the diagnostic value of three fish community indicators of Great Lakes wetland condition (with additional funding from GLRI GLIC: GLEI II and University of Windsor).
- Quantifying Aquatic Invasion Patterns Through Space and Time: A Relational Analysis of the Laurentian Great Lakes (University of Minnesota Duluth; with additional funding and data from USEPA)
- Novel Diagnostics for Biotransport of Aquatic Environmental Contaminants (University of Notre Dame, with additional funding from Advanced Diagnostics & Therapeutics program)
- Conservation of Common Terns in the Great Lakes Region (University of Minnesota; with additional funding from USFWS, MNDNR, and multiple smaller internal and external grants).
- Distribution of yellow perch in Great Lakes coastal wetlands (Grand Valley State University; with additional funding from GVSU).
- Variation in aquatic invertebrate assemblages in coastal wetland wet meadow zones of Lake Huron, of the Laurentian Great Lakes (University of Windsor; with additional funding from the University of Windsor).
- Influence of water level fluctuations and diel variation in dissolved oxygen concentrations on fish habitat use in Great Lakes coastal wetlands (University of Windsor; with additional funding from the University of Windsor).
- Bird community response to changes in wetland extent and lake level in Great Lakes coastal wetlands (University of Wisconsin-Green Bay with additional funding from Bird Studies Canada)
- Inferential measures for a quantitative ecological indicator of ecosystem health (University of Wisconsin-Green Bay)
- Per- and polyfluorinated alkyl substances (PFAS) in Great Lakes food webs and sportfish (University of Notre Dame)

Undergraduate Research with Leveraged Funding:

- Production of a short documentary film on Great Lakes coastal wetlands (University of Notre Dame; additional funding by the UND College of Arts and Letters).
- Heavy metal loads in freshwater turtle species inhabiting coastal wetlands of Lake Michigan (University of Notre Dame; additional funding by the UND College of Science, and ECI – Environmental Change Institute). [Online coverage](#), [TV](#) and [radio](#).
- Nitrogen-limitation in Lake Superior coastal wetlands (Northland College; additional funding from the Wisconsin DNR and Northland College).
- Patterns in chlorophyll-*a* concentrations in Great Lakes coastal wetlands (Northland College; additional funding provided by the college).
- *Phragmites australis* effects on coastal wetland nearshore fish communities of the Great Lakes basin (University of Windsor; with additional funding from GLRI GLIC: GLEI II).
- Sonar-derived estimates of macrophyte density and biomass in Great Lakes coastal wetlands (University of Windsor; with additional funding from GLRI GLIC: GLEI II presented at the International Association for Great Lakes Research annual meeting).
- Effects of disturbance frequency on the structure of coastal wetland macroinvertebrate communities (Lake Superior State University; with additional funding from LSSU's Undergraduate Research Committee; awarded Best Student Poster award at LSSU Research Symposium; presented at MI American Fisheries Society annual meeting).
- Resistance and resilience of macroinvertebrate communities in disturbed and undisturbed coastal wetlands (Lake Superior State University; with additional funding from LSSU's Undergraduate Research Committee, (presented at MI American Fisheries Society annual meeting and Midwest Fish and Wildlife Conference).
- Structure and function of restored Roxana Marsh in southern Lake Michigan (University of Notre Dame, with additional funding from the UND College of Science)
- Nutrient limitation in Great Lakes coastal wetlands (Central Michigan University, CMU Biological Station on Beaver Island)
- Effects of wetland size and adjacent land use on taxonomic richness (University of Minnesota Duluth, with additional funding from UMD's UROP program)
- Water depth optima and tolerances for St. Louis River estuary wetland plants (University of Wisconsin-Superior, with additional funding from WI Sea Grant)
- Mapping Wetland Areal Change in the St. Louis River Estuary Using GIS (University of Wisconsin-Superior, with additional funding from WI Sea Grant)
- An analysis of Microcystin concentrations in Great Lakes coastal wetlands (Central Michigan University; additional funding by CMU College of Science and Engineering).

- Bathymetry and water levels in lagoonal wetlands of the Apostle Islands National Lakeshore (Northland College; additional funding from the National Park Service). Several presentations at regional meetings and IAGLR.
- Non-native fish use of Great Lakes coastal wetlands (Northland College funding). Poster presentations by Northland College students at Wisconsin Wetland Science Meeting and IAGLR.

Graduate Research without Leveraged Funding:

- Impacts of drainage outlets on Great Lakes coastal wetlands (Central Michigan University).
- Effects of anthropogenic disturbance affecting coastal wetland vegetation (Central Michigan University).
- Great Lakes coastal wetland seed banks: what drives compositional change? (Central Michigan University).
- Spatial scale variation in patterns and mechanisms driving fish diversity in Great Lakes coastal wetlands (Central Michigan University).
- Building a model of macroinvertebrate functional feeding group community through zone succession: Does the River Continuum Concept apply to Great Lakes coastal wetlands? (Central Michigan University).
- Chemical and physical habitat variation within Great Lakes coastal wetlands; the importance of hydrology and dominant plant zonation (Central Michigan University)
- Macroinvertebrate-based Index of Biotic Integrity for Great Lakes coastal wetlands (Central Michigan University)
- Habitat conditions and invertebrate communities of Great Lakes coastal habitats dominated by Wet Meadow, and *Phragmites australis*: implications of macrophyte structure changes (Central Michigan University)
- The establishment of *Bithynia tentaculata* in coastal wetlands of the Great Lakes (Central Michigan University)
- Environmental covariates as predictors of anuran distribution in Great Lakes coastal wetlands (Central Michigan University)
- Impacts of muskrat herbivory in Great Lakes coastal wetlands (Central Michigan University).
- Mute swan interactions with native waterfowl in Great Lakes coastal wetlands (Central Michigan University).
- Effects of turbidity regimes on fish and macroinvertebrate community structure in coastal wetlands (Lake Superior State University and Oakland University).
- Scale dependence of dispersal limitation and environmental species sorting in Great Lakes wetland invertebrate meta-communities (University of Notre Dame).

- Spatial and temporal trends in invertebrate communities of Great Lakes coastal wetlands, with emphasis on Saginaw Bay of Lake Huron (University of Notre Dame).
- Model building and a comparison of the factors influencing sedge and marsh wren populations in Great Lakes coastal wetlands (University of Minnesota Duluth).
- The effect of urbanization on the stopover ecology of Neotropical migrant songbirds on the western shore of Lake Michigan (University of Minnesota Duluth).
- Assessing the role of nutrients and watershed features in cattail invasion (*Typha angustifolia* and *Typha x glauca*) in Lake Ontario wetlands (The College at Brockport).
- Developing captive breeding methods for bowfin (*Amia calva*) (The College at Brockport).
- Water chestnut (*Trapa natans*) growth and management in Lake Ontario coastal wetlands (The College at Brockport).
- Functional diversity and temporal variation of migratory land bird assemblages in lower Green Bay (University of Wisconsin-Green Bay).
- Effects of invasive *Phragmites* on stopover habitat for migratory shorebirds in lower Green Bay, Lake Michigan (University of Wisconsin-Green Bay).
- Plant species associations and assemblages for the whole Great Lakes, developed through unconstrained ordination analyses (Oregon State University).
- Genetic barcoding to identify black and brown bullheads (Grand Valley State University).
- Coastal wetland – nearshore linkages in Lake Michigan for sustaining sport fishes (University of Notre Dame)
- Anthropogenic disturbance effects on bird and anuran communities in Lake Ontario coastal wetlands (The College at Brockport)
- A fish-based index of biotic integrity for Lake Ontario coastal wetlands (The College at Brockport)
- Modeling potential nutria habitat in Great Lakes coastal wetlands (Central Michigan University)
- Modeling of Eurasian ruffe (*Gymnocephalus cernua*) habitat preferences to predict future invasions (University of Minnesota Duluth in collaboration with USEPA MED)
- Modeling species-specific habitat associations of Great Lakes coastal wetland birds (University of Minnesota)
- The effect of urbanization on the stopover ecology of Neotropical migrant songbirds on the western shore of Lake Michigan (University of Minnesota Duluth).
- Nutrient limitation in Great Lakes coastal wetlands: gradients and their influence (Central Michigan University; with additional funding from the CMU College of Science and Engineering)

- Invasive *Phragmites australis* management (Central Michigan University; with additional funding from the CMU College of Science and Technology)
- The relationship between vegetation and ice formation in Great Lakes coastal wetlands (Central Michigan University; with additional funding from CMU College of Science and Engineering)
- PFAS accumulation by Dressenidae *spp* in Great Lakes Coastal Wetlands (Central Michigan University)
- Development of a vegetation based IBI for Great Lakes Coastal Wetlands (Central Michigan University)
- Development of a model for Great-Lakes wide invasive plant harvest for bioenergy production and nutrient recycling (Loyola Chicago and Oregon State University)
- Updating the Macroinvertebrate-based Index of Biotic Integrity for Great Lakes coastal wetlands (Central Michigan University)
- Great Lakes coastal wetland bird and anuran habitat associations (UW-Green Bay)

Undergraduate Research without Leveraged Funding:

- Sensitivity of fish community metrics to net set locations: a comparison between Coastal Wetland Monitoring and GLEI methods (University of Minnesota Duluth).
- Larval fish usage and assemblage composition between different wetland types (Central Michigan University).
- Determining wetland health for selected Great Lakes Coastal Wetlands and incorporating management recommendations (Central Michigan University).
- Invertebrate co-occurrence trends in the wetlands of the Upper Peninsula and Western Michigan and the role of habitat disturbance levels (Central Michigan University).
- Is macroinvertebrate richness and community composition determined by habitat complexity or variation in complexity? (University of Windsor, complete; Published in *Ecosphere*).
- Modeling American coot habitat relative to faucet snail invasion potential (Central Michigan University).
- Nutrient uptake by *Phragmites australis* and native wetland plants (Central Michigan University).
- Comparison of the diagnostic accuracy two aquatic invertebrate field collection and laboratory sorting methods (University of Windsor, complete).
- Validation of a zoobenthic assemblage condition index for Great Lakes coastal wetlands (University of Windsor, complete).

- Water depth-related variation in net ecosystem production in a Great Lakes coastal wet meadow (University of Windsor, complete).
- Anuran habitat use in the Lower Green Bay and Fox River Area of Concern (University of Wisconsin-Green Bay with support from GLRI/AOC funding).
- Impacts of European frog-bit invasion on wetland macroinvertebrate communities (Lake Superior State University; presented at Midwest Fish and Wildlife Conference).
- Effects of European frog-bit on water quality and fish assemblages in St. Marys River coastal wetlands (Lake Superior State University; presented at Midwest Fish and Wildlife Conference).
- Functional diversity of macroinvertebrates in coastal wetlands along the St. Marys River (Lake Superior State University; awarded Best Student Poster award at LSSU Research Symposium; presented at Midwest Fish and Wildlife Conference).
- A comparison of macroinvertebrate assemblages in coastal wetlands exposed to varying wave disturbance (Lake Superior State University; presented at MI American Fisheries Society annual meeting).
- Coastal wetlands as nursery habitat for young-of-year fishes in the St. Marys River (Lake Superior State University; presented at MI American Fisheries Society annual meeting)
- Relationship between water level and fish assemblage structure in St. Marys River coastal wetlands (Lake Superior State University; presented at MI American Fisheries Society annual meeting)
- Dominance patterns in macroinvertebrate communities in Great Lakes coastal wetlands: does environmental stress lead to uneven community structure? Northland College.
- Understanding drivers of chlorophyll-a in Great Lakes coastal wetlands. University of Notre Dame
- Evaluating fish assemblage changes throughout the summer in St. Marys River coastal wetlands (Lake Superior State University)
- Quantifying litter decomposition in wetlands of varying condition (Lake Superior State University)

JOBS CREATED/RETAINED (2020)

- Principal Investigators (partial support): 22
- Post-doctoral researchers (partial support): 4
- Total graduate students supported on project (part-time): 19
- Unpaid undergraduate internship (summer): Not possible in 2020 due to Covid-19
- Undergraduate students (paid; summer and/or part-time): 21

- Technicians, jr. scientists (summer and/or partial support): 39
- Volunteers: Could not have volunteers in 2020 or 2021 due to Covid-19

Total jobs at least partially supported in 2020: 105.

Students and post-doctoral researchers trained in 2020: 44.

JOBS CREATED/RETAINED (CUMULATIVE SINCE 2011, LAST UPDATED 2020)

- Principal Investigators (partial support): 20 (average per year)
- Post-doctoral researchers (partial support; cumulative): 8
- Total graduate students supported on project (part-time; cumulative): 113
- Unpaid undergraduate internship (summer, cumulative): 35
- Undergraduate students (paid; summer and/or part-time; cumulative): 194
- Technicians, jr. scientists (summer and/or partial support; cumulative): 135
- Volunteers (cumulative): 47

Total jobs at least partially supported: 469.

Students and post-doctoral researchers trained: 349.

At our annual meetings in 2021 and 2023, we conducted a formal discussion session on Diversity, Equity, and Inclusion (DEI). In 2021, we split into 10 breakout groups to discuss three questions related to best practices for enhancing DEI in the CWMP workforce. In brief, the three questions concerned 1) current practices used to enhance DEI, 2) perceived barriers to enhancing DEI, and 3) potential mechanisms for enhancing DEI in the future. These discussion notes were compiled and organized, and then redistributed to all CWMP participants. In 2023 we focused our discussion on how to increase crew safety as field crews diversify, acknowledging that people from differing backgrounds, ethnicities, and identities may be treated differently and feel less safe. Our goal, as always, is for all field crew members to both feel and be safe. CWMP leadership will continue to monitor and encourage DEI goals for the program.

PRESENTATIONS ABOUT THE COASTAL WETLAND MONITORING PROJECT **(INCEPTION THROUGH 2023)**

Albert, Dennis. 2013. Use of Great Lakes Coastal Wetland Monitoring data in restoration projects in the Great Lakes region. 5th Annual Conference on Ecosystem Restoration, Schaumburg, IL. July 30, 2013. 20 attendees, mostly managers and agency personnel.

Albert, Dennis. 2013. Data collection and use of Great Lakes Coastal Wetland Monitoring data by Great Lakes restorationists. Midwestern State Wetland Managers Meeting, Kellogg Biological Station, Gull Lake, MI, October 31, 2013. 40 attendees; Great Lakes state wetland managers.

Albert, Dennis, N. Danz, D. Wilcox, and J. Gathman. 2014. Evaluating Temporal Variability of Floristic Quality Indices in Laurentian Great Lakes Coastal Wetlands. Society of Wetland Scientists, Portland, OR. June.

Albert, Dennis, et al. 2015. Restoration of wetlands through the harvest of invasive plants, including hybrid cattail and *Phragmites australis*. Presented to Midwestern and Canadian biologists. June.

Albert, Dennis, et al. 2015. Great-Lakes wide distribution of bulrushes and invasive species. Coastal and Estuarine Research Federation Conference in Portland, Oregon. November.

Amatangelo, K., D. Wilcox, R. Schultz, M. Altenritter, M. Chislock, and G. Lawrence. 2021. Application of the Great Lakes Coastal Wetlands Monitoring Program to Restoration Projects in Lake Ontario Wetlands. State of Lake Ontario Conference. March 9-11, 2021, online.

Baldwin, R., B. Currell, and A. Moerke. 2014. Effects of disturbance history on resistance and resilience of coastal wetlands. Midwest Fish and Wildlife Conference, January, Kansas City, MO.

Baldwin, R., B. Currell, and A. Moerke. 2014. Effects of disturbance history on resistance and resilience of coastal wetlands. MI American Fisheries Society annual meeting, February, Holland, MI.

Bergen, E., E. Shively, M.J. Cooper. Non-native fish species richness and distributions in Great Lakes coastal wetlands. International Association for Great Lakes Research Annual Conference, June 10-14, 2019, Brockport, NY. (poster)

Bergen, E., E. Shively, M.J. Cooper. Drivers of non-native fish species richness and distribution in the Laurentian Great Lakes. February 19-21, 2019. Madison, WI. (poster)

Bozimowski, S. and D.G. Uzarski. 2016. The Great Lakes coastal wetland monitoring program. 2016 Wetlands Science Summit, Richfield, OH. September, Oral Presentation.

Bozimowski, A.A., B.A. Murry, and D.G. Uzarski. 2012 Invertebrate co-occurrence patterns in the wetlands of northern and eastern Lake Michigan: the interaction of the harsh-benign hypothesis and community assembly rules. 55th International Conference on Great Lakes Research, Cornwall, Ontario.

Bozimowski, A. A., B. A. Murry, P. S. Kourtev, and D. G. Uzarski. 2014. Aquatic macroinvertebrate co-occurrence patterns in the coastal wetlands of the Great Lakes: the interaction of the harsh-benign hypothesis and community assembly rules. Great Lakes Science in Action Symposium, Central Michigan University, Mt. Pleasant, MI. April.

Bozimowski, A.A., B.A. Murry, P.S. Kourtev, and D.G. Uzarski. 2015. Aquatic macroinvertebrate co-occurrence patterns in the coastal wetlands of the Great Lakes. 58th International Conference on Great Lakes Research, Burlington, VT.

Bozimowski, A.A. and D.G. Uzarski. 2017. Monitoring a changing ecosystem: Great Lakes coastal wetlands. Saginaw Bay Watershed Initiative Network's State of the Bay Conference.

Bracey, A. M., R. W. Howe, N.G. Walton, E. E. G. Giese, and G. J. Niemi. Avian responses to landscape stressors in Great Lakes coastal wetlands. 5th International Partners in Flight Conference and Conservation Workshop. Snowbird, UT, August 25-28, 2013.

Brady, V., D. Uzarski, and M. Cooper. 2013. Great Lakes Coastal Wetland Monitoring: Assessment of High-variability Ecosystems. USEPA Mid-Continent Ecology Division Seminar Series, May 2013. 50 attendees, mostly scientists (INVITED).

Brady, V., G. Host, T. Brown, L. Johnson, G. Niemi. 2013. Ecological Restoration Efforts in the St. Louis River Estuary: Application of Great Lakes Monitoring Data. 5th Annual Conference on Ecosystem Restoration, Schaumburg, IL. July 30, 2013. 20 attendees, mostly managers and agency personnel.

Brady, V. and D. Uzarski. 2013. Great Lakes Coastal Wetland Fish and Invertebrate Condition. Midwestern State Wetland Managers Meeting, Kellogg Biological Station, Gull Lake, MI, October 31, 2013. 40 attendees; Great Lakes state wetland managers.

Brady, V., D. Uzarski, T. Brown, G. Niemi, M. Cooper, R. Howe, N. Danz, D. Wilcox, D. Albert, D. Tozer, G. Grabas, C. Ruetz, L. Johnson, J. Ciborowski, J. Haynes, G. Neuderfer, T. Gehring, J. Gathman, A. Moerke, G. Lamberti, C. Normant. 2013. A Biotic Monitoring Program for

Great Lakes Coastal Wetlands. Society of Wetland Scientists annual meeting, Duluth, MN, June 2013. 25 attendees, mostly scientists, some agency personnel.

Brady, V., D. Uzarski, T. Brown, G. Niemi, M. Cooper, R. Howe, N. Danz, D. Wilcox, D. Albert, D. Tozer, G. Grabas, C. Ruetz, L. Johnson, J. Ciborowski, J. Haynes, G. Neuderfer, T. Gehring, J. Gathman, A. Moerke, G. Lamberti, C. Normant. 2013. Habitat Values Provided by Great Lakes Coastal Wetlands: based on the Great Lakes Coastal Wetland Monitoring Project. Society of Wetland Scientists annual meeting, Duluth, MN, June 2013. 20 attendees, mostly scientists.

Brady, V.J., D.G. Uzarski, M.J. Cooper, D.A. Albert, N. Danz, J. Domke, T. Gehring, E. Giese, A. Grinde, R. Howe, A.H. Moerke, G. Niemi, H. Wellard-Kelly. 2018. How are Lake Superior's wetlands? Eight years, 100 wetlands sampled. State Of Lake Superior Conference. Houghton, MI. Oral Presentation.

Brady, V., G. Niemi, J. Dumke, H. Wellard Kelly, M. Cooper, N. Danz, R. Howe. 2019. The role of monitoring data in coastal wetland restoration: Case studies from Duluth and Green Bay. International Association of Great Lakes Research Annual Meeting, Brockport, NY, June 2019. Invited oral presentation.

Buckley, J.D., and J.J.H. Ciborowski. 2013. A comparison of fish indices of biological condition at Great Lakes coastal margins. 66th Canadian Conference for Freshwater Fisheries Research, Windsor, ON, January 3-5 2013. Poster Presentation.

Chorak, G.M., C.R. Ruetz III, R.A. Thum, J. Wesolek, and J. Dumke. 2015. Identification of brown and black bullheads: evaluating DNA barcoding. Poster presentation at the Annual Meeting of the Michigan Chapter of the American Fisheries Society, Bay City, Michigan. January 20-21.

Cooper, M.J. Great Lakes coastal wetland monitoring: chemical and physical parameters as co-variates and indicators of wetland health. Biennial State of the Lakes Ecosystem Conference, Erie, PA, October 26-27, 2011. Oral presentation.

Cooper, M.J. Coastal wetland monitoring: methodology and quality control. Great Lakes Coastal Wetland Monitoring Workshop, Traverse City, MI, August 30, 2011. Oral presentation.

Cooper, M.J., D.G. Uzarski, and G.L. Lamberti. GLRI: coastal wetland monitoring. Michigan Wetlands Association Annual Conference, Traverse City, MI, August 30-September 2, 2011. Oral presentation.

Cooper, M.J. Monitoring the status and trends of Great Lakes coastal wetland health: a basin-wide effort. Annual Great Lakes Conference, Institute of Water Research, Michigan State University, East Lansing, MI, March 8, 2011. Oral presentation.

Cooper, M.J., G.A. Lamberti, and D.G. Uzarski. Monitoring ecosystem health in Great Lakes coastal wetlands: a basin-wide effort at the intersection of ecology and management. Entomological Society of America, Reno, NV, November 13-16, 2011. Oral presentation

Cooper, M.J., and G.A. Lamberti. Taking the pulse of Great Lakes coastal wetlands: scientists tackle an epic monitoring challenge. Poster session at the annual meeting of the National Science Foundation Integrative Graduate Education and Research Traineeship Program, Washington, D.C., May 2012. Poster presentation.

Cooper, M.J., J.M. Kosiara, D.G. Uzarski, and G.A. Lamberti. Nitrogen and phosphorus conditions and nutrient limitation in coastal wetlands of Lakes Michigan and Huron. Annual meeting of the International Association for Great Lakes Research. Cornwall, Ontario. May 2012. Oral presentation.

Cooper, M.J., G.A. Lamberti, and D.G. Uzarski. Abiotic drivers and temporal variability of Saginaw Bay wetland invertebrate communities. International Association for Great Lakes Research, 56th annual meeting, West Lafayette, IN. June 2013. Oral presentation.

Cooper, M.J., D.G. Uzarski, J. Sherman, and D.A. Wilcox. Great Lakes coastal wetland monitoring program: support of restoration activities across the basin. National Conference on Ecosystem Restoration, Chicago, IL. July 2013. Oral presentation.

Cooper, M.J. and J. Kosiara. Great Lakes coastal wetland monitoring: Chemical and physical parameters as co-variates and indicators of wetland health. US EPA Region 5 Annual Wetlands Program Coordinating Meeting and Michigan Wetlands Association Annual Meeting. Kellogg Biological Station, Hickory Corners, MI. October 2013. Oral presentation.

Cooper, M.J. Implementing coastal wetland monitoring. Inter-agency Task Force on Data Quality for GLRI-Funded Habitat Projects. CSC Inc., Las Vegas, NV. November 2013. Web presentation, approximately 40 participants.

Cooper, M.J. Community structure and ecological significance of invertebrates in Great Lakes coastal wetlands. SUNY-Brockport, Brockport, NY. December 2013. Invited seminar.

Cooper, M.J. Great Lakes coastal wetlands: ecological monitoring and nutrient-limitation. Limno-Tech Inc., Ann Arbor, MI. December 2013. Invited seminar.

Cooper, M.J., D.G. Uzarski, and V.J. Brady. A basin-wide Great Lakes coastal wetland monitoring program: Measures of ecosystem health for conservation and management. Great Lakes Wetlands Day, Toronto, Ont. Canada, February 4, 2014. Oral presentation.

Cooper, M.J., G.A. Lamberti, and D.G. Uzarski. Supporting Great Lakes coastal wetland restoration with basin-wide monitoring. Great Lakes Science in Action Symposium. Central Michigan University. April 4, 2014.

Cooper, M.J. Expanding fish-based monitoring in Great Lakes coastal wetlands. Michigan Wetlands Association Annual Meeting. Grand Rapids, MI. August 27-29, 2014.

Cooper, M.J. Structure and function of Great Lakes coastal wetlands. Public seminar of Ph.D. dissertation research. University of Notre Dame. August 6, 2014.

Cooper, M.J., D.G. Uzarski, and T.N. Brown. Developing a decision support system for protection and restoration of Great Lakes coastal wetlands. Biodiversity without Borders Conference, NatureServe. Traverse City, MI. April 27, 2015.

Cooper, M.J. and D.G. Uzarski. Great Lakes coastal wetland monitoring for protection and restoration. Lake Superior Monitoring Symposium. Michigan Technological University. March 19, 2015.

Cooper, M.J. Where worlds collide: ecosystem structure and function at the land-water interface of the Laurentian Great Lakes. Central Michigan University Department of Biology. Public Seminar. February 5, 2015.

Cooper, M.J. Where worlds collide: ecosystem structure and function at the land-water interface of the Laurentian Great Lakes. Sigurd Olson Environmental Institute, Northland College. Public Seminar. May 4, 2015.

Cooper, M.J., and D.G. Uzarski. Great Lakes coastal wetland monitoring for protection and restoration. Lake Huron Restoration Meeting. Alpena, MI. May 14, 2015.

Cooper, M.J., D.G. Uzarski, and V.J. Brady. Developing a decision support system for restoration and protection of Great Lakes coastal wetlands. Wisconsin Wetlands Association Annual Meeting. February 24-25, 2016. Green Bay, WI.

Cooper, M.J., Stirratt, H., B. Krumwiede, and K. Kowalski. Great Lakes Resilient Lands and Waters Initiative, Deep Dive. Remote presentation to the White House Council on Environmental Quality and partner agencies, January 28, 2016.

Cooper, M., Redder, T., Brady, V. and D. Uzarski. 2016. Developing a decision support tool to guide restoration and protection of Great Lakes coastal wetlands. Annual Meeting of the Wisconsin Wetlands Association, Stevens Point, WI. February. Presentation.

Cooper, M.J.. Nutrient limitation in wetland ecosystems. Wisconsin Department of Natural Resources, February 12, 2016, Rhinelander, WI.

Cooper, M.J., D.G. Uzarski and V.J. Brady. 2016. Developing a decision support system for restoration and protection of Great Lakes coastal wetlands. Wisconsin Wetlands Association Annual Meeting, Green Bay, WI. February 24-25. Oral Presentation.

Cooper, M.J.. Monitoring biotic and abiotic conditions in Great Lakes coastal wetlands. Wisconsin DNR Annual Surface Water Quality Conference. May 2016, Tomahawk, WI.

Cooper, M.J. The Depth of Wisconsin's Water Resources. Panel Discussion, Wisconsin History Tour, Northern Great Lakes Visitors Center, June 15, 2016, Ashland, WI.

Cooper, M.J.. Great Lakes Coastal Wetlands. The White House Resilient Lands and Waters Initiative Roundtable. Washington, DC, November 17, 2016.

Cooper, M.J. Translating Science Into Action in the Great Lakes. Marvin Pertzik Lecture Series. Northland College, May 2016.

Cooper, M.C., C. Hippensteel, D.G. Uzarski, and T.M. Redder. Developing a decision support tool for Great Lakes coastal wetlands. LCC Coastal Conservation Working Group Annual Meeting, Great Lakes Environmental Research Laboratory, Ann Arbor, MI, Oct. 6, 2016.

Cooper, M.J., T.M. Redder, C. Hippensteel, V.J. Brady, D.G. Uzarski. Developing a decision support tool to guide restoration and protection of Great Lakes coastal wetlands. Midwest Fish and Wildlife Conference, Feb. 5-8, 2017, Lincoln, NE.

Cooper, M.J., T.M. Redder, V.J. Brady, D.G. Uzarski. Developing a decision support tool to guide restoration and protection of Great Lakes coastal wetlands. Wisconsin Wetlands Association Annual Conference, February 28-March 2, 2017, Steven's Point, WI.

Cooper, M.J. Coastal Wetlands as Metabolic Gates, Sediment Filters, Swiss Army Knife Habitats, and Biogeochemical Hotspots. Science on Tap, Ashland, WI, March 21, 2017.

Cooper, M.J., Brady, V.J., Uzarski, D.G., Lamberti, G.A., Moerke, A.H., Ruetz, C.R., Wilcox, D.A., Ciborowski, J.J.H., Gathman, J.P., Grabas, G.P., and Johnson, L.B. An Expanded Fish-Based Index of Biotic Integrity for Great Lakes Coastal Wetlands. International Association for Great Lakes Research 60th Annual Meeting, Detroit, MI, May 15-19, 2017.

Cooper, M.J., D.G. Uzarski, and A. Garwood. Great Lakes Coastal Wetland Monitoring.” Webinar hosted by Michigan Department of Environmental Quality, April 14, 2017. 78 attendees.

Cooper, M.J., A. Hefko, M. Wheeler. Nitrogen limitation of Lake Superior coastal wetlands. Society for Freshwater Science Annual Conference, May 20-24, 2018, Detroit, MI.

Cooper, M.J. The Role of Wetlands in Maintaining Water Quality. Briefing to the International Joint Commission, Ashland, WI, September 26, 2019.

Cooper, M.J., V.J. Brady, and D.G. Uzarski. Great Lakes Coastal Wetland Monitoring. Plenary Presentation, Great Lakes Coastal Wetland Symposium, Oregon, OH, September 19, 2019.

Cooper, M.J. and S. Johnson. Life on the Soggy Edges. Madeline Island Wilderness Preserve Lecture Series, Madeline Island Museum, La Pointe, WI, June 19, 2019.

Cooper, M.J., T.M. Redder, V.J. Brady, D.G. Uzarski. A data visualization tool to support protection and restoration of Great Lakes coastal wetlands. International Association for Great Lakes Research Annual Conference, June 10-14, 2019, Brockport, NY

Cooper, M.J., V.J. Brady, and D.G. Uzarski. 2022. Detecting Human Disturbance in Coastal Wetlands Across Temporal and Spatial Scales Using Biotic Indicators. Great Lakes Coastal Symposium. Sept. 19-21, 2022. Sault Ste. Marie, MI

Cooper, M.J., V.J. Brady, and D.G. Uzarski. 2023. Monitoring Great Lakes Coastal Wetlands. Michigan Wetlands Association Annual Meeting. Sept. 12-14, 2023. Kalamazoo, MI

Curell, Brian. 2014. Effects of disturbance frequency on macroinvertebrate communities in coastal wetlands. MI American Fisheries Society annual meeting, February, Holland, MI.

Dahlberg, N., N.P. Danz, and S. Schooler. 2015. Integrating prior vegetation surveys from the St. Louis River estuary. Poster presentation at the 2015 Annual St. Louis River Summit, Superior, WI.

Dahlberg, N., N.P. Danz, and S. Schooler. 2017. 2012 Flood Impacts on St. Louis River Plant Communities. Poster presentation at St. Louis River Summit, Superior, WI.

Danz, N.P. 2014. Floristic quality of Wisconsin coastal wetlands. Oral presentation at the Wisconsin Wetlands Association 19th Annual Wetlands Conference, LaCrosse, WI. Audience mostly scientists.

Danz, N.P. Floristic Quality of Coastal and Inland Wetlands of the Great Lakes Region. Invited presentation at the University of Minnesota Duluth, Duluth, MN.

Danz, N.P., S. Schooler, and N. Dahlberg. 2015. Floristic quality of St. Louis River estuary wetlands. Oral presentation at the 2015 Annual St. Louis River Summit, Superior, WI.

Danz, N.P. 2016. Floristic quality of St. Louis River estuary wetlands. Invited presentation at the Center for Water and the Environment, Natural Resources Research Institute, Duluth, MN.

Danz, N.P. 2017. Connections Between Human Stress, Wetland Setting, and Vegetation in the St. Louis River Estuary. Oral presentation at the Wetland Science Conference, Stevens Point, WI.

Danz, N.P. 2017. 10 Things We Learned from Your Vegetation Data. Oral presentation at the St. Louis River Summit, Superior, WI.

Daly, D., T. Dunn, and A. Moerke. 2016. Effects of European frog-bit on water quality and fish assemblages in St. Marys River wetlands. Midwest Fish and Wildlife Conference, Grand Rapids, MI. January 24-27.

Des Jardin, K. and D.A. Wilcox. 2014. Water chestnut: germination, competition, seed viability, and competition in Lake Ontario. New York State Wetlands Forum, Rochester, NY.

Dumke, J.D., V.J. Brady, J. Ciborowski, J. Gathman, J. Buckley, D. Uzarski, A. Moerke, C. Ruetz III. 2013. Fish communities of the upper Great Lakes: Lake Huron's Georgian Bay is an outlier. Society for Wetland Scientists, Duluth, Minnesota. 30 attendees, scientists and managers.

Dumke, J.D., V.J. Brady, R. Hell, A. Moerke, C. Ruetz III, D. Uzarski, J. Gathman, J. Ciborowski. 2013. A comparison of St. Louis River estuary and the upper Great Lakes fish communities (poster). Minnesota American Fisheries Society, St. Cloud, Minnesota. Attendees scientists, managers, and agency personnel.

Dumke, J.D., V.J. Brady, R. Hell, A. Moerke, C. Ruetz III, D. Uzarski, J. Gathman, J. Ciborowski. 2013. A comparison of wetland fish communities in the St. Louis River estuary and the upper Great Lakes. St. Louis River Estuary Summit, Superior, Wisconsin. 150 attendees, including scientists, managers, agency personnel, and others.

Dumke, J.D., V.J. Brady, J. Erickson, A. Bracey, N. Danz. 2014. Using non-degraded areas in the St. Louis River estuary to set biotic delisting/restoration targets. St. Louis River Estuary Summit, Superior, Wisconsin. 150 attendees, including scientists, managers, agency personnel, and others.

Dumke, J., C.R. Ruetz III, G.M. Chorak, R.A. Thum, and J. Wesolek. 2015. New information regarding identification of young brown and black bullheads. Oral presentation at the

Annual Meeting of the Wisconsin Chapter of the American Fisheries Society, Eau Claire, Wisconsin. February 24-26. 150 attendees, including scientists, managers, agency personnel, and others.

Dunn, T., D. Daly, and A. Moerke. 2016. Impacts of European frog-bit invasion on Great Lakes wetlands macroinvertebrate communities. Midwest Fish and Wildlife Conference, Grand Rapids, MI. January 24-27.

Dykstra, K.M., C.R. Ruetz III, M.J. Cooper, and D.G. Uzarski. 2018. Occupancy and detection of yellow perch in Great Lakes coastal wetlands. Poster presentation at the Annual Meeting of the Society for Freshwater Science, Detroit, Michigan. May 20-24.

Dykstra (Emelander), K.M., C.R. Ruetz III, M.J. Cooper, and D.G. Uzarski. 2018. Occupancy and detection of yellow perch in Great Lakes coastal wetlands: preliminary results. Poster presentation at the annual meeting of the Michigan Chapter of the American Fisheries Society, Port Huron, Michigan. February 13-14.

Elliot, L.H., A.M. Bracey, G.J. Niemi, D.H. Johnson, T.M. Gehring, E.E. Gnass Giese, G.P. Grabas, R.W. Howe, C.J. Norment, and D.C. Tozer. Habitat Associations of Coastal Wetland Birds in the Great Lakes Basin. American Ornithological Society Meeting, East Lansing, Michigan. Poster Presentation. 31 July-5 August 2017.

Elliott, L.H., A. Bracey, G. Niemi, D.H. Johnson, T. Gehring, E. Giese, G. Grabas, R. Howe, C. Norment, and D.C. Tozer. 2018. Hierarchical modeling to identify habitat associations of secretive marsh birds in the Great Lakes. IAGLR Conference, Toronto, Canada. Oral Presentation. 18-22 June 2018.

Fraley, E.F. and D.G. Uzarski 2017. The relationship between vegetation and ice formation in Great Lakes coastal wetlands. 60th Annual Meeting of the International Association of Great Lakes Research. Detroit, MI. Poster.

Fraley, E.F. and D.G. Uzarski. 2016. The Impacts of Ice on Plant Communities in Great Lakes Coastal Wetlands. 7th Annual Meeting of the Michigan Consortium of Botanists, Grand Rapids, MI. October. Poster.

Gathman, J.P. 2013. How healthy are Great Lakes wetlands? Using plant and animal indicators of ecological condition across the Great Lakes basin. Presentation to Minnesota Native Plant Society. November 7, 2013.

Gathman, J.P., J.J.J. Ciborowski, G. Grabas, V. Brady, and K.E. Kovalenko. 2013. Great Lakes Coastal Wetland Monitoring project: progress report for Canada. 66th Canadian Conference for Freshwater Fisheries Research, Windsor, ON, January 3-5, 2013. Poster Presentation.

- Gilbert, J.M., N. Vidler, P. Cloud Sr., D. Jacobs, E. Slavik, F. Letourneau, K. Alexander. 2014. *Phragmites australis* at the crossroads: Why we cannot afford to ignore this invasion. Great Lakes Wetlands Day Conference, Toronto, ON, February 4, 2014.
- Gilbert, J.M. 2013. *Phragmites* Management in Ontario. Can we manage without herbicide? Webinar, Great Lakes *Phragmites* Collaborative, April 5, 2013.
- Gilbert, J.M. 2012. *Phragmites australis*: a significant threat to Laurentian Great Lakes Wetlands, Oral Presentation, International Association of Great Lakes Wetlands, Cornwall, ON, May 2012
- Gilbert, J.M. 2012. *Phragmites australis*: a significant threat to Laurentian Great Lakes Wetlands, Oral Presentation to Waterfowl and Wetlands Research, Management and Conservation in the Lower Great Lakes. Partners' Forum, St. Williams, ON, May 2012.
- Gil de LaMadrid, D., and N.P. Danz. 2015. Water depth optima and tolerances for St. Louis River estuary wetland plants. Poster presentation at the 2015 Annual St. Louis River Summit, Superior, WI.
- Gnass Giese, E.E. 2015. Great Lakes Wetland Frog Monitoring. Annual Lower Fox River Watershed Monitoring Program Symposium at the University of Wisconsin-Green Bay, Green Bay, Wisconsin. April 14, 2015. Oral Presentation.
- Gnass Giese, E.E. 2015. Wetland Birds and Amphibians: Great Lakes Monitoring. Northeastern Wisconsin Audubon Society meeting at the Bay Beach Wildlife Sanctuary, Green Bay, Wisconsin. February 19, 2015. Oral Presentation.
- Gnass Giese, E.E., R.W. Howe, N.G. Walton, G.J. Niemi, D.C. Tozer, W.B. Gaul, A. Bracey, J. Shrovnal, C.J. Norment, and T.M. Gehring. 2016. Assessing wetland health using breeding birds as indicators. Wisconsin Wetlands Association Conference, Radisson Hotel & Convention Center, Green Bay, Wisconsin. February 24, 2016. Poster Presentation.
- Gnass Giese, E., R. Howe, A. Wolf, and G. Niemi. 2017. Breeding Birds and Anurans of Dynamic Green Bay Coastal Wetlands. State of Lake Michigan Conference, Green Bay, Wisconsin. Oral Presentation. 8 November 2017. Gnass Giese, E.E., R.W. Howe, A.T. Wolf, N.A. Miller, and N.G. Walton. An ecological index of forest health based on breeding birds. 2013. Webpage: <http://www.uwgb.edu/biodiversity/forest-index/>
- Gnass Giese, E.E., R.W. Howe, A.T. Wolf, N.A. Miller, and N.G. Walton. 2014. Using Bird Data to Assess Condition of Western Great Lakes Forests. Midwest Bird Conservation and Monitoring Workshop, Port Washington, Wisconsin. Poster Presentation. 4-8 August 2014. Gnass Giese, E.E. 2013. Monitoring forest condition using breeding birds in the

western Great Lakes region, USA. Editors: N. Miller, R. Howe, C. Hall, and D. Ewert. Internal Report. Madison, WI and Lansing, MI: The Nature Conservancy. 44 pp.

Gunsch, D., J.P. Gathman, and J.J.H. Ciborowski . 2018. Variation in dissolved-oxygen profiles along a depth gradient in Lake Huron coastal wet meadows relative to vegetation density and agricultural stress over 24 hours. IAGLR Conference, Toronto, Canada. Poster Presentation. 18-22 June 2018.

Gurholt, C.G. and D.G. Uzarski. 2013. Into the future: Great Lakes coastal wetland seed banks. IGLR Graduate Symposium, Central Michigan University, Mt. Pleasant, MI. March.

Gurholt, C.G. and D.G. Uzarski. 2013. Seed Bank Purgatory: What Drives Compositional Change of Great Lakes Coastal Wetlands. 56th International Association for Great Lakes Research Conference, Purdue University, West Lafayette, IN. June.

Harrison, A.M., M.J. Cooper, and D.G. Uzarski. 2019. Spatial and temporal (2011-2018) variation of water quality in Great Lakes coastal wetlands. International Association for Great Lakes Research. Brockport, NY. Presentation.

Hefko, A.G., M. Wheeler, M.J. Cooper. Nitrogen limitation of algal biofilms in Lake Superior coastal wetlands. International Association for Great Lakes Research Annual Conference, June 10-14, 2019, Brockport, NY. (poster)

Hein, M.C. and Cooper, M.J. Untangling drivers of chlorophyll a in Great Lakes coastal wetlands. International Association for Great Lakes Research 60th Annual Meeting, Detroit, MI, May 15-19, 2017.

Hirsch, B. E.E. Gnass Giese, and R. Howe. 2021. Anuran Occurrences in High and Low Water within the Lower Green Bay & Fox River AOC. Wisconsin Wetlands Association Conference, Virtual. Poster Presentation. February 2021.

Hohman, T., B. Howe, E. Giese, A. Wolf, and D. Tozer. 2019. Bird Community Response to Changes in Wetland Extent and Interspersion in Great Lakes Coastal Wetlands. Heckrodt Birding Club Meeting, Menasha, Wisconsin. Oral Presentation. 6 August 2019.

Hohman, T.R., R.W. Howe, A.T. Wolf, E.E.Gnass Giese, D.C. Tozer, T.M. Gehring, G.P. Grabas, G.J. Niemi, and C.J. Norment. 2019. Bird Community Response to Changes in Wetland Extent and Interspersion in Great Lakes Coastal Wetlands. Presented at the 62nd Annual Meeting of the International Association of Great Lakes Research (IAGLR), 12 June 2019, Brockport, NY.

Houghton, C.J., C.C. Moratz, P.S. Forsythe, G.A. Lamberti, D.G. Uzarski, and M.B. Berg. 2016. Relative use of wetland and nearshore habitats by sportfishes of Green Bay. 59th

International Conference on Great Lakes Research, Guelph, Ontario Canada. May. Oral Presentation.

Howe, R.W., R.P. Axler, V.J. Brady, T.N. Brown, J.J.H. Ciborowski, N.P. Danz, J.P. Gathman, G.E. Host, L.B. Johnson, K.E. Kovalenko, G.J. Niemi, and E.D. Reavie. 2012. Multi-species indicators of ecological condition in the coastal zone of the Laurentian Great Lakes. 97th Annual Meeting of the Ecological Society of America. Portland, OR.

Howe, B., E. Giese, A. Wolf, and B. Kupsky. 2019. Restoration Targets for Great Lakes Coastal Wetlands in the Lower Green Bay & Fox River AOC. International Association for Great Lakes Research, Brockport, New York. Oral Presentation. 12 June 2019.

Howe, R.W., G.J. Niemi, N.G. Walton, E.E.G. Giese, A.M. Bracey, V.J. Brady, T.N. Brown, J.J.H. Ciborowski, N.P. Danz, J.P. Gathman, G.E. Host, L.B. Johnson, K.E. Kovalenko, and E.D. Reavie. 2014. Measurable Responses of Great Lakes Coastal Wetland Biota to Environmental Stressors. International Association for Great Lakes Research Annual Conference, Hamilton, Ontario (Canada). May 26-30, 2014. Oral Presentation.

Howe, B., A. Wolf, E. Giese, V. Pappas, B. Kupsky, M. Grimm, and N. Van Helden. 2018. Lower Green Bay & Fox River Area of Concern Wildlife and Habitat Assessment Tools. AOC RAP Meeting, Green Bay, Wisconsin. Oral Presentation. 25 April 2018.

Howe, B., A. Wolf, E. Giese, V. Pappas, B. Kupsky, M. Grimm, and N. Van Helden. 2018. Assessing the Fish and Wildlife Habitat BUI for the Lower Green Bay and Fox River Area of Concern. Annual Great Lakes Areas of Concern Conference, Sheboygan, Wisconsin. Oral Presentation. 16 May 2018.

Howe, R.W., A.T. Wolf, and E.E. Gnass Giese. 2016. What's so special about Green Bay wetlands? Wisconsin Wetlands Association Conference, Radisson Hotel & Convention Center, Green Bay, Wisconsin. February 23-25, 2016. Oral Presentation.

Howe, R.W., N.G. Walton, E.G. Giese, G.J. Niemi, and A.M. Bracey. 2013. Avian responses to landscape stressors in Great Lakes coastal wetlands. Society of Wetland Scientists, Duluth, Minnesota. June 2-6, 2013. Poster Presentation.

Howe, R.W., N.G. Walton, E.E.G. Giese, G.J. Niemi, N.P. Danz, V.J. Brady, T.N. Brown, J.J.H. Ciborowski, J.P. Gathman, G.E. Host, L.B. Johnson, E.D. Reavie. 2013. How do different taxa respond to landscape stressors in Great Lakes coastal wetlands? Ecological Society of America, Minneapolis, Minnesota. August 4-9, 2013. Poster Presentation.

Howe, R.W., A.T. Wolf, J. Noordyk, and J. Stoll. 2017. Benefits and outcomes of Green Bay restoration: ecosystem and economic perspectives. Presented at the Summit on the

Ecological and Socio-Economic Tradeoffs of Restoration in the Green Bay, Lake Michigan, Ecosystem (July 18-20, 2017).

Howe, R.W., A.T. Wolf, and E.E. Giese. 2016. Proposed AOC de-listing process. Presentation to Lower Green Bay and Fox River AOC stakeholders. 16 December 2016.

Howe, R.W., A.T. Wolf, and E.E. Giese. 2017. Lower Green Bay & Fox River Area of Concern: A Plan for Delisting Fish and Wildlife Habitat & Populations Beneficial Use Impairments. A paper presented to AOC Technical Advisory Group. 3 August 2017.

Johnson, L., M. Cai, D. Allan, N. Danz, D. Uzarski. 2015. Use and interpretation of human disturbance gradients for condition assessment in Great Lakes coastal ecosystems. International Association for Great Lakes Research Conference, Burlington, VT.

Johnson, Z., M. Markel, and A. Moerke. 2019. Functional diversity of macroinvertebrates in coastal wetlands along the St. Marys River. Midwest Fish and Wildlife Conference, Cleveland, OH.

Kneisel, A.N., M.J. Cooper, and D.G. Uzarski. 2016. The impact of *Phragmites australis* invasion on macroinvertebrate communities in the coastal wetlands of Thunder Bay, MI. Institute for Great Lakes Research, 4th Annual Student Research Symposium, Central Michigan University, Mt. Pleasant, MI. February. Oral Presentation.

Kneisel, A.N., M.J. Cooper, and D.G. Uzarski. 2016. Impact of *Phragmites* invasion on macroinvertebrate communities in wetlands of Thunder Bay, MI. 59th International Conference on Great Lakes Research, Guelph, Ontario Canada. May. Oral Presentation.

Kosiara, J.M., M.J. Cooper, D.G. Uzarski, and G.A. Lamberti. 2013. Relationships between community metabolism and fish production in Great Lakes coastal wetlands. International Association for Great Lakes Research, 56th annual meeting. June 2-6, 2013. West Lafayette, IN. Poster presentation.

Kneisel, A.N., M.J. Cooper, and D.G. Uzarski. 2017. The impact of *Phragmites australis* invasion on Great Lakes coastal wetlands. 60th International Conference on Great Lakes Research, Detroit, MI. May. Presentation.

Kneisel, A.K., M.J. Cooper, D.G. Uzarski. 2018. Coastal wetland monitoring data as a resource for invasive species management. ELLS-IAGLR Big Lakes Small World Conference. Évian, France. September. Poster. Kosiara, J.K., J.J. Student, and D.G. Uzarski. 2017. Exploring coastal habitat-use patterns of Great Lakes yellow perch with otolith microchemistry. 60th International Conference on Great Lakes Research, Detroit, MI. May. Presentation.

Kosiara, J.M., J. Student and D.G. Uzarski. 2016. Assessment of yellow perch movement between coastal wetland and nearshore waters of the Great Lakes. 59th International Conference on Great Lakes Research, Guelph, Ontario Canada. May. Oral Presentation.

Kowalke, C.J. and D.G. Uzarski. 2019. Assessing the competitive impacts of invasive round goby on lake whitefish in northern Lake Michigan. International Association for Great Lakes Research. Brockport, NY. Poster.

Lamberti, G.A., D.G. Uzarski, V.J. Brady, M.J. Cooper, T.N. Brown, L.B. Johnson, J.J. Ciborowski, G.P. Grabas, D.A. Wilcox, R.W. Howe, and D. C. Tozer. An integrated monitoring program for Great Lakes coastal wetlands. Society for Freshwater Science Annual Meeting. Jacksonville, FL. May 2013. Poster presentation.

Lamberti, G.A. Pacific Salmon in Natal Alaska and Introduced Great Lakes Ecosystems: The Good, the Bad, and the Ugly. Department of Biology, Brigham Young University. Dec 5, 2013. Invited seminar.

Lamberti, G. A. The Global Freshwater Crisis. The Richard Stockton College of New Jersey and South Jersey Notre Dame Club. November 18, 2014.

Lamberti, G. A. The Global Freshwater Crisis. Smithsonian Journey Group and several University Alumni Groups. March 1, 2015.

Lamberti, G.A. The Global Freshwater Crisis. Newman University and Notre Dame Alumni Club of Wichita. September 28, 2016.

Lamberti, G.A. The Global Freshwater Crisis. Air and Wastewater Management Association and Notre Dame Alumni Club of Northeastern New York. December 2, 2016.

Lamberti, G.A. The Global Freshwater Crisis: Lessons for the Amazon. Association of University Alumni Clubs. Iquitos, Peru. September 9, 2019.

Lamberti, G. A. Pacific Salmon in Natal Alaska and Introduced Great Lakes Ecosystems: The Good, the Bad, and the Ugly. Annis Water Resources Institute, Grand Valley State University. December 12, 2014.

Lamberti, G.A., M.A. Brueseke, W.M. Conard, K.E. O'Reilly, D.G. Uzarski, V.J. Brady, M.J. Cooper, T.M. Redder, L.B. Johnson, J.H. Ciborowski, G.P. Grabas, D.A. Wilcox, R.W. Howe, D.C. Tozer, and T.K. O'Donnell. Great Lakes Coastal Wetland Monitoring Program: Vital resources for scientists, agencies and the public. Society for Freshwater Science Annual Meeting. Raleigh, NC. June 4-9, 2017. Poster.

- Langer, T.A., K. Pangle, B.A. Murray, and D.G. Uzarski. 2014. Beta Diversity of Great Lakes Coastal Wetland Communities: Spatiotemporal Structuring of Fish and Macroinvertebrate Assemblages. American Fisheries Society, Holland, MI. February.
- Langer, T., K. Pangle, B. Murray, D. Uzarski. 2013. Spatiotemporal influences, diversity patterns and mechanisms structuring Great Lakes coastal wetland fish assemblages. Poster. Institute for Great Lakes Research 1st Symposium, MI. March.
- Lemein, T.J., D.A. Albert, D.A. Wilcox, B.M. Mudrzynski, J. Gathman, N.P. Danz, D. Rokitnicki-Wojcik, and G.P. Grabas. 2014. Correlation of physical factors to coastal wetland vegetation community distribution in the Laurentian Great Lakes. Society of Wetland Scientists/Joint Aquatic Sciences Meeting, Portland, OR.
- MacDonald, J.L., L.S. Schoen, J.J. Student, and D.G. Uzarski. 2016. Variation in yellow perch (*Perca flavescens*) growth rate in the Great Lakes. 59th International Conference on Great Lakes Research, Guelph, Ontario Canada. May. Oral Presentation.
- Makish, C.S., K.E. Kovalenko, J.P. Gathman, and J.J.H. Ciborowski. 2013. invasive phragmites effects on coastal wetland fish communities of the Great Lakes basin. 66th Canadian Conference for Freshwater Fisheries Research, Windsor, ON, January 3-5, 2013. Poster Presentation.
- Markel, M., Z. Johnson, and A. Moerke. 2019. A comparison of macroinvertebrate assemblages in coastal wetlands exposed to varying wave disturbance. March 13-15, Gaylord, MI.
- McReynolds, A.T., K.E. O'Reilly, and G.A. Lamberti. 2016. Food web structure of a recently restored Indiana wetland. University of Notre Dame College of Science Joint Annual Meeting, Notre Dame, IN.
- Miranda, D.A., Zachritz, A.M., Whitehead, H.D., Peaslee, G.F., Cressman, S. R., Lamberti, G.A. PFAS Permeates Native and Introduced Salmonids from Lake Michigan, USA. Joint Aquatic Sciences Meeting, Grand Rapids, MI. May 2022.
- Miranda, D.A., Zachritz, A.M., Whitehead, H.D., Peaslee, G.F., Cressman, S. R., Lamberti, G.A., A Survey of Sportfish for Per-and Polyfluoroalkyl Substances (PFAS): An Emerging Contaminant in the Great Lakes. Portage, IN, October 2022
- Miranda, D.A., Zachritz, A.M., Whitehead, H.D., Peaslee, G.F., Cressman, S. R., Lamberti, G.A. "PFAS in Prey and Predator Fish from Lake Michigan", USA. SETAC North America 43 rd. Annual Meeting. November 2022.
- Miranda, D.A., Zachritz, A.M., Whitehead, H.D., Cressman, S., Klepinger, S., Peaslee, G.F. Lamberti, G.A. "Biomagnification of PFAS in Lake Michigan food web". Colleges of Science

and Engineering Joint Annual Meeting, Notre Dame IN. December 9, 2022.

Miranda, D.A., PFAS in Lake Michigan Fish, Annual Great Lakes Conference, Institute of Water Research– Michigan State University MI. March 7, 2023.

Moerke, A. 2015. Coastal wetland monitoring in the Great Lakes. Sault Naturalist meeting, Sault Sainte Marie, MI; approximately 40 community members present.

Monks, A., S. Lishawa, D. Albert, B. Mudrzynski, D.A. Wilcox, and K. Wellons. 2019. Innovative management of European frogbit and invasive cattail. International Association for Great Lakes Research. Brockport, NY

Moore, L.M., M.J. Cooper, and D.G. Uzarski. 2017. Nutrient limitation in Great Lakes coastal wetlands: gradients and their influence. 60th International Conference on Great Lakes Research, Detroit, MI. May 17. Presentation.

Mudrzynski, B.M., N.P. Danz, D.A. Wilcox, D.A. Albert, D. Rokitnicki-Wojcik, and J. Gathman. 2016. Great Lakes wetland plant Index of Biotic Integrity (IBI) development: balancing broad applicability and accuracy. Society of Wetland Scientists, Corpus Christi, TX.

Mudrzynski, B.M., D.A. Wilcox, and A. Heminway. 2012. Habitats invaded by European frogbit (*Hydrocharis morsus-ranae*) in Lake Ontario coastal wetlands. INTECOL/Society of Wetland Scientists, Orlando, FL.

Mudrzynski, B.M., D.A. Wilcox, and A.W. Heminway. 2013. European frogbit (*Hydrocharis morsus-ranae*): current distribution and predicted expansion in the Great Lakes using niche-modeling. Society of Wetland Scientists, Duluth, MN.

Mudrzynski, B.M. and D.A. Wilcox. 2014. Effect of coefficient of conservatism list choice and hydrogeographic type on floristic quality assessment of Lake Ontario wetlands. Society of Wetland Scientists/Joint Aquatic Sciences Meeting, Portland, OR.

Mudrzynski, B.M., K. Des Jardin, and D.A. Wilcox. 2015. Predicting seed bank emergence within flooded zones of Lake Ontario wetlands under novel hydrologic conditions. Society of Wetlands Scientists. Providence, RI.

Newman, W.L., L.P. Moore, M.J. Cooper, D.G. Uzarski, and S.N. Francoeur. 2019. Nitrogen-Fixing Diatoms as Indicators of Historical Nitrogen Limitation in Laurentian Great Lakes Coastal Wetlands. Society for Freshwater Science. Salt Lake City, UT. Presentation.

O'Donnell, T.K., Winter, C., Uzarski, D.G., Brady, V.J., and Cooper, M.J. 2017. Great Lakes coastal wetland monitoring: moving from assessment to action. Ecological Society of America Annual Conference. Portland, OR. August 6-11. Presentation.

O'Donnell, T.K., D.G. Uzarski, V.J. Brady, and M.J. Cooper. 2016. Great Lakes Coastal Wetland Monitoring: Moving from Assessment to Action. 10th National Monitoring Conference; Working Together for Clean Water, Tampa, Florida. May. Oral Presentation.

O'Reilly, K.E., A. McReynolds, and G.A. Lamberti. Quantifying Lake Michigan coastal wetland-nearshore linkages for sustaining sport fishes using stable isotope mixing models. Annual Meeting of the Ecological Society of America. Baltimore, MD. August 9-14, 2015.

O'Reilly, K.E., A. McReynolds, C. Stricker, and G.A. Lamberti. Quantifying Lake Michigan coastal wetland-nearshore linkages for sustaining sport fishes. State of Lake Michigan Conference. Traverse City, MI. October 28-30, 2015.

O'Reilly, K.E., A. McReynolds, C. Stricker, and G.A. Lamberti. 2016. Quantifying Lake Michigan coastal wetland-nearshore linkages for sustaining sport fishes. Society for Freshwater Science, Sacramento, CA.

O'Reilly, K.E., A. McReynolds, C. Stricker, and G.A. Lamberti. 2016. Quantifying Lake Michigan coastal wetland-nearshore linkages for sustaining sport fishes. International Association for Great Lakes Research, Guelph, ON.

O'Reilly, K.E., J.J. Student, B.S. Gerig, and G.A. Lamberti. 2019. Metalheads: What can sport fish otoliths reveal about heavy metal exposure over time? Annual Meeting of the Society for Freshwater Science, Salt Lake City, UT.

Otto, M., J. Marty, E.G. Gnass Giese, R. Howe, and A. Wolf. Anuran habitat use in the Lower Green Bay and Fox River Area of Concern (Wisconsin). University of Wisconsin-Green Bay Academic Excellence Symposium, Green Bay, Wisconsin. April 6, 2017. Poster Presentation.

Otto, M., J. Marty, E.G. Gnass Giese, R. Howe, and A. Wolf. Anuran habitat use in the Lower Green Bay and Fox River Area of Concern (Wisconsin). Green Bay Conservation Partners Spring Roundtable Meeting, Green Bay, Wisconsin. April 25, 2017. Poster Presentation.

Redder, T.M., D.G. Uzarski, V.J. Brady, M.J. Cooper, and T.K. O'Donnell. 2018. Application of data management and decision support tools to support coastal wetland management in the Laurentian Great Lakes. National Conference on Ecosystem Restoration. New Orleans, LA. August 26-30, 2018. Oral Presentation.

Reisinger, L. S., Pangle, K. L., Cooper, M. J., Learman, D. R., Uzarski, D. G., Woolnough, D. A., Bugaj, M. R., Burck, E. K., Dollard, R. E., Goetz, A., Goss, M., Gu, S., Karl, K., Rose, V. A.,

Scheunemann, A. E., Webster, R., Weldon, C. R., and J., Yan. 2017. The influence of water currents on community and ecosystem dynamics in coastal Lake Michigan. 60th International Conference on Great Lakes Research, Detroit, MI. May. Presentation.

Reisinger, A. J., and D. G., Uzarski. 2017. Natural and anthropogenic disturbances affect water quality of Great Lakes coastal wetlands. 60th International Conference on Great Lakes Research, Detroit, MI. May. Presentation.

St.Pierre, J.I., K.E. Kovalenko, A.K. Pollock, and J.J.H. Ciborowski. 2013. Is macroinvertebrate richness and community composition determined by habitat complexity or variation in complexity? 66th Canadian Conference for Freshwater Fisheries Research, Windsor, ON, January 3-5, 2013. Poster Presentation.

Schmidt, N. C., Schock, N., and D. G. Uzarski. 2013. Modeling macroinvertebrate functional feeding group assemblages in vegetation zones of Great Lakes coastal wetlands. International Association for Great Lakes Research Conference, West Lafayette, IN. June.

Schmidt, N.C., N.T. Schock, and D.G. Uzarski. 2014. Influences of metabolism on macroinvertebrate community structure across Great Lakes coastal wetland vegetation zones. Great Lakes Science in Action Symposium, Central Michigan University, Mt. Pleasant, MI. April.

Schock, N.T. and D.G. Uzarski. Stream/Drainage Ditch Impacts on Great Lakes Coastal Wetland Macroinvertebrate Community Composition. 55th International Conference on Great Lakes Research, Cornwall, Ontario.

Schock N.T., Uzarski D.G., 2013. Habitat conditions and macroinvertebrate communities of Great Lakes coastal habitats dominated by wet meadow, *Typha* spp. and *Phragmites australis*: implications of macrophyte structure changes. International Association for Great Lakes Research Conference, West Lafayette, IN. June.

Schock, N.T., B.A. Murry, D.G. Uzarski 2014. Impacts of agricultural drainage outlets on Great Lakes coastal wetlands. Great Lakes Science in Action Symposium, Central Michigan University, Mt. Pleasant, MI. April.

Schock, N.T., Schuberg, D.H., and Uzarski, D.G. 2015. Chemical and physical habitat gradients within Great Lakes coastal wetlands. 58th International Association for Great Lakes Research Conference, Burlington, VT. May.

Schoen, L.S., J.J. Student, and D.G. Uzarski. 2014. Reconstruction of fish movements between Great Lakes coastal wetlands. American Fisheries Society, Holland, MI. February.

- Sherman, J.S., T.A. Clement, N.T. Schock, and D.G. Uzarski. 2012. A comparison of abiotic and biotic parameters of diked and adjacent open wetland complexes of the Erie Marsh Preserve. 55th International Conference on Great Lakes Research, Cornwall, Ontario.
- Sherman, J.J., and D.G. Uzarski. 2013. A Comparison of Abiotic and Biotic Parameters of Diked and Adjacent Open Wetland Complexes of the Erie Marsh Preserve. 56th International Conference on Great Lakes Research, West Lafayette, IN. June.
- Sierszen, M., Schoen, L., Hoffman, J., Kosiara, J., and D. Uzarski. 2017. Support of coastal fishes by nearshore and coastal wetland habitats. 60th International Conference on Great Lakes Research, Detroit, MI. May. Presentation.
- Sierzen, M., L. Schoen, J. Hoffman, J. Kosiara and D. Uzarski. 2018. Tracing multi-habitat support of coastal fishes. Association for the Sciences of Limnology and Oceanography-Ocean Sciences Meeting. Portland, OR. February 2018. Oral Presentation.
- Smith, D.L., M.J. Cooper, J.M. Kosiara, and G.A. Lamberti. 2013. Heavy metal contamination in Lake Michigan wetland turtles. International Association for Great Lakes Research, 56th annual meeting. June 2-6, 2013. West Lafayette, IN. Poster presentation.
- Stirratt, H., M.J. Cooper. Landscape Conservation Design for the Great Lakes. International Union for the Conservation of Nature World Conservation Congress, September 6-10, 2016, Honolulu, Hawai'i.
- Thoennes, J., and N.P. Danz. 2017. Mapping Wetland Areal Change in the St. Louis River Estuary Using GIS. Poster presentation at the St. Louis River Summit, Superior, WI.
- Tozer, D.C., and S.A. Mackenzie. Control of invasive *Phragmites* increases breeding marsh birds but not frogs. Long Point World Biosphere Research and Conservation Conference, Simcoe, Ontario, Canada. Oral Presentation. 8 November 2019.
- Tozer, D.C., M. Falconer, A. Bracey, E. Giese, T. Gehring, G. Grabas, R. Howe, G. Niemi, and C. Norment. 2018. Detecting and monitoring elusive marsh breeding birds in the Great Lakes. IAGLR Conference, Toronto, Canada. Oral Presentation. 18-22 June 2018. (INVITED).
- Trebitz, A., J. Hoffman, G. Peterson, G. Shepard, A. Frankiewicz, B. Gilbertson, V. Brady, R. Hell, H. Wellard Kelly, and K. Schmude. 2015. The faucet snail (*Bithynia tentaculata*) invades the St. Louis River Estuary. St. Louis River Estuary Summit, Superior, Wisconsin. Mar. 30 – Apr. 1.
- Tuttle, E., T.N. Brown, D.A. Albert, and *T.J. Lemein. 2013. Comparison of two plant indices: Floristic Quality Index (FQI) and an index based on non-native and invasive species. Annual Society of Wetland Scientists Conference, Duluth, MN. June 4, 2013.

Unitis, M.J., B.A. Murry and D.G. Uzarski. 2012. Use of coastal wetland types by juvenile fishes. Ecology and Evolutionary Ecology of Fishes, Windsor, Ontario. June 17-21.

Uzarski, D.G. 2011. Great Lakes Coastal Wetland Monitoring for Restoration and Protection: A Basin-Wide Effort. State Of the Lakes Ecosystem Conference (SOLEC). Erie, Pennsylvania. October 26.

Uzarski, D.G. 2011. Coastal Wetland Monitoring: Background and Design. Great Lakes Coastal Wetland Monitoring Meeting. MDEQ; ASWM. Acme, Michigan. August 29.

Uzarski, D.G., N.T. Schock, T.A. Clement, J.J. Sherman, M.J. Cooper, and B.A. Murry. 2012. Changes in Lake Huron Coastal Wetland Health Measured Over a Ten Year Period During Exotic Species Invasion. 55th International Conference on Great Lakes Research, Cornwall, Ontario.

Uzarski, D.G., M.J. Cooper, V.J. Brady, J. Sherman, and D.A. Wilcox. 2013. Use of a basin-wide Great Lakes coastal wetland monitoring program to inform and evaluate protection and restoration efforts. International Association for Great Lakes Research, West Lafayette, IN. (INVITED)

Uzarski, D.G. 2013. A Basin Wide Great Lakes Coastal Wetland Monitoring Plan. Region 5 State and Tribal Wetlands Meeting: Focusing on Wetland Monitoring and Assessment around the Great Lakes. October 31. Kellogg Biological Station, Hickory Corners, MI.

Uzarski, D.G. 2013. Great Lakes Coastal Wetland Assessments. Lake Superior Cooperative Science and Monitoring Workshop. September 24-25. EPA Mid-Continent Ecology Division Lab, Duluth, MN.

Uzarski, D.G. 2013. A Basin-Wide Great Lakes Coastal Wetland Monitoring Program. 5th National Conference on Ecosystem Restoration. July 29-August 2. Schaumburg, IL.

Uzarski, D.G., Cooper, M.J., Brady, V., Sherman, J.J., and D.A. Wilcox. 2013. Use of a Basin Wide Great Lakes Coastal Wetland Monitoring Program to inform and Evaluate Protection and Restoration Efforts. 56th International Conference on Great Lakes Research, West Lafayette, IN.

Uzarski, D., M. Cooper and V. Brady. 2014. Implementing a Basin-wide Great Lakes Coastal Wetland Monitoring Program. Webinar for Sustain Our Great Lakes, Jan. 29, 2014. On-line webinar for Great Lakes researchers, managers, agency personnel, and environmental groups. Attendance approximately 400.

- Uzarski, D.G., Schock, N.T., Schuberg, D.H., Clement, T.A., and Cooper, M.J. 2015. Interpreting multiple organism-based IBIs and disturbance gradients: Basin wide monitoring. 58th International Conference on Great Lakes Research, Burlington, VT. May.
- Uzarski, D.G., N. Schock, T.M. Gehring, and B.A. Wheelock. 2016. Faucet snail (*Bithynia tentaculata*) occurrence across the Great lakes basin in coastal wetlands. 59th International Conference on Great Lakes Research, Guelph, Ontario Canada. May. Oral Presentation.
- Uzarski, D.G., V.J. Brady, M.J. Cooper, D.A. Wilcox, A.A. Bozimowski. 2017. Leveraging landscape level monitoring and assessment program for developing resilient shorelines throughout the Laurentian Great Lakes. Society of Wetland Scientists Annual Meeting. San Juan, Puerto Rico. June. Presentation.
- Uzarski, D.G., V.J. Brady, and M.J. Cooper. 2017. The Great Lakes Coastal Wetland Monitoring Program: Seven Years of Implementation. 60th International Conference on Great Lakes Research, Detroit, MI. May. Presentation.
- Uzarski, D.G. 2017. Emerging Issues in Wetland Science. Michigan Wetland Association Conference. Gaylord, Michigan. Plenary Presentation.
- Uzarski, D.G. 2018. Monitoring multiple biological attributes in Great Lakes coastal wetlands: database access for invasive species management. Association of State Wetlands Managers. Webinar Presentation.
- Uzarski, D.G. Global Significance & Major Threats to the Great Lakes. 2018. Frey Foundation Strategic Learning Session. The Great Lakes: Global Significance, Major Threats & Innovative Solutions. Petoskey, MI.
- Uzarski, D.G., V.J. Brady, M.J. Cooper, et al. 2018. The Laurentian Great Lakes Coastal Wetland Monitoring Program: Landscape level assessment of ecosystem health. ELLS-IAGLR Big Lakes Small World Conference. Évian, France. September. Poster
- Uzarski, D.G. and M.J. Cooper. 2019. Using a decision tree approach to inform protection and restoration of Great Lakes coastal wetlands. International Association for Great Lakes Research. Brockport, NY.
- Walton, N.G., E.E.G. Giese, R.W. Howe, G.J. Niemi, N.P. Danz, V.J. Brady, T.N. Brown, J.H. Ciborowski, J.P. Gathman, G.E. Host, L.B. Johnson, E.D. Reavie, and K.E. Kovalenko. 2013. How do different taxa respond to landscape stressors in Great Lakes coastal wetlands? 98th Annual Meeting of the Ecological Society of America. Minneapolis, MN, August 4-9.

Webster, W.C. and D.G. Uzarski. 2012. Impacts of Low Water level Induced Disturbance on Coastal Wetland Vegetation. 55th International Conference on Great Lakes Research, Cornwall, Ontario.

Wheeler, R. and D.G. Uzarski. 2012. Spatial Variation of Macroinvertebrate Communities within Two Emergent Plant Zones of Great Lakes Coastal Wetlands. 55th International Conference on Great Lakes Research, Cornwall, Ontario.

Wheeler, R.L. and D.G. Uzarski. 2013. Effects of Vegetation Zone Size on a Macroinvertebrate-based Index of Biotic Integrity for Great Lakes Coastal Wetlands. 56th International Conference on Great Lakes Research, West Lafayette, IN. June.

Wheelock, B.A., T.M. Gehring, D.G. Uzarski, G.J. Niemi, D.C. Tozer, R.W. Howe, and C.J. Norment. 2016. Factors affecting current distribution of Anurans in Great Lakes coastal wetlands. 59th International Conference on Great Lakes Research, Guelph, Ontario Canada. May. Oral Presentation.

Wilcox, D.A. 2018. Application of the Great Lakes Coastal Wetland Monitoring Program to restoration projects in Lake Ontario wetlands. Society of Wetland Scientists, Denver, CO.

Wilcox, D.A. 2018. Wetland restorations in the Braddock Bay Fish and Wildlife Management Area of Lake Ontario. Great Lakes Coastal Wetland Monitoring Program. Midland, MI. (INVITED)

Wilcox, D.A. and B.M. Mudrzynski. 2011. Wetland vegetation sampling protocols under the Great Lakes Coastal Wetland Monitoring program: experience in Lake Ontario. State of the Lakes Ecosystem Conference, Erie, PA. (INVITED)

Wilcox, D.A. and B.M. Mudrzynski. 2012. Implementing Great Lakes coastal wetlands monitoring: southern Lake Ontario. SUNY Great Lakes Research Consortium Conference, Oswego, NY. (INVITED)

Wilcox, D.A., B.M. Mudrzynski, D.G. Uzarski, V.J. Brady, M.J. Cooper, and T.N. Brown. 2016. Great Lakes coastal wetland monitoring program assesses wetland condition across the basin. Society of Wetland Scientists, Corpus Christi, TX.

Wilcox, D.A., B.M. Mudrzynski, D.G. Uzarski, V.J. Brady, and M.J. Cooper. 2017. A second phase of the Great Lakes Coastal Wetland Monitoring Program to assess wetland health across the basin. Society of Wetland Scientists, San Juan, PR.

Wilcox, D.A. 2012. Wetland restoration options under the Great Lakes Restoration Initiative. SUNY Great Lakes Research Consortium Conference, Oswego, NY. (INVITED)

Wilcox, D.A., D.G. Uzarski, V.J. Brady, M.J. Cooper, and T.N. Brown. 2013. Great Lakes coastal wetland monitoring program assists restoration efforts. Fifth World Conference on Ecological Restoration, Madison, WI.

Wilcox, D.A., D.G. Uzarski, V.J. Brady, M.J. Cooper, and T.N. Brown. 2014. Wetland restoration enhanced by Great Lakes coastal wetland monitoring program. Society of Wetland Scientists, Portland, OR.

Wilcox, D.A., D.G. Uzarski, V.J. Brady, and M.J. Cooper. 2019. Student training in wetland science through the Great Lakes Coastal Wetland Monitoring Program. Society of Wetland Scientists, Baltimore, MD.

Wilcox, D.A. 2015. Wetland restorations in the Braddock Bay Fish and Wildlife Management Area of Lake Ontario. NY Waterfowl and Wetland Collaborative Network, Oswego, NY. (INVITED)

Winter, C., T.K. O'Donnell, D.G. Uzarski, V.J. Brady, M.J., Cooper, A. Garwood, J.L. Utz, and J. Neal. 2017. Great Lakes coastal wetland monitoring: moving from assessment to action. Ecological Society of America Annual Conference. Portland, OR. Oral Presentation.

Wood, N.J., T.M. Gehring, and D.G. Uzarski. 2016. The invasive mute swan impacts on submerged aquatic vegetation in Michigan's coastal wetlands. 59th International Conference on Great Lakes Research, Guelph, Ontario Canada. May. Oral Presentation.

Zachritz, A.M, Miranda, D.A., Whitehead, H.D., Peaslee, G.F., Cressman, S. R., Lamberti, G.A. PFAS in Lake Michigan (USA) Salmonids: Implications for Human Dietary Intake. Joint Aquatic Sciences Meeting, Grand Rapids, MI. May 2022.

Zachritz, A.M., Miranda, D.A., Whitehead, H.D., Peaslee, G.F., Rand A.A., Harris K.J., Conard W.M., Cressman S.R., Lamberti, G.A. PFAS in Lake Michigan Salmonids: Ecological and Human Health Perspectives. Michigan PFAS Summit, virtual, December 2022.

PUBLICATIONS/MANUSCRIPTS (INCEPTION THROUGH 2023)

- Bansal, S., S. Lishawa, S. Newman, B. Tangen, D.A. Wilcox, D.A. Albert, M. Anteau, M. Chimney, R. Cressey, S. DeKeyser, K. Elgersma, S.A. Finkelstein, J. Freeland, R. Grosshans, P. Klug, D. Larkin, B. Lawrence, G. Linz, J. Marburger, G. Noe, C. Otto, N. Reo, J. Richards, C.J. Richardson, L. Rogers, A. Schrank, D. Svedarsky, S. Travis, N. Tuchman, A.G. van der Valk, and L. Windham-Myers. 2019. Typha (cattail) invasion in North American wetlands: Biology, regional problems, impacts, desired services, and management. *Wetlands* 39:645-684.
- Carson, D.B., S.C. Lishawa, N.C. Tuchman, A.M. Monks, B.A. Lawrence, and D.A. Albert. 2018. Harvesting invasive plants to reduce nutrient loads and produce bioenergy: an assessment of Great Lakes coastal wetlands. *Ecosphere* 9(6):e02320. 10.1002/ecs2.2320
- Ciborowski, J.J.H., J. Landry, L. Wang and J. Tomal. 2020. Compiling and Assessing Environmental Stress and Biological Condition Data for the Detroit River Area of Concern. Prepared for Environment and Climate Change Canada, Toronto, ON.
- Ciborowski, J.J.H., P. Chow Fraser, M. Croft, L. Wang, J. Buckley, J.P. Gathman, L.B. Johnson, S. Parker, D. Uzarski and M. Cooper. 2015. Lake Huron coastal wetland status - Review, assessment and synopsis of the condition of coastal wetlands and associated habitats. Technical report prepared for The Lake Huron Binational Partnership.
- Conard et al. 2022 Maternal Offloading of Per- and Polyfluoroalkyl Substances to Eggs by Lake Michigan Salmonids. *Environmental Science & Technology Letters*. (in revision) <https://doi.org/10.1021/acs.estlett.2c00627>
- Cooper, M.J., and D.G. Uzarski. 2016. Invertebrates in Great Lakes Marshes. *Invertebrates in Freshwater Marshes: An International Perspective on their Ecology*: D. Batzer (ed). Springer.
- Cooper, M.J., G.A. Lamberti, and D.G. Uzarski. 2014. Spatial and temporal trends in invertebrate communities of Great Lakes coastal wetlands, with emphasis on Saginaw Bay of Lake Huron. *Journal of Great Lakes Research Supplement* 40:168–182.
- Cooper, M.J., G.M. Costello, S.N. Francoeur, and G.A. Lamberti. 2016. Nitrogen limitation of algal biofilms in coastal wetlands of Lakes Michigan and Huron. *Freshwater Science* 35(1):25–40.
- Cooper, M.J., G.A. Lamberti, A.H. Moerke, C.R. Ruetz, D.A. Wilcox, V.J. Brady, T.N. Brown, J.J.H. Ciborowski, J.P. Gathman, G.P. Grabas, L.B. Johnson, and D.G. Uzarski. 2018. An expanded fish-based index of biotic integrity for Great Lakes coastal wetlands. *Environmental Monitoring and Assessment* 190: 580.

- Danz, N.P., N. Dahlberg, and S. Schooler. 2017. The St. Louis River Estuary vegetation database. Lake Superior Research Institute Technical Report 2017-1, University of Wisconsin-Superior, Superior, WI. 8 pages.
- Denomme-Brown, S.T., G.E. Fiorino, T. M. Gehring, G. J. Lawrence, D. C. Tozer, and G. P. Grabas. 2023. Marsh birds as ecological performance indicators for Lake Ontario outflow regulation. *Journal of Great Lakes Research*, in press.
- Des Jardin, K. 2015. Water chestnut: field observations, competition, and seed germination and viability in Lake Ontario coastal wetlands. M.S. Thesis. SUNY-The College at Brockport, Brockport, NY.
- Diller, S.N., A.M. Harrison, K.P. Kowalski, V.J. Brady, J.J.H. Ciborowski, M.J. Cooper, J.D. Dumke, J.P. Gathman, C.R. Ruetz III, D.G. Uzarski, D.A. Wilcox, J.S. Schaeffer. 2022. Influences of seasonality and habitat quality on Great Lakes coastal wetland fish community composition and diets. *Wetlands Ecology & Management*. DOI: 10.1007/s11273-022-09862-8
- Dumke, J., V. Brady, N. Danz, A. Bracey, G. Niemi. 2014. St. Louis River Report: Clough Island. NRRI TR2014/26 for Wisconsin DNR.
- Dumke, J.D., G.M. Chorak, C.R. Ruetz III, R.A. Thum, and J.N. Wesolek. 2020. Identification of Black Bullhead (*Ameiurus melas*) and Brown Bullhead (*A. nebulosus*) from the Western Great Lakes: Recommendations for Small Individuals. *The American Midland Naturalist* 183: 90-104.
- Dybiec, J.M., D.A. Albert, N.P. Danz, D.A. Wilcox, and D.G. Uzarski. 2020. Development of a preliminary vegetation-based indicator of ecosystem health for coastal wetlands of the Laurentian Great Lakes. *Ecological Indicators*. 119: 106768.
- Elliott, L.H., A.M. Bracey, G.J. Niemi, D.H. Johnson, T.M. Gehring, E.E. Gnass Giese, G.P. Grabas, R.W. Howe, C.J. Norment, D.C. Tozer, L.D. Igl LD. 2023. Application of habitat association models across regions: useful explanatory power retained in wetland bird case study. *Ecosphere*. In press.
- Gaul, W. 2017. Inferential measures for a quantitative ecological indicator of ecosystem health. M.Sci. Thesis, University of Wisconsin, Green Bay, Wisconsin. 35 pp.
- Gehring, T.M., C.R., Blass, B.A. Murry, and D.G. Uzarski. 2020. Great Lakes coastal wetlands as suitable habitat for invasive mute swans. *Journal of Great Lakes Research* 46:323-329.
- Gentine, J., W. Conard, K. O'Reilly, M. Cooper, G. Fiorino, A. Harrison, M. Hein, A. Moerke, C. Ruetz, D. Uzarski, and G. Lamberti. 2022. Environmental predictors of phytoplankton

chlorophyll-a in Great Lakes coastal wetlands. Journal of Great Lakes Research 48(4):927-934. <https://doi.org/10.1016/j.jglr.2022.04.015>

Gnass Giese, E.E., R.W. Howe, A.T. Wolf, N.A. Miller, and N.G. Walton. 2015. Sensitivity of breeding birds to the “human footprint” in western Great Lakes forest landscapes. Ecosphere 6(6):90. <http://dx.doi.org/10.1890/ES14-00414.1>

Gnass Giese, E.E., R.W. Howe, A.T. Wolf, and G.J. Niemi. 2018. Breeding birds and anurans of dynamic coastal wetlands in Green Bay, Lake Michigan. Journal of Great Lakes Research (Green Bay Special Issue): 44(5):950-959. <https://doi.org/10.1016/j.jglr.2018.06.003>

Grand, J., S.P. Saunders, N.L. Michel, L. Elliott, S. Beilke, A. Bracey, T.M. Gehring, E.R. Gnass Giese, R.W. Howe, B. Kasberg, N. Miller, G.J. Niemi, C.J. Norment, D.C. Tozer, J. Wu, and C. Wilsey. 2020. Prioritizing coastal wetlands for marsh bird conservation in the U. S. Great Lakes. Biological Conservation 249: 108708. <https://doi.org/10.1016/j.biocon.2020>

Harrison, A.M., A.J. Reisinger, M.J. Cooper, V.J. Brady, J.J. Ciborowski, K.E. O'Reilly, C.R. Ruetz, D.A. Wilcox, and D.G. Uzarski. 2020. A Basin-Wide Survey of Coastal Wetlands of the Laurentian Great Lakes: Development and Comparison of Water Quality Indices. Wetlands, 40:465-477. <https://doi.org/10.1007/s13157-019-01198>

Heminway, A.W. 2016. Response of *Typha x glauca* to phosphorus, hydrology, and land use in Lake Ontario coastal wetlands. M.S. Thesis. SUNY-The College at Brockport, Brockport, NY.

Heminway, A.W. and D.A. Wilcox. 2022. Response of *Typha* to phosphorus, hydrology, and land use in Lake Ontario coastal wetlands and a companion greenhouse study. Wetlands Ecology and Management 30:1167-1180.

Hilts, D.J., M.W. Belitz, T.M. Gehring, K.L. Pangle, and D.G. Uzarski. 2019. Climate change and nutria range expansion in the Eastern United States. Journal of Wildlife Management 83:591-598.

Hohman, T. 2019. Bird community response to change in wetland extent and lake level in Great Lakes coastal wetlands. M.Sci. Thesis, University of Wisconsin, Green Bay, Wisconsin. 41 pp.

Hohman, T.R., R.W. Howe, D.C. Tozer, E.E. Gnass Giese, A.T. Wolf, G.J. Niemi, T.M. Gehring, G.P. Grabas, and C.J. Norment. 2021. Influence of lake levels on water extent, interspersions, and marsh birds in Great Lakes coastal wetlands. Journal of Great Lakes Research 47(2):534-545. <https://doi.org/10.1016/j.jglr.2021.01.006>

Horton, D.J., K.R. Theis, D.G. Uzarski, D.R. Learman 2018. Microbial community structure and microbial networks correspond to nutrient gradients within coastal wetlands of the Great Lakes. bioRxiv, 217919

- Howe, R.W., E.E. Gnass Giese, and A.T. Wolf. 2018. Quantitative restoration targets for fish and wildlife habitats and populations in the Lower Green Bay and Fox River AOC. *Journal of Great Lakes Research* (Green Bay Special Issue): 44(5):883-894.
<https://doi.org/10.1016/j.jglr.2018.05.002>
- Howe, R.W., G.J. Niemi, L. Elliott, A.M. Bracey, W. Gaul, T.M. Gehring, E.E. Gnass Giese, G.P. Grabas, C.J. Normont, H. Panci, D. Tozer, and N.G. Walton. 2020. Birds as Indicators of Great Lakes Wetland Quality. In preparation for submission to *Ecological Indicators*.
- Kneisel, A.N., M.J. Cooper, A.K. Monfils, S. Haidar, and D.G. Uzarski. 2020. Ecological data as a resource for invasive species management in U.S. Great Lakes coastal wetlands. *Journal of Great Lakes Research*. 46 (4): 910-919.
- Kovalenko, K.E., L.B. Johnson, V.J. Brady, J.H.H. Ciborowski, M.J. Cooper, J.P. Gathman, G.A. Lamberti, A.H. Moerke, C.R. Ruetz III, and D.G. Uzarski. 2019. Hotspots and bright spots in functional and taxonomic fish diversity. *Freshwater Science*. 38:480-490.
doi.org/10.1086/704713
- Langer, T. A., B. A. Murry, K.L. Pangle, and D. G. Uzarski. 2016. Species turnover drives β -diversity patterns across multiple spatial and temporal scales in Great Lakes Coastal Wetland Communities. *Hydrobiologia*, DOI 10.1007/s10750-016-2762-2.
- Langer, T.A., M.J. Cooper, L.S. Reisinger, A.J. Reisinger, and D. G. Uzarski. 2017. Water depth and lake-wide water level fluctuation influence on α - and β -diversity of coastal wetland fish communities. *Journal of Great Lakes Research*, In Press. 44(1): 71-76.
- Lemein, T., D.A. Albert, and E.D. Tuttle. 2017. Coastal wetland vegetation community classification and distribution across environmental gradients through the Laurentian Great Lakes. *Journal of Great Lakes Research* 43 (4): 658-669.
- Lishawa, S.C., B.A. Lawrence, D.A. Albert, N.C. Tuchman. 2015. Biomass harvest of invasive *Typha* promotes plant diversity in a Great Lakes coastal wetland. *Restoration Ecology* Vol. 23 (3):228-237.
- Miranda, D. A., Peaslee, G. F., Zachritz, A. M., Lamberti, G. A. 2022. A worldwide evaluation of trophic magnification of per- and polyfluoroalkyl substances in aquatic ecosystems. *Integr. Environ. Assess. Manag.* 00, 1–13.doi.org/10.1002/ieam.4579
- Monks, A.M., S.C. Lishawa, K.C. Wellons, D.A. Albert, B. Mudrzynski, and D.A. Wilcox. 2019. European frogbit (*Hydrocharis morsus-ranae*) invasion facilitated by non-native cattails (*Typha*) in the Laurentian Great Lakes. *Journal of Great Lakes Research* 45:912-918.

- Monacelli, K. and D.A. Wilcox. 2021. Competition between two floating-leaved aquatic plants. *Aquatic Botany* 172:103390.
- O'Reilly, K.O., M.J. Cooper, P.S. Forsythe, C.J. Houghton, J.S. Shrovnal, J.J. Student, D.G. Uzarski, and G.A. Lamberti. 2023. Lakescape connectivity: Mobile fish consumers link Lake Michigan coastal wetland and nearshore food webs. *Ecosphere* 14 (2): e4333
doi.org/10.1002/ecs2.4333
- Panci, H., G.J. Niemi, R.R. Regal, D.C. Tozer, R.W. Howe, C.J. Norment, T.M. Gehring. 2017. Influence of local-and landscape-scale habitat on Sedge and Marsh Wren occurrence in Great Lakes coastal wetlands. *Wetlands: in press*.
- Podoliak, J.M. 2018. Amphibian and bird communities of Lake Ontario coastal wetlands: disturbance effects and monitoring efficiencies. M.S. Thesis. SUNY-Brockport, Brockport, NY.
- Reisinger, A.J., A.M. Harrison, M.J. Cooper, C.R. Ruetz, D.G. Uzarski, D.A. Wilcox. In Press. A basin-wide survey of coastal wetlands of the Laurentian Great Lakes: Development and comparison of water quality indices. *Wetlands*. Early Online August 5, 2019.
- Rutherford, R., Hartsock, J.A., and Danz, N.P. 2022. Physical and plant community changes at a Lake Michigan coastal marsh related to a two-meter increase in lake level. *Wetlands Ecology and Management* 30(3):547-560.
- Schock, N.T. A.J. Reisinger, L.S. Reisinger, M.J. Cooper, J.J.H. Cibrowski, T.M. Gehring, A. Moerke, D.G. Uzarski. 2019. Relationships between the distribution of the invasive faucet snail (*Bithynia tentaculata*) and environmental factors in Laurentian Great Lakes coastal wetlands. *Biological Invasions*. <https://doi.org/10.1007/s10530-019-02000-1>.
- Schoen, D. G. Uzarski. 2016. Reconstructing fish movements between coastal wetlands and nearshore habitats of the Great Lakes. *Limnology and Oceanography*, LO-15-0273.R1.
- Sierszen M.E., L.S. Schoen, J.M. Kosiara*, J.C. Hoffman, M.J. Cooper, and D.G. Uzarski. 2018. Relative contributions of nearshore and wetland habitats to coastal food webs in the Great Lakes. *J. Great Lakes Res*, <https://doi.org/10.1016/j.jglr.2018.11.006>
- Smith, D.L, M.J. Cooper, J.M. Kosiara, and G.A. Lamberti. 2016. Body burdens of heavy metals in Lake Michigan wetland turtles. *Environmental Monitoring and Assessment* 188:128.
- Suir, G.M. and D.A. Wilcox. 2021. Evaluating the use of hyperspectral imagery to

calculate raster-based wetland vegetation condition indicator. *Aquatic Ecosystem Health and Management* 24:100-114.

Suir, G.M., D.A. Wilcox, and M. Reif. 2021. Classification of *Typha*-dominated wetlands using airborne hyperspectral imagery along Lake Ontario, USA. *Aquatic Ecosystem Health and Management* 24:140-155.

Tozer, D.C., C.M. Falconer, A.M. Bracey, E.E. Gnass Giese, G.J. Niemi, R.W. Howe, T.M. Gehring, and C.J. Normont. 2017. Influence of call broadcast timing within point counts and survey duration on detection probability of marsh breeding birds. *Avian Conservation and Ecology* 12(2):8.

Tozer, D.C., R.W. Howe, G.J. Niemi, E.E. Gnass Giese, N.G. Walton, A.M. Bracey, W. Gaul, C.J. Normont, and T.M. Gehring. 2017. Coastal Wetland Amphibians in State of the Great Lakes 2017 Technical Report: Indicators to assess the status and trends of the Great Lakes ecosystem on pages 146-162.

Tozer, D.C., R.W. Howe, G.J. Niemi, E.E. Gnass Giese, N.G. Walton, A.M. Bracey, W. Gaul, C.J. Normont, and T.M. Gehring. 2017. Coastal Wetland Birds in State of the Great Lakes 2017 Technical Report: Indicators to assess the status and trends of the Great Lakes ecosystem on pages 163-179.

Tozer, D.C., and S.A. Mackenzie. Control of invasive *Phragmites* increases breeding marsh birds but not frogs. *Canadian Journal of Wildlife Management* 8:66-82.

Uzarski, D.G., V.J. Brady, M.J. Cooper, D.A. Wilcox, D.A. Albert, R. Axler, P. Bostwick, T.N. Brown, J.J.H. Ciborowski, N.P. Danz, J. Gathman, T. Gehring, G. Grabas, A. Garwood, R. Howe, L.B. Johnson, G.A. Lamberti, A. Moerke, B. Murry, G. Niemi, C.J. Normont, C.R. Ruetz III, A.D. Steinman, D. Tozer, R. Wheeler*, T.K. O'Donnell, and J.P. Schneider. 2017. Standardized measures of coastal wetland condition: implementation at the Laurentian Great Lakes basin-wide scale. *Wetlands*, DOI:10.1007/s13157-016-0835-7.

Uzarski, D.G., D.A. Wilcox, V.J. Brady, M.J. Cooper, D.A. Albert, J.J.H. Ciborowski, N.P. Danz, A. Garwood, J.P. Gathman, T.M. Gehring, G.P. Grabas, R.W. Howe, G.A. Lamberti, A.H. Moerke, G.J. Niemi, C.R. Ruetz, D.C. Tozer, and T.K. O'Donnell, ACCEPTED FOR PUBLICATION. Leveraging landscape level monitoring and assessment program for developing resilient shorelines throughout the Laurentian Great Lakes. *Wetlands*.

Wilcox, D.A., E.L. Polzer, A. Graham, R.K. Booth, and B. Mudrzynski. 2023. Fen development along the southern shore of Lake Ontario. *Journal of Great Lakes Research* 49: (in press).

REFERENCES

- Albert, D. A. 2008. Chapter 3: Vegetation Community Indicators, In: Great Lakes Coastal Wetland Monitoring Plan. Great Lakes Coastal Wetland Consortium, Great Lakes Commission. Ann Arbor, MI.
- Bailey, R. G.; Cushwa, C T. 1981. Ecoregions of North America (map). (FWS/OBS-81/29.) Washington, DC: U.S. Fish and Wildlife Service. 1:12,000,000.
- Bourdaghs, M., C.A. Johnston, and R.R. Regal. 2006. Properties and performance of the floristic quality index in Great Lakes coastal wetlands. *Wetlands*, 26:718-735.
- CEC, 1997, Ecoregions of North America, Commission for Environmental Cooperation Working Group (CEC) http://www.eoearth.org/article/Ecoregions_of_North_America_%28CEC%29
- Cooper, M. G.A. Lamberti, A.H. Moerke, C.R. Ruetz III, D.A. Wilcox, V. J. Brady, T.N. Brown, J.J.H. Ciborowski, J.P. Gathman, G.P. Grabas, L.B. Johnson, D.G. Uzarski. 2018. An Expanded Fish-Based Index of Biotic Integrity for Great Lakes Coastal Wetlands. *Env. Monit. Assess.* 190:580. DOI: <https://doi.org/10.1007/s10661-018-6950-6>.
- Crewe, T.L. and Timmermans, S.T.A. 2005. Assessing Biological Integrity of Great Lakes Coastal Wetlands Using Marsh Bird and Amphibian Communities. Bird Studies Canada, Port Rowan, Ontario. 89pp.
- Danz, N.P., G.J. Niemi, R. R. Regal, T. Hollenhorst, L. B. Johnson, J.M. Hanowski, R.P. Axler, J.J.H. Ciborowski, T. Hrabik, V.J. Brady, J.R. Kelly, J.A. Morrice, J.C. Brazner, R.W. Howe, C.A. Johnston and G.E. Host. 2007. Integrated Measures of Anthropogenic Stress in the U.S. Great Lakes Basin. *Environ Manage.* 39:631–647.
- Dybiec, J.M., D.A. Albert, N.P. Danz, D.A. Wilcox, and D.G. Uzarski. 2020. Development of a preliminary vegetation-based indicator of ecosystem health for coastal wetlands of the Laurentian Great Lakes. *Ecological Indicators* 119: 106768.
- Elias, J. E, R. Axler, and E. Ruzyski. 2008. Water quality monitoring protocol for inland lakes. Version 1.0. National Park Service, Great Lakes Inventory and Monitoring Network. Natural Resources Technical Report NPS/GLKN/NRTR—2008/109. National Park Service, Fort Collins, Colorado.
- Elliott, L.H. 2019. Habitat associations and conservation of wetland-obligate birds. Ph.D. Dissertation, University of Minnesota. 180 p.
- Elliott, L.H., A. Bracey, G.J. Niemi, D.J. Johnson, T.M. Gehring, E.E. Gnass Giese, G.R. Fiorino, R.W. Howe, G.J. Lawrence, C.J. Norment, D.C. Tozer, and L.D. Igl. 2022. Wetland bird case study for application of habitat association models across Great Lakes and Prairie Pothole regions. University of Minnesota Data Repository. <https://conservancy.umn.edu/handle/11299/250105>
- Farnsworth, G.L., K.H. Pollock, J.D. Nichols, T.R. Simons, J.E. Hines, and J.R. Sauer. 2002. A removal model for estimating detection probabilities from point-count surveys. *Auk* 119:414-425.
- Gaul, W. 2017. Inferential measures for a quantitative ecological indicator of ecosystem health. M.Sc. Thesis. University of Wisconsin-Green Bay, Green Bay, WI. 184 pp.

- Gnass Giese, E.E., R.W. Howe, A.T. Wolf, N.A. Miller, N.G. Walton. 2015. Sensitivity of breeding birds to the “human footprint” in western Great Lakes forest landscapes. *Ecosphere* 6: 90.
<http://dx.doi.org/10.1890/ES14-00414.1>.
- Howe, R.W., A. Bracey, L.H. Elliott, G.J. Niemi, T.M. Gehring, E.E. Gnass Giese, G. Grabas, G. Lawrence, C. Norment, and D. Tozer. 2021. A multi-species ecological indicator of Great Lakes coastal wetland quality based on breeding birds. In prep.
- Howe, R.W., R. R. Regal, J.M. Hanowski, G.J. Niemi, N.P. Danz, and C.R. Smith. 2007a. An index of ecological condition based on bird assemblages in Great Lakes coastal wetlands. *Journal of Great Lakes Research* 33 (Special Issue 3): 93-105.
- Howe, R.W., R. R. Regal, G.J. Niemi, N.P. Danz, J.M. Hanowski. 2007b. A probability-based indicator of ecological condition. *Ecological Indicators* 7:793-806.
- Jung, J.A., H.N. Rogers, and G.P. Grabas. 2020. Refinement of an index of ecological condition for marsh bird communities in lower Great Lakes coastal wetlands. *Ecological Indicators* 113:
<https://www.sciencedirect.com/science/article/abs/pii/S1470160X20300340?via%3Dihub>
- Karr, J.R., 1981. Assessment of biotic integrity using fish communities. *Fisheries* 6: 21–27.
- Lyons, J. 2012. Development and validation of two fish-based indices of biotic integrity for assessing perennial coolwater streams in Wisconsin, USA. *Ecological Indicators* 23: 402-412.
- Meyer, SW, JW Ingram, and GP Grabas. 2006. The marsh monitoring program: evaluating marsh bird survey protocol modifications to assess Lake Ontario coastal wetlands at a site-level. Technical Report Series 465. Canadian Wildlife Service, Ontario Region, Ontario.
- Morrice, J.A., N.P. Danz, R.R. Regal, J.R. Kelly, G.J. Niemi, E.D. Reavie, T. Hollenhorst, R.P. Axler, A.S. Trebitz, A.M. Cotter, and G.S. Peterson. 2008. Human influences on water quality in Great Lakes coastal wetlands. *Environmental Management* 41:347–357.
- Omernik, J.M. 1987. Ecoregions of the conterminous United States. Map (scale 1:7,500,000). *Annals of the Association of American Geographers* 77(1):118-125.
- Panci, H.G., Niemi, G.J., Regal, R.R., Tozer, D.C., Gehring, T.M., Howe, R.W. and Norment, C.J. 2017. Influence of Local, Landscape, and Regional Variables on Sedge and Marsh Wren Occurrence in Great Lakes Coastal Wetlands. *Wetlands*, 37(3): 447-459.
- R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL. <https://www.R-project.org/>.
- Reavie, E., R. Axler, G. Sgro, N. Danz, J. Kingston, A. Kireta, T. Brown, T. Hollenhorst and M. Ferguson. 2006. Diatom-base weighted-averaging models for Great Lakes coastal water quality: Relationships to watershed characteristics. *J. Great Lakes Research* 32:321–347.
- Schock, N.T., A.J. Reisinger, L.S. Reisinger, M.J. Cooper, J.J.H. Cibrowski, T.M. Gehring, A. Moerke, D.G. Uzarski.

2019. Relationships between the distribution of the invasive faucet snail (*Bithynia tentaculata*) and environmental factors in Laurentian Great Lakes coastal wetlands. *Biological Invasions*. <https://doi.org/10.1007/s10530-019-02000-1>. Early Online May 7, 2019.

Tozer, D.C., R.W. Howe, G.J. Niemi, E.E. Gnass Giese, N.G. Walton, A.M. Bracey, W. Gaul, C.J. Norment, and T.M. Gehring. 2015. Coastal Wetland Birds. In State of the Great Lakes 2017, Environmental Canada and U.S. Environmental Protection Agency, draft report. Tozer, D.C., 2016. *Marsh bird occupancy dynamics, trends, and conservation in the southern Great Lakes basin: 1996 to 2013*. J. Great Lakes Res. 42, in press.

Trebitz, A., G. Shepard, V. Brady, K. Schmude. 2015. The non-native faucet snail (*Bithynia tentaculata*) makes the leap to Lake Superior. J. Great Lakes Res. 41, 1197-1200.

United States Environmental Protection Agency. 2002. Methods for Evaluating Wetland Condition: Developing Metrics and Indexes of Biological Integrity. Office of Water, United States Environmental Protection Agency. Washington, DC. EPA-822-R-02-016.

Uzarski, D.G., V.J. Brady, M.J. Cooper, D.A. Wilcox, D.A. Albert, R.P. Axler, P. Bostwick, T.N. Brown, J.J. Ciborowski, N.P. Danz, J.P. Gathman, T.M. Gehring, G.P. Grabas, A. Garwood, R.W. Howe, L.B. Johnson, G.A. Lamberti, A.H. Moerke, B.A. Murry, G.J. Niemi, C.J. Norment, C.R. Ruetz III, A.D. Steinman, D.C. Tozer, R. Wheeler, T.K. O'Donnell, and J.P. Schneider. 2017. Standardized Measures of Coastal Wetland Condition: Implementation at a Laurentian Great Lakes Basin-Wide Scale. Wetlands 37: 15-32.

Uzarski, D.G., T.M. Burton, and J.J.H. Ciborowski. 2008. Chemical/Physical and Land Use/Cover Measurements, in Great Lakes Coastal Wetlands Monitoring Plan, T.M. Burton, et al. (editors), Great Lakes Coastal Wetland Consortium Final Report to Great Lakes Commission (GLC) and U.S. Environmental Protection Agency – Great Lakes National Program Office (EPA-GLNPO). www.glc.org/wetlands (March 2008).

APPENDIX

News articles about faucet snail detection in Great Lakes coastal wetlands.

1. <http://www.upnorthlive.com/news/story.aspx?id=1136758>
2. <http://www.wgmt.com/news/features/top-stories/stories/Snail-harmful-to-ducks-spreading-in-Great-Lakes-63666.shtml>
3. <http://fox17online.com/2014/12/16/gvsu-researchers-find-more-of-invasive-snail-species-in-lake-michigan/>
4. http://www.ourmidland.com/news/cmu-scientists-identify-spread-of-invasive-species/article_e9dc5876-00f4-59ff-8bcd-412007e079e8.html
5. <http://www.therepublic.com/view/story/4cde108b10b84af7b9d0cfcba603cf7a/MI--Invasive-Snails>
6. <http://media.cmich.edu/news/cmu-institute-for-great-lakes-research-scientists-identify-spread-of-invasive-species>
7. <http://www.veooz.com/news/qHv4acl.html>
8. <http://www.gvsu.edu/gvnow/index.htm?articleId=1E55A5C5-D717-BBE7-E79768C5213BB277>
9. http://hosted2.ap.org/OKDUR/99dded7a373f40a5aba743ca8e3d4951/Article_2014-12-16-MI--Invasive%20Snails/id-b185b9fd71ea4fa895aee0af983d7dbd
10. <http://whitehallmontague.wzzm13.com/news/environment/327493-my-town-waterfowl-killer-spreads-great-lakes-basin>
11. <http://www.timesunion.com/news/science/article/Snail-harmful-to-ducks-spreading-in-Great-Lakes-5959538.php>
12. <http://grandrapids-city.com/news/articles/gvsu-researchers-find-more-of-invasive-snail-species-in-lake-michigan>
13. <http://myinforms.com/en-us/a/8645879-gvsu-researchers-find-more-of-invasive-snail-species-in-lake-michigan/>
14. <http://usnew.net/invasive-snail-in-the-great-lakes-region.html>
15. http://www.cadillacnews.com/ap_story/?story_id=298696&issue=20141216&ap_cat=2
16. <http://theoryoflife.com/connect/researchers-track-invasive-9251724/>
17. <http://snewsi.com/id/1449258811>
18. <http://www.newswalk.info/muskegon-mich-new-scientists-say-742887.html>
19. http://www.petoskeynews.com/sports/outdoors/snail-harmful-to-ducks-spreading-in-great-lakes/article_b94f1110-9572-5d18-a5c7-66e9394a9b24.html
20. <http://www.chron.com/news/science/article/Snail-harmful-to-ducks-spreading-in-Great-Lakes-5959538.php>
21. <http://usa24.mobi/news/snail-harmful-to-ducks-spreading-in-great-lakes>
22. <http://www.wopular.com/snail-harmful-ducks-spreading-great-lakes>
23. <http://www.news.nom.co/snail-harmful-to-ducks-spreading-in-14203127-news/>
24. http://www.mlive.com/news/muskegon/index.ssf/2014/12/hard_to_kill_invasive_faucet_s.html
25. <http://wkar.org/post/researchers-eye-spread-invasive-faucet-snails>

26. <http://www.greenfieldreporter.com/view/story/4cde108b10b84af7b9d0cfcba603cf7a/MI--Invasive-Snails>
27. <http://www.natureworldnews.com/articles/11259/20141217/invasive-snails-killing-great-lake-birds.htm>
28. <http://www.wsbt.com/news/local/snail-harmful-to-ducks-spreading-in-great-lakes/30251286>
29. <http://www.wtkg.com/articles/wood-news-125494/invasive-and-deadly-snail-found-in-13073963>
30. <http://www.techtimes.com/articles/22378/20141218/invasive-snail-problem-in-great-lakes-difficult-to-deal-with-says-experts.htm>
31. <http://perfsience.com/content/214858-invasive-snails-kill-birds-great-lakes>
32. <http://www.hollandsentinel.com/article/20141216/NEWS/141219279>
33. <http://www.woodradio.com/articles/wood-news-125494/invasive-and-deadly-snail-found-in-13073963>
34. <http://www.full-timewhistle.com/science-27/great-lake-invasive-snails-kill-birds-265.html>
35. <http://www.islamabadglobe.com/invasive-deadly-snails-are-more-dangerous-than-we-thought-805.html>
36. <http://americanlivewire.com/2014-12-17-invasive-snail-species-attack-birds-great-lakes/>
37. <http://www.seattlepi.com/news/science/article/Snail-harmful-to-ducks-spreading-in-Great-Lakes-5959538.php>
38. <http://www.pendletontimespost.com/view/story/4cde108b10b84af7b9d0cfcba603cf7a/MI--Invasive-Snails/>
39. <http://www.wilx.com/home/headlines/Invasive-Snail-Spreading-in-Great-Lakes-285933261.html>
40. <http://www.watertowndailytimes.com/article/20150119/NEWS03/150118434>
41. <http://howardmeyerson.com/2015/01/15/scientists-invasive-snail-more-prevalent-than-thought-poses-grave-danger-to-waterfowl/>

Mock-up of press release produced by collaborating universities.

FOR IMMEDIATE RELEASE: December 9, 2014

CONTACT: June Kallestad, NRRI Public Relations Manager, 218-720-4300

USEPA-sponsored project greatly expands known locations of invasive snail

DULUTH, Minn. – Several federal agencies carefully track the spread of non-native species. This week scientists funded by the Great Lakes Restoration Initiative in partnership with USEPA's Great Lakes National Program Office greatly added to the list of known locations of faucet snails (*Bithynia tentaculata*) in the Great Lakes. The new locations show that the snails have invaded many more areas along the Great Lakes coastline than anyone realized.

The spread of these small European snails is bad news for water fowl: They are known to carry intestinal flukes that kill ducks and coots.

"We've been noting the presence of faucet snails since 2011 but didn't realize that they hadn't been officially reported from our study sites," explained Valerie Brady, NRRI aquatic ecologist who is collaborating with a team of researchers in collecting plant and animal data from Great Lakes coastal wetlands.

Research teams from 10 universities and Environment Canada have been sampling coastal wetlands all along the Great Lakes coast since 2011 and have found snails at up to a dozen sites per year [See map 1]. This compares to the current known locations shown on the [USGS website](#) [see map 2].

"Our project design will, over 5 years, take us to every major coastal wetland in the Great Lakes. These locations are shallow, mucky and full of plants, so we're slogging around, getting dirty, in places other people don't go. That could be why we found the snails in so many new locations," explained Bob Hell, NRRI's lead macroinvertebrate taxonomist. "Luckily, they're not hard to identify."

The small snail, 12 – 15 mm in height at full size, is brown to black in color with a distinctive whorl of concentric circles on the shell opening cover that looks like tree rings. The tiny size of young snails means they are easily transported and spread, and they are difficult to kill.

According to the Minnesota Department of Natural Resources, the faucet snail carries three intestinal trematodes that cause mortality in ducks and coots. When waterfowl consume the infected snails, the adult trematodes attack the internal organs, causing lesions and hemorrhage. Infected birds appear lethargic and have difficulty diving and flying before eventually dying.

Although the primary purpose of the project is to assess how Great Lakes coastal wetlands are faring, detecting invasives and their spread is one of the secondary benefits. The scientific team expects to

report soon on the spread of non-native fish, and has helped to locate and combat invasive aquatic plants.

“Humans are a global species that moves plants and animals around, even when we don’t mean to. We’re basically homogenizing the world, to the detriment of native species,” Brady added, underscoring the importance of knowing how to keep from spreading invasive species. Hell noted, “We have to make sure we all clean everything thoroughly before we move to another location.”

For more information on how to clean gear and boats to prevent invasive species spread, go to www.protectyourwaters.net.